

Disclaimer: The manuscript and its contents are confidential, intended for journal review purposes only, and not to be further disclosed.

URL: <http://hype-submit.aha-journals.org>

Title: ANGIOTENSIN II PROMOTES K^V7.4 CHANNELS DEGRADATION

THROUGH REDUCED INTERACTION WITH HSP90

Manuscript number: HYPE201811116DR2

Author(s): Vincenzo Barrese, St George's University of London

Jennifer Stott, St George's University of London

Hericka Figueiredo, St George's University of London

Aisah Aubdool, Barts and the London School of Medicine and Dentistry, Queen Mary University of London

Adrian Hobbs, Barts & The London School of Medicine, QMUL

Thomas Jepps, University of Copenhagen

Alistair McNeish, University of Reading

Iain Greenwood, St George's University of London

TITLE PAGE

ANGIOTENSIN II PROMOTES $K_v7.4$ CHANNELS DEGRADATION THROUGH REDUCED INTERACTION WITH HSP90

Vincenzo Barrese¹, Jennifer B Stott¹, Hericka B Figueiredo¹, Aisah A Aubdool², Adrian J
Hobbs², Thomas A Jepps³, Alister J McNeish⁴, Iain A Greenwood¹

¹Vascular Research Centre, Institute of Molecular & Clinical Sciences, St George's,
University of London, London, UK

²William Harvey Research Institute, Barts & The London School of Medicine, Queen Mary,
University of London, London, UK

³Department of Biomedical Sciences, University of Copenhagen, Copenhagen, Denmark

⁴Reading School of Pharmacy, University of Reading, Reading, UK

Short title: Angiotensin II induces $K_v7.4$ degradation

Manuscript word count: 5915

Abstract word count: 250

Number of figures: 6

Corresponding author: Vincenzo Barrese, MD, PhD

email: vbarrese@sgul.ac.uk. Tel: +44 (0) 2087252857

Vascular Research Centre, Institute of Molecular & Clinical
Sciences, St George's, University of London, Cranmer Terrace,
SW17 0RE, London, United Kingdom

ABSTRACT

Voltage gated Kv7.4 channels have been implicated in vascular smooth muscle cells (VSMCs) activity as they modulate basal arterial contractility, mediate responses to endogenous vasorelaxants, and are down-regulated in several arterial beds in different models of hypertension. Angiotensin II (Ang II) is a key player in hypertension that affects the expression of several classes of ion channels. In this study we evaluated the effects of Ang II on the expression and function of vascular Kv7.4. Western blot and quantitative PCR revealed that in whole rat mesenteric artery Ang II incubation for 1-7h decreased Kv7.4 protein expression without reducing transcript levels. Moreover, Ang II decreased XE991 (Kv7) – sensitive currents, and attenuated membrane potential hyperpolarization and relaxation induced by the Kv7 activator ML213. Ang II also reduced Kv7.4 staining at the plasma membrane of VSMCs. Proteasome inhibition with MG132 prevented Ang II-induced decrease of Kv7.4 levels, and counteracted the functional impairment of ML213-induced relaxation in myography experiments. Proximity Ligation Assays showed that Ang II impaired the interaction of Kv7.4 with the molecular chaperone HSP90, enhanced the interaction of Kv7.4 with the E3 ubiquitin ligase CHIP, and increased Kv7.4 ubiquitination. Similar alterations were found in mesenteric VSMCs isolated from Ang II-infused mice. The effect of Ang II was emulated by 17-AAG that inhibits HSP90 interactions with client proteins. These results show that Ang II downregulates Kv7.4 by altering protein stability through a decrease of its interaction with HSP90. This leads to the recruitment of CHIP and Kv7.4 ubiquitination and degradation via the proteasome.

Keywords: Angiotensin II/Heat shock protein/KCNQ/Kv7.4/Proteasome

INTRODUCTION

Hypertension is associated with increased arterial contractility and resistance to receptor-mediated vasodilators. K_v7 channels ($K_v7.1 - K_v7.5$) are voltage-gated potassium (K^+) channels encoded by *KCNQ* genes that regulate the contractile state of vascular smooth muscle at rest and contribute markedly to receptor-mediated vasorelaxations^{1, 2}. Arterial smooth muscle cells express $K_v7.1$, $K_v7.4$ and $K_v7.5$, with negligible $K_v7.2$ and $K_v7.3$ levels³, and blockers of these channels such as XE991 or linopirdine can contract most arteries or enhance vasoconstrictor responses⁴⁻⁶. Conversely, agents that enhance K_v7 activity are effective vasorelaxants⁴⁻⁷. In addition, K_v7 blockers, as well as knockdown of $K_v7.4$ impairs vasorelaxations produced by receptor agonists in various arteries^{2, 8, 9}. Notably, many receptor-mediated vasorelaxations are impaired in arteries from hypertensive animals, and any remaining relaxation is no longer sensitive to K_v7 blockade^{8, 10}. The functional impairment is linked to a reduction of $K_v7.4$ protein abundance in renal, mesenteric and coronary arteries from hypertensive animals, which is not correlated with a reduction in gene transcript (*KCNQ4*), suggesting the involvement of post transcriptional mechanisms. Whilst some molecular determinants of the pathological changes in K_v7 channel expression occurring in the vasculature during hypertension have started to be unveiled¹¹, the mechanisms that dictate the membrane abundance of $K_v7.4$ channels in smooth muscle are yet to be fully elucidated. K^+ channel trafficking can be regulated by several processes, such as the protein internalization/endosomal recycling pathway¹², and altered protein folding via changes in the interaction with molecular chaperones such as Heat Shock Proteins (HSPs)¹³. Here, we aim to uncover the processes governing the handling of $K_v7.4$ in vascular smooth muscle following treatment with Angiotensin II (Ang II), a molecule strongly associated with

cardiovascular dysfunction, to gain insight into the processes that control $K_v7.4$ membrane abundance during hypertension.

METHODS

Detailed methods are available in the Online Supplement (<http://hyper.ahajournals.org>). The authors declare that all supporting data are available within the article (and its online supplementary files).

Animals

All experiments were performed in accordance with the UK Animals (Scientific Procedures) Act (1986) and were approved by the local ethics committees (St George's Animal Welfare Committee, Barts and The London School of Medicine, University of Reading Animal Welfare and Ethical Review Board). 10-12 weeks-old male Wistar rats (Charles River, U.K.) and C57BL/6J mice (12-18 weeks of age) were killed by cervical dislocation. Animals were housed in a climatically controlled environment, on a 12h light/dark cycle, with free access to water and standard food ad libitum.

Statistical analysis

All data are expressed as mean \pm s.e.m. One- or two-way ANOVA test followed by a Dunnett's or Tukey's multiple comparisons test, and Student's t-test (paired or unpaired) were used to determine statistical significance between groups, according to the different experiments. Differences were considered statistically significant when $p < 0.05$.

RESULTS

Effects of Ang II on $K_v7.4$ levels and function

We first investigated the effects of Ang II on gross $K_v7.4$ levels in protein lysates from whole mesenteric arteries. Western blot experiments showed that 100nmol/L Ang II reduced $K_v7.4$ protein levels by ~50% after 1h compared to control vessels, by ~70% after 3h, and by ~80% after 7h treatment (Fig 1A). Incubation with Ang II did not reduce *KCNQ4* mRNA levels even after 7h (Fig 1B), suggesting a post-transcriptional mechanism. Similar results were also observed in rat aorta (Fig S1A-B, <http://hyper.ahajournals.org>). To circumvent any possible non-specific binding of the $K_v7.4$ antibody we investigated the effects of Ang II treatment on a human smooth muscle cell line (SGVSM-9) transfected with EGFP-tagged $K_v7.4$, using an anti-GFP antibody. In SGVSM-9 cells transfected with EGFP- $K_v7.4$, in-cell western blot experiments showed that treatment for 1h with 100nmol/L Ang II induced a ~20% decrease in EGFP- $K_v7.4$ with respect to control cells (Fig 1C). Moreover, in CHO cells expressing the angiotensin II receptor type 1 (AT1R) and EGFP- $K_v7.4$, incubation of 100nmol/L Ang II for 1h decreased EGFP- $K_v7.4$ protein expression by ~30%, as measured by Western blot (Fig S1C, <http://hyper.ahajournals.org>).

We then evaluated the effects of Ang II on K_v7 function. Whole cell K^+ currents were recorded from isolated mesenteric artery myocytes and the K_v7 current was identified as the component sensitive to the selective blocker XE991 (1 μ mol/L) (Fig S2A, <http://hyper.ahajournals.org>). Incubation of 100nmol/L Ang II for 3h decreased the amplitude of XE991-sensitive current by about 60% (Fig 1D). ML213 is an activator of $K_v7.2$, $K_v7.4$ and $K_v7.5$ channels that has been characterized in several smooth muscles, including VSMCs from mesenteric arteries¹⁴. In sharp microelectrode studies on whole mesenteric arteries ML213 hyperpolarized the membrane potential by approximately 17mV; a subsequent application of the K_{ATP} channel activator levcromakalin (10 μ mol/L) further increased membrane resting potential (Fig 1E and Fig S2B, <http://hyper.ahajournals.org>, control). Treatment with Ang II for 2h blunted ML213-induced hyperpolarization (increase of

membrane potential by $\sim 2\text{mV}$) but did not affect the membrane hyperpolarization caused by levcromakalin (Fig 1E and Fig S2B, <http://hyper.ahajournals.org>, Ang II). ML213-induced relaxation of pre-contracted mesenteric arteries is abolished by the K_v7 blockers XE991 (Fig S3, <http://hyper.ahajournals.org>) and linopirdine¹⁴. Incubation of mesenteric arteries with Ang II impaired the relaxation of pre-contracted mesenteric arteries produced by ML213 (Fig 1F and S2D, Fig S3, <http://hyper.ahajournals.org>), consistent with the loss of $K_v7.4$ function.

Effects on Ang II on $K_v7.4$ subcellular distribution

$K_v7.4$ subunits consistently show a clear and abundant peripheral distribution, following the shape of the cell membrane of the VSMC, whilst $K_v7.1$ was mainly intracellular and $K_v7.5$ signal was altogether less apparent and only occasionally localized at the plasma membrane (Fig S3, <http://hyper.ahajournals.org>). The $K_v7.4$ staining overlapped with the signal of the plasma membrane marker wheat germ agglutinin in control conditions (WGA, Fig. 2A). Treatment with 100nmol/L Ang II (3h) reduced the membrane localization of $K_v7.4$ concomitant with a decrease in the global level of $K_v7.4$ (Fig 2A). Fluorescence-intensity profiles along cross-sections of VSMCs showed that in control cells the $K_v7.4$ signal was higher in the membrane than cytosol (membrane:cytoplasm ratio 1.57 ± 0.12), while in Ang II treated cells $K_v7.4$ fluorescence was similar between the two regions (membrane:cytoplasm ratio 0.88 ± 0.10) (Fig 2B).

Role of endosomal and proteasomal pathways in Ang II-mediated $K_v7.4$ degradation

To gain insight about the cellular mechanisms that mediated the shift in $K_v7.4$ localization and reduction in total protein levels, mesenteric arteries were incubated with the endosome inhibitor dynasore or the proteasome inhibitor MG132 in presence or absence of Ang II. In Western blot experiments co-incubation of dynasore (DYN), used at concentrations

(100 μ mol/L) shown to be effective in previous studies¹⁵, did not fully prevent the reduction of K_v7.4 produced by 3h incubation with Ang II (Fig. 2C). Conversely, MG132 (20 μ mol/L) fully prevented the Ang II-induced reduction of K_v7.4 (Fig. 2C). MG132 also counteracted the impairment of ML213-induced relaxation of mesenteric arteries by Ang II in myography experiments (Fig 2D).

Role of oxidative stress in Ang II-induced K_v7.4 degradation

Ang II is known to increase oxidative stress¹⁶ which induces proteasome-mediated protein degradation¹⁷. We therefore evaluated whether the observed increase of K_v7.4 protein degradation induced by Ang II was mediated by oxidative stress. In VSMCs, treatment with Ang II for 30min increased the production of reactive oxygen species (ROS) measured by the fluorescence dye ROS Deep Red (by ~50% compared to time-matched control), similarly to that caused by 1mmol/L H₂O₂ (increase by ~70%, Fig 3A). In contrast, longer incubation (3h) with Ang II did not significantly enhance ROS levels with respect to time-matched control, whereas H₂O₂ was still able to increase ROS levels (by ~40%, (Fig 3A). The ROS scavenger N-Acetyl-L-Cysteine (NAC, 1mmol/L) counteracted Ang II-induced ROS production (Fig S5, <http://hyper.ahajournals.org>), but did not prevent either K_v7.4 protein down-regulation (Fig 3B) or the impairment of ML213-induced relaxation in mesenteric arteries when co-incubated with Ang II (Fig 3C).

Role of HSPs in Ang II-mediated K_v7.4 degradation

HSP70/90 machinery is abundantly expressed in eukaryotic cells, and is a key regulator of protein homeostasis in both physiologic and pathologic conditions¹⁸. Moreover, HSP70/90 interact with K_v7.4 in heterologous-expression systems¹⁹. Neither HSP70 nor HSP90 transcript/protein levels changed in Ang II-treated mesenteric arteries compared to control

vessels (Fig S6, <http://hyper.ahajournals.org>). Therefore, we investigated whether Ang II-induced downregulation of K_v7.4 was mediated by an altered interaction with HSPs using Proximity Ligation Assay (PLA). No significant differences in the number of interactions between K_v7.4 and HSP70 were observed in isolated VSMCs treated with Ang II when compared to controls (Fig 4A). In contrast, a significant reduction (by ~40%) in K_v7.4:HSP90 interactions was detected in VSMCs upon incubation with Ang II (Fig 4B). HSP70/90 machinery is linked to the proteasome system via the co-chaperone and E3 ubiquitin-ligase CHIP (C-terminus of Hsp70-Interacting Protein). Incubation of VSMCs with Ang II enhanced the interaction of K_v7.4 with CHIP by ~30% (Fig 4C), and increased the number of ubiquitin molecules interacting with K_v7.4 by ~65% when compared to control (Fig 4D). These data suggested that Ang II reduced K_v7.4:HSP90 interactions thus enhancing protein ubiquitination via an increased interaction of K_v7.4:HSP70 complex with CHIP. To corroborate this, we studied the effect of 17-AAG (17-Demethoxy-17-(2-propenylamino) geldanamycin), a geldanamycin-analog which inhibits HSP90 function, on total K_v7.4 protein levels. Incubation of mesenteric arteries with 1 μmol/L 17-AAG caused a reduction of K_v7.4 protein by ~60% (Fig 5A), increased the number of interactions between K_v7.4 and CHIP (Fig 5B), and the number of ubiquitin molecules interacting with K_v7.4 (Fig 5C).

Role of HSPs-CHIP-ubiquitin system in K_v7.4 degradation in vivo

Since K_v7.4 protein is down-regulated in Ang II-infused mice⁷, we investigated whether the observed alterations of HSPs-CHIP-ubiquitin system could be detected in the same *in vivo* model of prolonged exposure to Ang II. PLAs showed that the number of interactions of K_v7.4 with HSP70 in VSMCs isolated from mesenteric arteries of Ang II-infused mice was not significantly different from those isolated from saline-infused mice (Fig 6A), whereas a significant decrease in the number of K_v7.4:HSP90 interactions was observed in VSMCs

from Ang II- with respect to saline-infused mice (Fig 6B). Moreover, an increased interaction of K_v7.4 with CHIP (Fig 6C) and ubiquitin (Fig 6D) was detected in VSMCs from Ang II- mice when compared to control groups.

DISCUSSION

Ang II reduces K_v7 expression and function via increased proteasome-mediated degradation

In this study, we present novel and detailed mechanistic information showing how Ang II, a key mediator of the pathological changes occurring in vascular disease¹⁶, modulates K_v7.4 channel expression, which is down-regulated in hypertension⁷. Ang II has been demonstrated to acutely (up to 25 min) suppress the XE991-sensitive K_v current in mesenteric VSMCs²⁰, but no insight regarding the molecular mechanisms involved was provided, and the effects of longer exposure of Ang II on K_v7 expression and function were not studied. Our data show that treatment with Ang II (1-7h) reduced the expression of K_v7.4 protein, decreased the amplitude of voltage-dependent currents sensitive to the pan-K_v7 channels blocker XE991 in VSMCs, blunted the hyperpolarization of the VSMCs membrane induced by the K_v7 activator ML213, and impaired the vasorelaxant effects induced by the same drug in whole mesenteric arteries. In contrast, Ang II did not affect the ability of the K_{ATP} channel activator levcromakalin to hyperpolarize VSMCs. ML213 effects in smooth muscle are abolished by K_v7 blockers like XE991 (Fig S3, <http://hyper.ahajournals.org>) or linopirdine¹⁴, strongly suggesting the involvement of K_v7 channels in ML213-induced arterial relaxation. ML213 has been reported to be more specific for channels formed by K_v7.2 (whose expression in VSMCs is negligible³) or K_v7.4 subunits²¹, but one study suggested that it also enhances the current mediated by K_v7.5²², which contributes with K_v7.4 to the formation of the mature channel in the vasculature^{1, 23}. Due to the lack of K_v7 subunit-specific blockers, we could not rule out the involvement of K_v7.5 in ML213-induced relaxation. However, our

immunofluorescence data (Fig S4, <http://hyper.ahajournals.org>) together with previous evidence^{2, 8, 9, 23} infer that the reduction of vasorelaxant effects of ML213 prompted by Ang II was mediated by decreased $K_v7.4$ levels in VSMCs. Interestingly, ML213 still produced some relaxation in Ang II-treated arteries even though its ability to hyperpolarize was largely (but not entirely) lost by incubation with Ang II. A previous study revealed that 1 μ M ML213 produced considerable relaxation of pre-contracted arteries without marked membrane hyperpolarization¹⁴. This apparent discrepancy in electro-mechanical coupling probably reflects the differences in experimental conditions, as isometric tension studies involved a pre-contracted and therefore depolarized artery compared to the microelectrode recordings performed on unstimulated arteries. Like other $K_v7.2-7.5$ activators ML213 works through stabilization of the open configuration that will be more apparent under depolarized conditions. However, further investigation regarding the role of K_v7 channels in regulating the electro-mechanical coupling in VSMCs is required.

$K_v7.4$ down-regulation was also confirmed in clonal cells expressing heterologous $K_v7.4$ channels, and was not accompanied by a reduction in mRNA levels, indicating that Ang II reduced $K_v7.4$ by post-transcriptional mechanisms. Modulation of protein trafficking, recycling and degradation has a considerable impact on K^+ channels expression and function²⁴ and dysregulation of recycling/degradation pathways has been linked to channelopathies such as hereditary arrhythmias or cystic fibrosis²⁴. Our findings reveal that degradation of $K_v7.4$ protein by Ang II treatment and the subsequent functional impairment of K_v7 -dependant vasorelaxation was prevented by proteasome inhibition. Ang II has been shown to induce proteasome degradation of large conductance calcium activated BK channels via increased endosome-mediated internalization in VSMCs, and $K_v11.1$ (ERG) channels in over-expression systems^{15, 25}. Interestingly, in contrast to the above studies where Ang II effects were noticeable after prolonged treatment (7-24h), we observed

downregulation and altered subcellular distribution of K_v7.4 after as little as 1h. Moreover, the lack of effect of dynasore in fully restoring K_v7.4 protein levels suggest that the endosomal pathway does not play a primary role in Ang II-induced K_v7.4 down-regulation in contrast to BK_{Ca} channels¹⁵. Previous studies have established that Ang II causes oxidative stress in target cells¹⁶, and that the increased ROS levels activate the ubiquitin-proteasome system, as a protective mechanism to prevent the accumulation/aggregation of oxidized proteins¹⁷. In our experiments, Ang II caused a rapid increase in ROS levels (30min) which did not persist with longer exposure (3h), and the ROS scavenger N-Acetyl-L-Cysteine did not prevent the down-regulation of K_v7.4 levels and function prompted by Ang II. Although we cannot exclude the possibility that sustained oxidative stress reduces K_v7.4 levels, our data suggest that the early burst of ROS production induced by Ang II was not sufficient to alter K_v7.4 abundance, and that the activation of the proteasome system triggered by oxidative stress was not the main mechanisms involved in Ang II-induced downregulation of K_v7.4.

Ang II-induced decrease of K_v7.4:HSP90 interaction promotes the ubiquitination of K_v7.4

Ang II affects protein folding in hypertensive mice²⁶, regulates the activity of HSPs, which mediate some maladaptive responses occurring in hypertension²⁷, and also regulate ion channel folding. HSP90 in particular, plays a pivotal role in the folding of the cystic fibrosis transmembrane conductance regulator (CFTR), the ClC-2 chloride channel²⁸, HERG channel¹³ and the K_{ATP} potassium channel²⁹. Moreover, enhancement of the HERG:HSP90 interaction is responsible for the increased membrane trafficking induced by estradiol³⁰. Reduced binding of HSP90 decreases the interaction of client proteins with the HSP40-HSP70-HOP chaperone pathway that promotes forward trafficking, and increases interaction with HSP70-ubiquitin ligase CHIP complexes, which lead to proteasome-mediated degradation³¹. Our data reveal that the molecular chaperones HSP70/90 interact with K_v7.4 in

native VSMCs, and that Ang II treatment selectively reduced the interaction of K_v7.4 with HSP90 in this physiological environment. In heterologous expression systems, modulating HSP90 isoforms levels either by overexpression or silencing strategies revealed a differential role in K_v7.4 maturation of the two major HSP90 isoforms, namely α and β , with β -isoform promoting protein folding, and α -isoform enhancing CHIP-mediated proteasomal degradation³². Although some distinctive features exist, HSP90 isoforms exert similar actions, with many functions being characteristic of both isoforms and only associated more frequently to one of them³³. Moreover, their expression is differentially regulated, with a constitutive form (HSP90 β) and an inducible one (HSP90 α). Although we did not investigate the contribution of α - or β -isoform, in our experiments we did not observe any change in HSP90 levels, suggesting that HSP90 expression was not modified by Ang II incubation, and that a different mechanism was responsible for the alteration in K_v7.4:HSP90 binding. Covalent modifications, including phosphorylation, acetylation and nitrosylation, influence the activity of HSP90 and consequently the maturation of selected clients. Phosphorylation and acetylation slow down HSP90 activity, resulting in a reduced maturation of client proteins^{34, 35}. Therefore, Ang II might activate different pathways ultimately altering HSP90 and/or K_v7.4, modifying their ability to interact. Irrespective of the molecular mechanism, the ultimate effect was an increased interaction of K_v7.4 with CHIP and an enhanced ubiquitination of K_v7.4 as determined by PLA. This hypothesis was corroborated by studies with 17-AAG, a pharmacological inhibitor of HSP90, that in our experiments reproduced the effects of Ang II in terms of K_v7.4 down-regulation, interaction with CHIP and ubiquitination. Our results are in line with previous evidence showing that pharmacological inhibition of HSP90 induces rapid degradation of client proteins through the ubiquitin-proteasome pathway^{13, 36}.

PERSPECTIVES

We show for the first time in vascular smooth muscle that Ang II enhances K_v7.4 degradation via the ubiquitin-proteasome system, through a reduced interaction with HSP90. The specific pathway(s) activated by Ang II, as well as the possible molecular modifications occurring in all the interactors (including additional co-chaperones) will need to be further investigated. Indeed, Ang II can activate diverse intracellular pathways, including non G-protein signaling, and modulate several processes, such as inflammation¹⁶, which might in turn regulate HSPs activity and therefore protein stability and degradation. However, the present study has identified a novel and dynamic regulation of K_v7.4 that impacts considerably on vascular physiology. Noticeably, we showed that these alterations also occur *in vivo* after chronic exposure to Ang II, a well-known model of hypertension where K_v7.4 expression and function is reduced⁷, suggesting that the observed reduction of K_v7.4 protein stability and enhanced degradation play a role also in the pathological changes occurring in hypertension. Since modulators of HSP90 function are being developed therapeutically³⁷⁻³⁹, these data provide a possible new therapeutic strategy for the treatment of hypertension.

ACKNOWLEDGEMENTS

We thank Prof. Guy Whitley (St George's, University of London) for providing SGVSM-9 cells, and Prof. Riccardo Olcese (University of California Los Angeles) for sharing pcDNA-AT1R plasmid.

SOURCES OF FUNDING

This work has been funded by the British Heart Foundation for JS (PG/15/97/31862), AM (FS/14/33/30799) and AA (RG/16/7/32357), and by the Medical Research Council (MR/K019074/1) for VB. TAJ was funded by a Carlsberg Foundation Fellowship (CF16-0136).

DISCLOSURE

None

For Hypertension Peer Review. Do not distribute.
Destroy after use.

REFERENCES

1. Haick JM, Byron KL. Novel treatment strategies for smooth muscle disorders: Targeting kv7 potassium channels. *Pharmacol Ther.* 2016;165:14-25
2. Stott JB, Barrese V, Greenwood IA. Kv7 channel activation underpins epac-dependent relaxations of rat arteries. *Arterioscler Thromb Vasc Biol.* 2016;36:2404-2411
3. Barrese V, Stott JB, Greenwood IA. Kcnq-encoded potassium channels as therapeutic targets. *Annu Rev Pharmacol Toxicol.* 2018;58:625-648
4. Yeung SY, Pucovsky V, Moffatt JD, Saldanha L, Schwake M, Ohya S, Greenwood IA. Molecular expression and pharmacological identification of a role for k(v)7 channels in murine vascular reactivity. *Br J Pharmacol.* 2007;151:758-770
5. Mackie AR, Brueggemann LI, Henderson KK, Shiels AJ, Cribbs LL, Scrogin KE, Byron KL. Vascular kcnq potassium channels as novel targets for the control of mesenteric artery constriction by vasopressin, based on studies in single cells, pressurized arteries, and in vivo measurements of mesenteric vascular resistance. *J Pharmacol Exp Ther.* 2008;325:475-483
6. Zhong XZ, Harhun MI, Olesen SP, Ohya S, Moffatt JD, Cole WC, Greenwood IA. Participation of kcnq (kv7) potassium channels in myogenic control of cerebral arterial diameter. *J Physiol.* 2010;588:3277-3293
7. Jepps TA, Chadha PS, Davis AJ, Harhun MI, Cockerill GW, Olesen SP, Hansen RS, Greenwood IA. Downregulation of kv7.4 channel activity in primary and secondary hypertension. *Circulation.* 2011;124:602-611
8. Chadha PS, Zunke F, Zhu HL, Davis AJ, Jepps TA, Olesen SP, Cole WC, Moffatt JD, Greenwood IA. Reduced kcnq4-encoded voltage-dependent potassium channel

- activity underlies impaired beta-adrenoceptor-mediated relaxation of renal arteries in hypertension. *Hypertension*. 2012;59:877-884
9. Lee S, Yang Y, Tanner MA, Li M, Hill MA. Heterogeneity in kv7 channel function in the cerebral and coronary circulation. *Microcirculation*. 2015;22:109-121
 10. Mani BK, O'Dowd J, Kumar L, Brueggemann LI, Ross M, Byron KL. Vascular kcnq (kv7) potassium channels as common signaling intermediates and therapeutic targets in cerebral vasospasm. *J Cardiovasc Pharmacol*. 2013;61:51-62
 11. Carr G, Barrese V, Stott JB, Povstyan OV, Jepps TA, Figueiredo HB, Zheng D, Jamshidi Y, Greenwood IA. MicroRNA-153 targeting of kcnq4 contributes to vascular dysfunction in hypertension. *Cardiovasc Res*. 2016;112:581–589
 12. Seeböhm G, Strutz-Seeböhm N, Ureche ON, Henrion U, Baltaev R, Mack AF, Korniychuk G, Steinke K, Tapken D, Pfeufer A, Kaab S, Bucci C, Attali B, Merot J, Tavaré JM, Hoppe UC, Sanguinetti MC, Lang F. Long qt syndrome-associated mutations in kcnq1 and kcne1 subunits disrupt normal endosomal recycling of iks channels. *Circ Res*. 2008;103:1451-1457
 13. Ficker E, Dennis AT, Wang L, Brown AM. Role of the cytosolic chaperones hsp70 and hsp90 in maturation of the cardiac potassium channel hERG. *Circ Res*. 2003;92:e87-100
 14. Jepps TA, Bentzen BH, Stott JB, Povstyan OV, Sivaloganathan K, Dalby-Brown W, Greenwood IA. Vasorelaxant effects of novel kv 7.4 channel enhancers ml213 and ns15370. *Br J Pharmacol*. 2014;171:4413-4424
 15. Leo MD, Bulley S, Bannister JP, Kuruvilla KP, Narayanan D, Jaggar JH. Angiotensin II stimulates internalization and degradation of arterial myocyte plasma membrane BK channels to induce vasoconstriction. *Am J Physiol Cell Physiol*. 2015;309:C392-402

16. Karnik SS, Unal H, Kemp JR, Tirupula KC, Eguchi S, Vanderheyden PM, Thomas WG. International union of basic and clinical pharmacology. Xcix. Angiotensin receptors: Interpreters of pathophysiological angiotensinergic stimuli [corrected]. *Pharmacol Rev.* 2015;67:754-819
17. Demasi M, Laurindo FR. Physiological and pathological role of the ubiquitin-proteasome system in the vascular smooth muscle cell. *Cardiovasc Res.* 2012;95:183-193
18. Schopf FH, Biebl MM, Buchner J. The hsp90 chaperone machinery. *Nat Rev Mol Cell Biol.* 2017;18:345-360
19. Gao Y, Yechikov S, Vazquez AE, Chen D, Nie L. Distinct roles of molecular chaperones hsp90alpha and hsp90beta in the biogenesis of kcnq4 channels. *PLoS One.* 2013;8:e57282
20. Schleifenbaum J, Kassmann M, Szijarto IA, Hercule HC, Tano JY, Weinert S, Heidenreich M, Pathan AR, Anistan YM, Alenina N, Rusch NJ, Bader M, Jentsch TJ, Gollasch M. Stretch-activation of angiotensin ii type 1a receptors contributes to the myogenic response of mouse mesenteric and renal arteries. *Circ Res.* 2014;115:263-272
21. Yu H, Wu M, Townsend SD, Zou B, Long S, Daniels JS, McManus OB, Li M, Lindsley CW, Hopkins CR. Discovery, synthesis, and structure activity relationship of a series of n-aryl- bicyclo[2.2.1]heptane-2-carboxamides: Characterization of ml213 as a novel kcnq2 and kcnq4 potassium channel opener. *ACS Chem Neurosci.* 2011;2:572-577
22. Brueggemann LI, Haick JM, Cribbs LL, Byron KL. Differential activation of vascular smooth muscle kv7.4, kv7.5, and kv7.4/7.5 channels by ml213 and ica-069673. *Mol Pharmacol.* 2014;86:330-341

23. Chadha PS, Jepps TA, Carr G, Stott JB, Zhu HL, Cole WC, Greenwood IA. Contribution of kv7.4/kv7.5 heteromers to intrinsic and calcitonin gene-related peptide-induced cerebral reactivity. *Arterioscler Thromb Vasc Biol.* 2014;34:887-893
24. Curran J, Mohler PJ. Alternative paradigms for ion channelopathies: Disorders of ion channel membrane trafficking and posttranslational modification. *Annu Rev Physiol.* 2015;77:505-524
25. Cai Y, Wang Y, Xu J, Zuo X, Xu Y. Down-regulation of ether-a-go-go-related gene potassium channel protein through sustained stimulation of at1 receptor by angiotensin ii. *Biochem Biophys Res Commun.* 2014;452:852-857
26. Kassan M, Galan M, Partyka M, Saifudeen Z, Henrion D, Trebak M, Matrougui K. Endoplasmic reticulum stress is involved in cardiac damage and vascular endothelial dysfunction in hypertensive mice. *Arterioscler Thromb Vasc Biol.* 2012;32:1652-1661
27. Meier M, King GL, Clermont A, Perez A, Hayashi M, Feener EP. Angiotensin at(1) receptor stimulates heat shock protein 27 phosphorylation in vitro and in vivo. *Hypertension.* 2001;38:1260-1265
28. Hinzpeter A, Lipecka J, Brouillard F, Baudoin-Legros M, Dadlez M, Edelman A, Fritsch J. Association between hsp90 and the clc-2 chloride channel upregulates channel function. *Am J Physiol Cell Physiol.* 2006;290:C45-56
29. Yan FF, Pratt EB, Chen PC, Wang F, Skach WR, David LL, Shyng SL. Role of hsp90 in biogenesis of the beta-cell atp-sensitive potassium channel complex. *Mol Biol Cell.* 2010;21:1945-1954
30. Anneken L, Baumann S, Vigneault P, Biliczki P, Friedrich C, Xiao L, Girmatsion Z, Takac I, Brandes RP, Kissler S, Wiegratz I, Zumhagen S, Stallmeyer B, Hohnloser SH, Klingenheben T, Schulze-Bahr E, Nattel S, Ehrlich JR. Estradiol regulates human

- qt-interval: Acceleration of cardiac repolarization by enhanced *kcnh2* membrane trafficking. *Eur Heart J*. 2016;37:640-650
31. Kundrat L, Regan L. Balance between folding and degradation for hsp90-dependent client proteins: A key role for chip. *Biochemistry*. 2010;49:7428-7438
 32. Gao Y, Yechikov S, Vazquez AE, Chen D, Nie L. Impaired surface expression and conductance of the *kcnq4* channel lead to sensorineural hearing loss. *J Cell Mol Med*. 2013;17:889-900
 33. Sreedhar AS, Kalmar E, Csermely P, Shen YF. Hsp90 isoforms: Functions, expression and clinical importance. *FEBS Lett*. 2004;562:11-15
 34. Wandinger SK, Suhre MH, Wegele H, Buchner J. The phosphatase *ppt1* is a dedicated regulator of the molecular chaperone hsp90. *EMBO J*. 2006;25:367-376
 35. Zhao YG, Gilmore R, Leone G, Coffey MC, Weber B, Lee PW. Hsp90 phosphorylation is linked to its chaperoning function. Assembly of the reovirus cell attachment protein. *J Biol Chem*. 2001;276:32822-32827
 36. Isaacs JS, Jung YJ, Mimnaugh EG, Martinez A, Cuttitta F, Neckers LM. Hsp90 regulates a von hippel lindau-independent hypoxia-inducible factor-1 alpha-degradative pathway. *J Biol Chem*. 2002;277:29936-29944
 37. Guo J, Zhang X, Hu Z, Zhuang Z, Zhu Z, Chen Z, Chen W, Zhao Z, Zhang C, Zhang Z. A422t mutation in *herg* potassium channel retained in er is rescuable by pharmacologic or molecular chaperones. *Biochem Biophys Res Commun*. 2012;422:305-310
 38. Pratt WB, Gestwicki JE, Osawa Y, Lieberman AP. Targeting hsp90/hsp70-based protein quality control for treatment of adult onset neurodegenerative diseases. *Annu Rev Pharmacol Toxicol*. 2015;55:353-371

39. Yang B, Zhao D, Verkman AS. Hsp90 inhibitor partially corrects nephrogenic diabetes insipidus in a conditional knock-in mouse model of aquaporin-2 mutation. *FASEB J.* 2009;23:503-512

For Hypertension Peer Review. Do not distribute.
Destroy after use.

NOVELTY AND SIGNIFICANCE

What is new. We showed that Ang II decreases the expression and function of Kv7.4 potassium channels in vascular smooth muscle cells by reducing its interaction with chaperone protein HSP90 and altering protein stability.

What is relevant. Kv7.4 is an important mediator of arterial relaxation and is down-regulated in hypertension, a condition where Ang II is dysregulated. Increasing HSP90 activity might represent a new strategy to treat hypertension.

Summary. Ang II down-regulates Kv7.4 by disrupting its interaction with HSP90, which lead to the recruitment of the ubiquitin-ligase CHIP and the subsequent protein degradation via the proteasome.

FIGURE LEGENDS

Fig. 1. Effects of Ang II on K_v7.4 expression and function. (A) Top: representative western blot showing K_v7.4 (detected with an anti-K_v7.4 antibody) and β-actin in mesenteric arteries after incubation with 100nmol/L Ang II for 1, 3 or 7 hours. Bottom: Quantification of western blot experiments. n=4-6. *=*p*<0.05; **=*p*<0.01 (Student's t-test). (B) Quantitative PCR showing *kcnq4* mRNA levels after incubation with 100nmol/L Ang II for 7h. Data are expressed using the 2^{-ΔCt} formula. (C) In cell western blot showing the levels of EGFP-K_v7.4 (detected with an anti-GFP antibody) in transfected SGVSM-9 after incubation with 100nmol/L Ang II for 1h. Data are expressed as percentage of the average of the controls for each experimental session. n=12-14 wells per experimental point from 3-4 sessions. ***=*p*<0.001 (Student's t-test). The inset shows a SGVSM9 cell transfected with EGFP-K_v7.4, with GFP fluorescence (green pseudocolor) and DAPI staining (blue pseudocolor), proving the effective expression of EGFP-K_v7.4 protein. Scale bar: 5μm. (D) Mean current-voltage relationship (I-V) of the XE991-sensitive current in mesenteric VSMCs after 3 hours incubation in absence (control, black) or in presence (Ang II, grey) of 100nmol/L Ang II. n=5 cells per experimental point from 5 rats. **=*p*<0.01; ****=*p*<0.0001 (two-way ANOVA). (E) Changes in resting membrane potential (ΔE_m) of VSMCs in whole mesenteric artery induced by 10μmol/L ML213 (black columns) and 10μmol/L levromakalim (grey columns) after 2 hours incubation in absence (control) or in presence (Ang II) of 100nmol/L. (F) Isometric tension recordings in mesenteric arteries incubated for 3 hours in absence (CTL, black) or presence (Ang II, grey) of 100nmol/L Ang II. The graph shows the relaxation induced by increasing concentration of ML213 in arteries pre-contracted with 1μmol/L U46619. Data are expressed as percentage of the maximum contraction to U46619. n=4-5. **=*p*<0.01; ***=*p*<0.001 (two-way ANOVA).

Fig. 2. Effects of Ang II on K_v7.4 subcellular distribution in mesenteric artery myocytes and role of endocytosis and proteasome in Ang II-induced down-regulation of K_v7.4. (A) Immunofluorescence experiments showing K_v7.4 (red pseudocolor) in absence (CTL) or in presence (Ang II) of 100nmol/L Ang II. Staining of the plasma membrane marker wheat germ agglutinin (WGA, green pseudocolor) and nuclear marker DAPI (blue pseudocolor) are also shown. Scale bar: 5μm. (B) Top: fluorescence intensity profiles for K_v7.4 (red) and WGA (green) along the yellow line drawn in the pictures, expressed as Arbitrary Units (AU), in control (CTL) or Ang II-treated VSMCs. Regions where WGA intensity was <5AU were considered as cytosol (C), while portions where intensity was above the threshold were identified as plasma membrane (M). Bottom: bar graph showing the ratio between the average fluorescence intensities in the plasma-membrane and cytosol in VSMCs. For each cell, the mean of the ratios obtained along three random lines was calculated. n=15 cells per experimental point, obtained from 3-4 rats in 3-4 sessions. ***=p<0.001 (Student's t-test). (C) Top: representative western blot showing K_v7.4 and β-actin proteins in mesenteric arteries after incubation for 3h with different combinations of 100nmol/L Ang II, 100μmol/L dynasore (DYN), and 20μmol/L MG132, as indicated. Bottom: quantification of western blot experiments. Normalized K_v7.4 intensities are expressed as percentage of control. n=6: **=p<0.01, ns=non significant (one-way ANOVA). (D) Isometric tension recordings in mesenteric arteries incubated for 3 hours in control conditions or with 100nmol/L Ang II in absence or presence of 20μmol/L MG132. The graph shows the relaxation to ML213 in arteries pre-contracted with 300nmol/L U46619. Data are expressed as percentage of the maximum contraction to U46619. n=4-6. *=p<0.05; **=p<0.01; ****=p<0.0001 (two-way ANOVA).

Fig. 3. Role of oxidative stress in the downregulation of K_v7.4 induced by Ang II. (A) Left: representative images showing the fluorescence of the ROS-sensitive dye ROS Deep

Red in mesenteric VSMCs after incubation for 30 min (upper panels) or 3 hours (lower panels) in control medium (CTL) or in presence of 100nmol/L Ang II or 1mmol/L H₂O₂. Right: quantification of ROS Deep Red fluorescence intensity in VSMCs, expressed as Arbitrary Units (AU). n=35-48 cells obtained from 4-5 rats per experimental point in 4-5 sessions. *=p<0.05, ns=not significant (one-way ANOVA). (B) The inset shows a representative western blot of K_v7.4 and β-actin proteins in mesenteric arteries after incubation for 3 hours with different combination of 100nmol/L Ang II and 1mmol/L of N-Acetyl-L-Cysteine (NAC), as indicated. The bar graph shows the quantification of western blot experiments. n=6. *=p<0.05 (one-way ANOVA). (C) Isometric tension recordings in mesenteric arteries incubated for 3 hours in control condition or with 100nmol/L Ang II in absence of presence of 1mmol/L NAC. The graph shows the relaxation to ML213 in arteries pre-contracted with 1μmol/L U46619. Data are expressed as percentage of the maximum contraction to U46619. n=6. **=p<0.01, ***=p<0.001 ****=p<0.0001 (two-way ANOVA).

Fig. 4. Effects of Ang II on K_v7.4 interaction with Heat Shock Proteins and ubiquitin.







Proximity Ligation Assay showing the interaction of K_v7.4 with HSP70 (A), HSP90 (B), CHIP (C), and ubiquitin (D) in mesenteric artery myocytes. Representative fluorescence and bright field (insets) confocal mid-cell xy sections of mesenteric VSMCs in absence (CTL) or presence (Ang II) of 100nmol/L Ang II are shown for each panel. Red puncta indicate target proteins are in close proximity (<40 nm). Bar graphs show the quantification of the mean number of PLA signals per mid-cell xy section. n=17-40 cells from 3-4 rats per experimental point in 3-4 sessions. *=p<0.05, ***=p<0.001, ****=p<0.0001 (Student's t-test). Nuclei (DAPI staining, blue) are also shown. Scale bar: 5μm.





Fig. 5. Effects of HSP90 inhibition on K_v7.4 levels, interaction with CHIP, and ubiquitination.


(A) Left: representative western blot showing K_v7.4 and β-actin in mesenteric arteries incubated with DMSO (CTL) or 1μmol/L 17-AAG. Right: quantification

of western blot experiments. $n=5$. $*=p<0.05$ (Student's t-test). (B-C) Proximity Ligation Assay showing the interaction of $K_v7.4$ with CHIP (B) and ubiquitin (C) in mesenteric artery myocytes. Representative fluorescence and bright field (insets) confocal mid-cell xy sections of mesenteric artery myocytes incubated with DMSO (CTL) or $1\mu\text{mol/L}$ 17-AAG are shown for both panels. Red puncta indicate target proteins are in close proximity ($<40\text{nm}$). Nuclei (DAPI staining) are shown in blue pseudocolor. Scale bar: $5\mu\text{m}$. Bar graphs show the quantification of the mean number of PLA signals per mid-cell xy section. $n=24-44$ cells from 3-4 rats per experimental point in 3-4 sessions. $*=p<0.05$, $****=p<0.0001$ (Student's t-test).




Fig. 6. $K_v7.4$ interaction with Heat Shock Proteins, CHIP and ubiquitin in Ang II-infused mice. Proximity Ligation Assay showing the interaction of $K_v7.4$ with HSP70 (A), HSP90 (B), CHIP (C), and ubiquitin (D) in mesenteric artery myocytes obtained from mice infused for 28 days with saline solution or Ang II. Representative fluorescence and bright field (insets) confocal mid-cell xy sections of mesenteric VSMCs from saline- or Ang II-infused mice are shown for each panel. Red puncta indicate target proteins are in close proximity ($<40\text{ nm}$). Bar graphs show the quantification of the mean number of PLA signals per mid-cell xy section. $n=35-40$ cells from 4 mice per experimental group. $*=p<0.05$, $**=p<0.01$, $****=p<0.0001$ (Student's t-test). Nuclei (DAPI staining, blue) are also shown. Scale bar: $5\mu\text{m}$.





A



Corrected Total Cell Fluorescence (CTCF, Arbitrary Unit)

**B****C**

A**K_v7.4:HSP70**



CTL

Ang II

**B****K_v7.4:HSP90**



CTL

Ang II

**C****K_v7.4:CHIP**



CTL







Ang II




**D****K_v7.4:Ubiquitin**

CTL

Ang II



A**B****K_v7.4:CHIP****C****K_v7.4:ubiquitin**

A**K_v7.4:HSP70****B****K_v7.4:HSP90****C****K_v7.4:CHIP****D****K_v7.4:Ubiquitin**