

COVID-19 and preterm birth

In their Comment (September, 2020), Vimla Kumari and colleagues¹ analyse the impact of the COVID-19 pandemic in four tertiary teaching hospitals in western India. The authors report that, compared with before the pandemic, the demographics of pregnant women delivering in their facilities has changed (ie, there are more educated and nulliparous women), and there is an increased number of in-hospital maternal deaths, and an increased number of late pregnancy fetal losses. During the lockdown period there was a 43·2% reduction in hospital admissions when compared with the control period (ie, 10 weeks before lockdown; 6209 vs 3527), and a 66·4% reduction in referred obstetric emergencies compared with the same calendar period in the previous year (905 vs 304).¹

These results are interesting, as the increase in maternal deaths combined with the observed reduction in the number of emergency referrals during lockdown compared with before lockdown suggest that, not only are fewer women seeking care, the local maternity health system is also failing to seek care for these women. This observation continues to illustrate the indirect effects of COVID-19 on maternal and perinatal health, which we had previously warned about,² and that there is an urgent need to comprehensively report all important obstetric and neonatal outcomes to highlight both predictable and understandable outcomes (eg, an increase in the incidence of stillbirth³) and unexpected patterns of outcomes (eg, the contradictory global evidence about preterm birth rates^{3–5}). Lower preterm birth rates during the COVID-19 pandemic compared with before the pandemic would not be explained by reduced care-seeking behaviour, but could reflect countervailing reductions in maternal activity (high amounts of

which could potentially be associated with an increased risk of preterm birth), changes in clinical decision making that might reduce iatrogenic prematurity, or both. Do Kumari and colleagues¹ have information about changes in the number of preterm births during lockdown to help present a more complete picture?

We declare no competing interests.

Copyright © 2019 The Author(s). Published by Elsevier Ltd. This is an Open Access article under the CC BY 4.0 license.

Laura A Magee, Peter von Dadelszen,

**Asma Khalil*

akhalil@sgul.ac.uk

Vascular Biology Research Centre, Molecular and Clinical Sciences Research Institute, St George's University of London, London, UK (LAM, AK); School of Life Course Sciences, Faculty of Life Sciences and Medicine, King's College London, London, UK (PvD); and Fetal Medicine Unit, Department of Obstetrics and Gynaecology, St George's University Hospitals NHS Foundation Trust, London SW17 0QT, UK (AK)

- 1 Kumari V, Mehta K, Choudhary R. COVID-19 outbreak and decreased hospitalisation of pregnant women in labour. *Lancet Glob Health* 2020; **8**: e1116–17.
- 2 von Dadelszen P, Khalil A, Wolfe I, Kametas NA, O'Brien P, Magee LA. "Women and children last"—effects of the covid-19 pandemic on reproductive, perinatal, and paediatric health. *BMJ* 2020; **369**: m2287.
- 3 Khalil A, von Dadelszen P, Draycott T, Ugwumadu A, O'Brien P, Magee L. Change in the incidence of stillbirth and preterm delivery during the COVID-19 pandemic. *JAMA* 2020; **324**: 705–06.
- 4 Hedermann G, Hedley PL, Baekvad-Hansen M, et al. Danish premature birth rates during the COVID-19 lockdown. *Arch Dis Child Fetal Neonatal Ed* 2020; published online Aug 9. <https://doi.org/10.1136/archdischild-2020-319990>.
- 5 KC A, Gurung R, Kinney MV, et al. Effect of the COVID-19 pandemic response on intrapartum care, stillbirth, and neonatal mortality outcomes in Nepal: a prospective observational study. *Lancet Glob Health* 2020; **8**: e1273–81.



Lancet Glob Health 2020

Published Online
November 20, 2020
[https://doi.org/10.1016/S2214-109X\(20\)30457-5](https://doi.org/10.1016/S2214-109X(20)30457-5)