Supplemental Material

Long-Term Exposure to Fine Particle Elemental Components and Natural and Cause-Specific Mortality—a Pooled Analysis of Eight European Cohorts within the ELAPSE Project

Jie Chen, Sophia Rodopoulou, Kees de Hoogh, Maciej Strak, Zorana J. Andersen, Richard Atkinson, Mariska Bauwelinck, Tom Bellander, Jørgen Brandt, Giulia Cesaroni, Hans Concin, Daniela Fecht, Francesco Forastiere, John Gulliver, Ole Hertel, Barbara Hoffmann, Ulla Arthur Hvidtfeldt, Nicole A. H. Janssen, Karl-Heinz Jöckel, Jeanette Jørgensen, Klea Katsouyanni, Matthias Ketzel, Jochem O. Klompmaker, Anton Lager, Karin Leander, Shuo Liu, Petter Ljungman, Conor J. MacDonald, Patrik K.E. Magnusson, Amar Mehta, Gabriele Nagel, Bente Oftedal, Göran Pershagen, Annette Peters, Ole Raaschou-Nielsen, Matteo Renzi, Debora Rizzuto, Evangelia Samoli, Yvonne T. van der Schouw, Sara Schramm, Per Schwarze, Torben Sigsgaard, Mette Sørensen, Massimo Stafoggia, Anne Tjønneland, Danielle Vienneau, Gudrun Weinmayr, Kathrin Wolf, Bert Brunekreef, and Gerard Hoek

Table of Contents

Table S7. Characteristics of the Cooperative Health Research in the Region of Augsburg (KORA).

Table S8. Characteristics of the Vorarlberg Health Monitoring and Prevention Programme (VHM&PP).

Table S9. Performance of Europe-wide $PM_{2.5}$ composition models in five-fold hold-out validation.

Table S10. Truncation frequency (Truncation performed for model 3 population, N=323,782).

Table S11. Exposure distribution of $PM_{2.5}$ composition in the pooled cohort.

Table S12. Spearman correlation coefficient between $PM_{2.5}$ composition and $PM_{2.5}$ mass (N=323,782).

Table S13. Spearman correlation coefficients between $PM_{2.5}$ composition and NO_2 (N=323,782).

Table S14. Associations of $PM_{2.5}$ composition with natural mortality in single pollutant and two-pollutant models.

Table S15. Associations of $PM_{2.5}$ composition with natural mortality with increasing control for covariates.

Table S16. Associations of $PM_{2.5}$ composition with cardiovascular mortality in single pollutant and two-pollutant models.

Table S17. Associations of $PM_{2.5}$ composition with non-malignant respiratory mortality in single pollutant and two-pollutant models.

Table S18. Associations of $PM_{2.5}$ composition with lung cancer mortality in single pollutant and two-pollutant models.

Table S19. Sensitivity analysis: Associations of $PM_{2.5}$ composition with natural mortality with restricted follow-up period.

Table S20. Sensitivity analysis: Associations between $PM_{2.5}$ composition and natural mortality with additional adjustment for individual level socio-economic status.

Table S21. Sensitivity analysis: Associations between $PM_{2.5}$ composition and natural mortality with additional adjustment for neighborhood-level socio-economic status.

Figure S1. Map of study areas.

Figure S2. Average of cohort-specific Spearman correlation coefficients between $PM_{2.5}$ composition.

Figure S3. Natural spline (3 degree of freedom) for associations between PM$_{2.5}$ composition and natural cause mortality.

References