

B. Vallet  
M. R. Pinsky  
M. Cecconi

## Resuscitation of patients with septic shock: please “mind the gap”!

Received: 30 May 2013

Accepted: 7 June 2013

Published online: 29 June 2013

© The Author(s) 2013. This article is published with open access at Springerlink.com

B. Vallet (✉)

Department of Anesthesiology and Critical Care Medicine  
University Hospital of Lille, Univ Nord de France, Lille, France  
e-mail: benoit.vallet@chru-lille.fr  
Tel.: +33-3-20444447  
Fax: +33-3-20444440

M. R. Pinsky

Department of Critical Care Medicine, University of Pittsburgh,  
Pittsburgh, PA, USA


M. Cecconi

Department of Intensive Care Medicine, St George's Healthcare  
NHS Trust and St George's University of London, London, UK

Van Beest et al. [1] performed a post hoc analysis of 53 patients with severe sepsis or septic shock to investigate the interchangeability of mixed and central venous-to-arterial carbon dioxide (CO<sub>2</sub>) differences (mvaCO<sub>2</sub>gap and cvaCO<sub>2</sub>gap, respectively) and the relation between the cvaCO<sub>2</sub>gap (“pCO<sub>2</sub>gap” or the “gap”), cardiac index (CI), and outcome. The authors observed a strong agreement between pCO<sub>2</sub> measured from either mixed venous or central venous sites with relatively small limits of agreement. The authors claim that combining ScvO<sub>2</sub> values, as easily obtained from a central venous catheter, as a surrogate for global tissue hypoxia, and pCO<sub>2</sub>gap as a surrogate for CI, obtained from the same central venous catheter, may be useful in assessing cardiovascular state during resuscitation in critically ill patients. We cannot

agree more and propose thereafter a tentative “ScvO<sub>2</sub>-cvaCO<sub>2</sub>gap-guided protocol” (Fig. 1). Cuschieri et al. [2] previously demonstrated in a mixed population of critically ill patients that the relationships between the mvaCO<sub>2</sub>gap or the cvaCO<sub>2</sub>gap and the CI were equivalent. Since central venous blood is readily available from a central venous catheter, whereas mixed venous blood requires a pulmonary artery catheter, the cvaCO<sub>2</sub>gap, as an easily available clinical monitoring tool, is attractive. At ICU admission, 24 patients in the van Beest et al. study had a pCO<sub>2</sub>gap greater than 0.8 kPa (or 6 mmHg). Persistence of such a large pCO<sub>2</sub>gap after 24 h of treatment was predictive of higher mortality.

These data are in line with results of Vallée et al. [3] who prospectively tested this hypothesis in 56 septic shock patients resuscitated to an ScvO<sub>2</sub> greater than 70 % (according to the Rivers' study results [4]). They found that patients who still had altered tissue perfusion (assessed by serum lactate levels greater than 2 mmol L<sup>-1</sup>) in spite of a normalized ScvO<sub>2</sub> displayed a large cvaCO<sub>2</sub>gap (greater than 6 mmHg). CO<sub>2</sub> is the end product of aerobic metabolism and its concentration in the venous blood reflects the global tissue blood flow relative to metabolic demand. Since CO<sub>2</sub> is about 20 times more soluble than O<sub>2</sub> the likelihood of it diffusing out of ischemic tissues and into the venous effluent is great, making it a very sensitive marker of hypoperfusion. Thus, in situations where an O<sub>2</sub> diffusion barrier exists (resulting from non-functional and obliterated capillaries), “masking” poor O<sub>2</sub> extraction (O<sub>2</sub>ER) and increased tissue O<sub>2</sub> debt, CO<sub>2</sub> still diffuses to the venous effluent, “unmasking” the low perfusion state for the clinician when venous-to-arterial CO<sub>2</sub> difference is evaluated. Consistently Vallée et al. [3] evidenced that patients with high cvaCO<sub>2</sub>gap values had lower lactate clearance and CI values, and presented a significant lower decrease in Sepsis-related Organ Failure Assessment score than patients with a low cvaCO<sub>2</sub>gap. Thus, the cvaCO<sub>2</sub>gap



**Fig. 1** The ScvO<sub>2</sub>-cvaCO<sub>2</sub>gap-guided protocol. ScvO<sub>2</sub> venous central O<sub>2</sub> saturation, cvCO<sub>2</sub>gap central venous-to-arterial pCO<sub>2</sub> difference, SaO<sub>2</sub> arterial O<sub>2</sub> saturation, CI cardiac index, SVV stroke volume variability, PPV pulse pressure variability

represents a useful complementary tool to identify patients who remain inadequately resuscitated when the 70 % ScvO<sub>2</sub> threshold value has been reached.

The obvious limitation of ScvO<sub>2</sub> is therefore that normal/high values cannot discriminate whether delivery is adequate or in excess to demand. High ScvO<sub>2</sub> profiles have even been shown to be related to elevated blood lactate concentration and poor survival rates [5]. Although ScvO<sub>2</sub> may thus not miss any global oxygen delivery (DO<sub>2</sub>) disorder, it may remain “blind” to local perfusion disorders, which abound in sepsis because of impaired microcirculation. Under conditions where O<sub>2</sub> uptake (VO<sub>2</sub>) does not meet O<sub>2</sub> demand, tissue dysoxia occurs, leading to organ failure and death. The crucial point here is that these tissues might remain however accessible to conventional therapeutic hemodynamic management (inotropes and fluid infusion).

Whether the resultant effect on pCO<sub>2</sub> gap depends in principle on the flow state or anaerobic CO<sub>2</sub> production was tested by Vallet et al. [6] in an experimental model of isolated limb in which ischemic hypoxia (IH) and hypoxic hypoxia (HH) were compared. The authors demonstrated that when DO<sub>2</sub> was reduced beyond its critical threshold in IH (dysoxia), this was associated with an increased limb venous-to-arterial pCO<sub>2</sub>gap [6]. Conversely, in HH, pCO<sub>2</sub>gap did not increase in spite of a marked VO<sub>2</sub>

**Table 1** Lactate-ScvO<sub>2</sub>-cvaCO<sub>2</sub>gap as shock diagnostic tools

Shock type	Lactate	O <sub>2</sub> ER	ScvO <sub>2</sub>	cvaCO <sub>2</sub> gap
Cardiogenic hypovolemic	HI	HI	LO	HI
Anemic hypoxemic	HI	HI	LO	LO
Distributive	HI	LO	HI	HI
Cytopathic	HI	LO	HI	LO

ScvO<sub>2</sub> venous central O<sub>2</sub> saturation, cvCO<sub>2</sub>gap central venous-to-arterial pCO<sub>2</sub> difference, O<sub>2</sub>ER O<sub>2</sub> extraction ratio, HI high, LO low

and VCO<sub>2</sub> reduction, clearly evidencing the gap as a marker of adequacy of venous blood flow to remove CO<sub>2</sub> produced rather than a marker of tissue hypoxia or dysoxia.

In conclusion, determining the gap during resuscitation of critically ill patients is useful when deciding when to stop resuscitation despite persistent evidence of organ ischemia and an ScvO<sub>2</sub> of greater than 70 % (see Fig. 1, Table 1). All forms of circulatory stress are potentially associated with hyperlactatemia, but hyperlactatemia is not a discriminatory factor in defining the cause of that stress. A goal of a gap lower than 6 could be a useful complementary tool to evaluate the adequacy of blood flow to global metabolic demand. In this regard it can help to titrate inotropes in order to adapt DO<sub>2</sub> to VCO<sub>2</sub>, or to choose between hemoglobin correction or fluid/inotrope

infusion. Whatever way you use it or like it, from mixed or central venous, in patients with septic shock, please “mind the gap”!

**Conflicts of interest** None.

**Open Access** This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.

## References

1. Van Beest PA, Lont MC, Holman ND, Loef B, Kuiper MA, Boerma EC (2013) Central venous-arterial pCO<sub>2</sub> difference as a tool in resuscitation of septic patients. *Intensive Care Med* 39:1034–1039
2. Cuschieri J, Rivers EP, Donnino MW, Katilius M, Jacobsen G, Nguyen HB, Pamukov N, Horst HM (2005) Central venous-arterial carbon dioxide difference as an indicator of cardiac index. *Intensive Care Med* 31:818–822
3. Vallée F, Vallet B, Mathe O, Parraguette J, Mari A, Silva S, Samii K, Fourcade O, Genestal M (2008) Central venous-to-arterial carbon dioxide difference: an additional target for goal-directed therapy in septic shock? *Intensive Care Med* 34:2218–2225
4. Rivers E, Nguyen B, Havstad S, Ressler J, Muzzin A, Knoblich B, Peterson E, Tomlanovich M, Early Goal-Directed Therapy Collaborative Group (2001) Early goal-directed therapy in the treatment of severe sepsis and septic shock. *N Engl J Med* 345:1368–1377
5. Puskarich MA, Trzeciak S, Shapiro NI, Arnold RC, Heffner AC, Kline JA, Jones AE, Emergency Medicine Shock Research Network (EMSHOCKNET) (2012) Prognostic value and agreement of achieving lactate clearance or central venous oxygen saturation goals during early sepsis resuscitation. *Acad Emerg Med* 19:252–258
6. Vallet B, Teboul JL, Cain S, Curtis S (2000) Venoarterial CO<sub>2</sub> difference during regional ischemic or hypoxic hypoxia. *J Appl Physiol* 89:1317–1321