

REVIEW

## Editor's Choice — The Implications of Non-compliance to Endovascular Aneurysm Repair Surveillance: A Systematic Review and Meta-analysis

Matthew Joe Grima<sup>a,b,\*</sup>, Mourad Boufi<sup>a,c,e</sup>, Martin Law<sup>d</sup>, Dan Jackson<sup>d</sup>, Kate Stenson<sup>a,b</sup>, Benjamin Patterson<sup>a,b</sup>, Ian Loftus<sup>a,b</sup>, Matt Thompson<sup>a,b</sup>, Alan Karthikesalingam<sup>a,b</sup>, Peter Holt<sup>a,b</sup>

<sup>a</sup> St George's Vascular Institute, St George's Hospital, NHS Foundation Trust, London, UK

<sup>b</sup> Molecular and Clinical Sciences Research Institute, St George's, University of London, London, UK

<sup>c</sup> Aix-Marseille Université, CNRS, IRPHE UMR 7342, Marseille, France

<sup>d</sup> MRC Biostatistics Unit, Institute of Public Health, Cambridge, UK

<sup>e</sup> APHM, Department of Vascular Surgery, University Hospital Nord, Marseille, France

### WHAT THIS PAPER ADDS

Surveillance imaging is considered mandatory after endovascular aneurysm repair (EVAR), but many patients are either non-compliant or lost to follow-up, and the impact of this is poorly understood. This review highlights and confirms the great variability in published EVAR surveillance compliance rates. This study also suggests that although compliance may be associated with increased re-interventions after EVAR, surveillance does not appear to confer a survival advantage to compliant patients in the first 5 years after EVAR.

**Objective/background:** Increasingly, reports show that compliance rates with endovascular aneurysm repair (EVAR) surveillance are often suboptimal. The aim of this study was to determine the safety implications of non-compliance with surveillance.

**Methods:** The study was carried out according to the Preferred Items for Reporting of Systematic Reviews and Meta-Analyses (PRISMA) guidelines. An electronic search was undertaken by two independent authors using Embase, MEDLINE, Cochrane, and Web of Science databases from 1990 to July 2017. Only studies that analysed infrarenal EVAR and had a definition of non-compliance described as weeks or months without imaging surveillance were analysed. Meta-analysis was carried out using the random-effects model and restricted maximum likelihood estimation.

**Results:** Thirteen articles (40,730 patients) were eligible for systematic review; of these, seven studies (14,311 patients) were appropriate for comparative meta-analyses of mortality rates. Three studies (8316 patients) were eligible for the comparative meta-analyses of re-intervention rates after EVAR and four studies (12,995 patients) eligible for meta-analysis for abdominal aortic aneurysm related mortality (ARM). The estimated average non-compliance rate was 42.0% (95% confidence interval [CI] 28–56%). Although there is some evidence that non-compliant patients have better survival rates, there was no statistically significant difference in all cause mortality rates (year 1: odds ratio [OR] 5.77, 95% CI 0.74–45.14; year 3: OR 2.28, 95% CI 0.92–5.66; year 5: OR 1.81, 95% CI 0.88–3.74) and ARM (OR 1.47, 95% CI 0.99–2.19) between compliant and non-compliant patients in the first 5 years after EVAR. The re-intervention rate was statistically significantly higher in compliant patients from 3 to 5 years after EVAR (year 1: OR 6.36, 95% CI 0.23–172.73; year 3: OR 3.94, 85% CI 1.46–10.69; year 5: OR 5.34, 95% CI 1.87–15.29).

**Conclusion:** This systematic review and meta-analysis suggests that patients compliant with EVAR surveillance programmes may have an increased re-intervention rate but do not appear to have better survival rates than non-compliant patients.

© 2017 The Author(s). Published by Elsevier Ltd on behalf of European Society for Vascular Surgery. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Article history: Received 10 March 2017, Accepted 27 November 2017, Available online 5 January 2018

**Keywords:** Abdominal aortic aneurysm, Endovascular procedures, Epidemiology, Meta-analysis, Review, Stents

\* Corresponding author. St George's Vascular Institute, Room 0.231, Corridor 10, Ground Floor, Jenner Wing, St George's, University of London, Cranmer Terrace, London, SW17 0RE, UK.

E-mail address: [matthewjoe.grima@gmail.com](mailto:matthewjoe.grima@gmail.com) (Matthew Joe Grima).

Twitter: [MJ\\_Grima](#)

1078-5884/© 2017 The Author(s). Published by Elsevier Ltd on behalf of European Society for Vascular Surgery. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

<https://doi.org/10.1016/j.ejvs.2017.11.030>

## INTRODUCTION

Despite endovascular aneurysm repair (EVAR) being the modern preferred first choice for repair of abdominal aortic aneurysms (AAA),<sup>1,2</sup> studies show that re-interventions after EVAR are common and are undertaken in around 20% of patients within 5 years.<sup>3,4</sup> Consequently, guidelines from learned societies recommend lifelong annual imaging in order to identify and treat aortic complications to prevent aneurysm rupture and death.<sup>5–7</sup> However, published population and observational studies show that patients are not always compliant with their surveillance programmes.<sup>8,9</sup> A number of studies have attempted to evaluate patient characteristics that may be associated with poor compliance rates.<sup>8,10</sup> Despite this, little is known about the consequence of non-compliance with surveillance. Thus, a systematic review and meta-analysis was undertaken to study the implications of non-compliance with EVAR surveillance programmes. The primary outcomes were overall compliance, all cause mortality (ACM), and re-intervention rates and the secondary outcome was aneurysm related mortality (ARM).

## METHODS

The study was carried out according to the Preferred Items for Reporting of Systematic Reviews and Meta-Analyses (PRISMA) guidelines.<sup>11</sup> An electronic search was undertaken by two independent authors using the Embase, Medline, and Cochrane databases from 1990 to July 2017. Studies that assessed the compliance rate with surveillance after EVAR and analysed the relationship of re-intervention and mortality rates with compliance rates were identified. The search terms (including medical subject sub-headings) “abdominal aortic aneurysm”, “aneurysm”, “AAA”, “EVAR”, “endovascular repair”, “compliance”, “surveillance”, “follow-up”, and “survey” were used in combination with Boolean operators AND or OR. The reference lists of articles obtained were investigated to identify relevant citations. Conference abstracts from major vascular meetings, when published online, were also scrutinised through the Web of Science database (full search history is available in the [Supplementary Material](#)).

Inclusion criteria encompassed all studies describing endovascular repair of infrarenal AAA. The studies needed to have a definition of non-compliance described as weeks or months without imaging surveillance. Exclusion criteria included non-English language papers, thoraco-abdominal aortic aneurysm repair, suprarenal AAA, fenestrated grafts, parallel grafts, iliac aneurysms, and patients treated with the endovascular aneurysm sealing technique.

Studies that provided follow-up data using statistical methods for survival analysis were used for comparative meta-analyses of ACM and re-intervention rates. Quality assessment was carried out independently by two authors using Grading of Recommendations, Assessment, Development and Evaluations (GRADE),<sup>12</sup> and differences were resolved through discussion between the two authors. Outcome data were obtained from at risk scores provided with the tables and graphs when available, but if not, data

were extracted from Kaplan–Meier curves. Attempts were made to contact the authors whenever data required were not readily available.

ARM was standardised by Chaikof et al. as deaths secondary to aneurysm rupture, EVAR to open conversion, and the index or secondary procedure (see [Table 7](#)).<sup>13</sup>

As different institutions and studies used different definitions for non-compliance, a laxity index was developed by the authors at the outset of the study. The laxity index is a measure of the stringency of the studies’ definition of non-compliance. The laxity index was based on the number of scans missed and the number of months without imaging. A laxity value (from 0 to 1) was attributed to studies. A low laxity index suggests a very rigid application of the surveillance protocol, such that minimal deviation was labelled as non-compliance (detailed explanation is available in the [Supplementary Material](#)).

## Statistical analysis

Statistical analysis was carried out using ‘R package metafor’. Random effects meta-analyses were performed using restricted maximum likelihood estimation.<sup>14</sup> The meta-analysis for the non-compliance rates was performed using the observed rates and standard methods for a non-comparative proportion. For four of the papers,<sup>8,15–17</sup> reported longer-term compliance rates were used to determine the necessary outcome data.

The comparative meta-analyses of ACM and re-intervention rates were performed at five time points (1, 2, 3, 4, and 5 years) after intervention. The outcome data were empirical log-odds ratios (ORs) that compare the all cause mortality and re-intervention rates of the compliant and non-compliant patients. In order to include outcome data where Kaplan–Meier curves indicated that the event rate was negligible, the corresponding outcome data were analysed using the rate where half a person had experienced the event. Random effects meta-analyses were done using standard methods, where a conservative sample size was used for calculating the within study variances, so that censoring resulted in the maximum possible loss of information.<sup>18</sup> This sample size calculation requires the number at risk at each time point. Most papers gave these or values at adjacent time points that could be used for interpolation. Where numbers at risk were not given in study reports, for the purposes of calculating within study variances, the sample sizes were reduced by the average percentage reduction across the other studies that contribute to the analysis. Pooled estimates were transformed to the OR scale, where an OR that is  $> 1$  indicates that the mortality or re-intervention rate is higher in compliant patients. To account for confounding factors, matched cohorts were used where possible. This included the studies by Garg et al. and Hicks et al. for comparative meta-analyses of mortality,<sup>16,19</sup> and the study by Garg et al. for comparative meta-analyses of re-intervention rates.<sup>19</sup>

Five random effects meta-regression models were fitted where the overall survival log-ORs were regressed on the

laxity index. Here the regression coefficient is a log-OR that is associated with a change in the laxity index from its minimum (0) to its maximum (1). A positive regression coefficient indicates that a greater laxity index is associated with larger ORs. A single comparative meta-analysis of the rates of ARM was performed. Here a random effects meta-analysis was performed, using empirical log-ORs as outcome data, and as in the other comparative meta-analyses.

For three studies,<sup>16,20,21</sup> Kaplan–Meier curves for “non-compliant” groups and “lost to follow-up” were provided separately. It was agreed that, although the “lost to follow-up” group is a more non-compliant group, this particular survival curve would not be analysed for meta-analysis. This decision was taken as this provides a more conservative analysis and avoids statistical issues involved in reconstructing Kaplan–Meier curves *de novo* (which would be a combination of non-compliant and lost to follow-up). Thus, the results produced will be more transparent and easily reproducible.

Elective and emergency cases of AAA repair could not be distinguished to statistically assess the impact of survival and re-intervention rate. Thus, the statistical outcomes do not refer to elective or rupture patients independently.

In the study by Hicks et al.,<sup>16</sup> an assumption was agreed by the authors, whereby patients who had telephone follow-up were considered as non-compliant, whereas patients who attended clinic for follow-up were considered as compliant. This was done in order to resemble the current real world scenario where patients physically attend follow-up sessions. Given that the recorded follow up in the Vascular Quality Initiative register is only for 1 year after EVAR, a sensitivity analysis for all cause mortality excluding the study by Hicks et al. was carried out.<sup>16</sup>

## RESULTS

After screening, the literature search identified 31 articles that discussed compliance after EVAR. Of these 31 articles, 18 were excluded for various reasons (Fig. 1).<sup>9,10,22–37</sup> Of the 13 articles that were suitable for systematic review,<sup>8,15–17,19–21,38–43</sup> seven studies were eligible for comparative meta-analysis of ACM.<sup>16,19–21,40–42</sup> Three were eligible for comparative meta-analyses of re-intervention rates after EVAR,<sup>19,40,42</sup> whereas four studies were eligible for random effects meta-analyses of ARM.<sup>19,20,41,42</sup> The study by Godfrey et al.,<sup>38</sup> despite providing a survival analysis curve, was not eligible for inclusion in meta-analysis owing to a different definition of non-compliance as compared with other papers. This was confirmed when the corresponding author was contacted.

### Primary outcomes

**Compliance.** Using the seven studies that contributed to the comparative meta-analysis, the estimated average non-compliance rate (according to the papers’ specific criteria for determining this) was 42% (95% confidence interval [CI] 28–56% [26,622 patients: 15,255 compliant; 11,367 non-compliant]) (Fig. 2). This finding is consistent even if the

non-compliance rates of the other four studies were included, where the non-compliance rate was 38% (95% CI 27–48%), (40,730 patients: 22,971 compliant; 17,759 non-compliant) (Fig. 2, Table 1). Overall quality of evidence is “moderate” (Table 2).<sup>12</sup>

**ACM.** In total, 14,311 patients (6947 compliant and 7364 non-compliant) contributed to meta-analysis of ACM. There is some evidence that compliant patients have higher ACM rates in the 5 years after EVAR (year 1, OR 5.77 [95% CI 0.74–45.14]; year 3, OR 2.28 [95% CI 0.92–5.66]; year 5, OR 1.81 [95% CI 0.88–3.74]); however, this did not reach statistical significance (Table 3, Fig. 3). Overall quality of evidence is “moderate” (Table 2).<sup>12</sup>

**ACM and laxity index.** Meta-regressions generally suggest that larger laxity indices are associated with larger ORs for overall survival (Tables 4 and 5, Figs. S11–S15). This observation strengthens the conclusions that ACM may be higher in more compliant patient groups, although no regression was statistically significant.

**Re-intervention after EVAR.** In total, 8316 (4298 compliant and 4018 non-compliant) patients contributed to meta-analysis of re-intervention after EVAR. There was no statistically significant difference in the intervention rate between the two groups for the first 2 years after EVAR (year 1, OR 6.36 [95% CI 0.23–172.73]; year 2, OR 8.08 [95% CI 0.25–262.57]). However, re-intervention rates were statistically significantly higher in compliant patients for years 3–5 after EVAR (year 3, OR 3.94 [95% CI 1.46–10.69]; year 4, OR 4.37 [95% CI 1.31–14.57]; year 5, OR 5.34 [95% CI 1.87–15.29]) (Table 6, Fig. 4). Overall quality of evidence is “low” (Table 2).<sup>12</sup>

### Secondary outcome


**AAA related mortality.** In total, 12,995 patients (5890 compliant and 7105 non-compliant) contributed to this analysis. Although there is some evidence that the rate of ARM is higher in compliant patients, this was not statistically significant (OR 1.47, 95% CI 0.99–2.19;  $p = .06$ ) (Tables 7 and 8). Overall quality of evidence is “very low” (Table 2).<sup>12</sup>

### Sensitivity analysis (excluding Hicks et al.<sup>16</sup>)

**ACM.** In total, 13,407 patients (6182 compliant and 7225 non-compliant) contributed. The ACM rate was statistically significantly higher in compliant patients compared with non-compliant patients in the 5 years after EVAR (year 1, OR 9.85 [95% CI 1.14–84.98]; year 2, OR 5.10 [95% CI 1.57–16.56]; year 3, OR 3.26 [95% CI 1.50–7.06]; year 4, OR 2.68 [95% CI 1.35–5.33]; year 5, OR 2.09 [95% CI 1.00–4.35]; see Table S3). These results were sensitive to the decision of whether or not to include this study.


## DISCUSSION

This study highlights the wide variation in published rates of compliance with surveillance after EVAR and confirms the



**Figure 1.** Preferred Items for Reporting of Systematic Reviews and Meta-Analyses (PRISMA) flowchart.

Note. EVAR = endovascular aneurysm repair.



**Figure 2.** Meta-analysis of proportions of non-compliance. Meta-analysis of proportions of non-compliance to endovascular aneurysm repair surveillance using data from the seven studies eligible for comparative meta-analysis, termed “main studies”, was carried out. An overall meta-analysis of proportions of non-compliance using the data from all the studies in the systematic review (even if not eligible for comparative meta-analysis) was carried out. The overall result is shown next to the “model for all studies”.

previously published reports of lack of adherence to surveillance programmes.<sup>8,19</sup> Despite the recommendation by international guidelines that EVAR surveillance is mandatory, it was found in the present review that only around 60% of patients are compliant with their surveillance programmes.

The safety implications of non-compliance have previously been ill defined. Although there was no statistically significant difference, the present results suggest that compliant patients may have higher ACM rates than non-compliant patients. However, the current analyses do not imply any causal link between surveillance and survival, and, as a result, the authors recommend that this finding be interpreted with caution and not be taken at face value.

Potential reasons for this result are likely to be multifactorial. This may be because sicker patients have more imaging for unrelated problems and therefore show a higher rate of overall mortality in the compliant group. This phenomenon was highlighted in a multicentre European study and a US population based study.<sup>19,41</sup> However, in the study by Schanzer et al.,<sup>8</sup> Medicare patients with comorbidities and cardiovascular risk factors were more non-compliant. The authors hypothesised that patients with competing medical pathologies become less inclined to attend EVAR surveillance. Subsequently, given the observation of different attitudes by patients with a high comorbid status, it may indicate that surveillance programmes need to be tailored according to the catchment population.

The mode of presentation to imaging surveillance could potentially explain the results of this meta-analysis. The

study by Karthikesalingam et al. showed that around 60% of patients who had re-intervention presented with symptoms,<sup>4</sup> whereas the systematic review by Nordon et al. showed that >90% of patients presented with symptoms.<sup>3</sup> Thus, patients who were asymptomatic and potentially healthier were less likely to attend surveillance. This may explain the trend of better survival rates in non-compliant patients and why complications were noted to be higher in compliant groups.<sup>21,41</sup> Kret et al. analysed a mixed cohort of treated aortic pathologies and observed that complications were higher in the compliant group,<sup>9</sup> which is in line with the results of this review.

The present study showed that the re-intervention rate was statistically significantly higher in compliant patients for years 3–5 after EVAR. In contrast to this outcome, despite no difference in re-interventions or endoleak, Jones et al. noted a significantly higher rate of major complications in non-compliant patients. Major complication was defined as any complication requiring urgent surgery; however, no definition was provided for re-intervention. This contrasts with the study by Garg et al.,<sup>19</sup> whereby the statistical difference between the two groups persisted, even when all complications, including late rupture or re-intervention, were analysed together. The results of this meta-analysis do not come as a surprise, as the aim of surveillance is to identify asymptomatic complications to allow re-intervention in order to prevent ARM.

The present study indicates that while compliant patients had higher rates of re-intervention, compliance with surveillance was not associated with a lower ARM. This raises a



**Table 1.** Meta-analysis of proportions, along with definitions of, non-compliance and the overall compliant and non-compliant numbers in each study.

Study	Definition of non-compliance	Total number of patients	Non-compliant patients (n)	Compliant patients (n)	Main study
Schanzer et al. <sup>8</sup>	Defined as “prolonged time period”: a patient who did not undergo at least one imaging study during each 2 y interval that they were alive after EVAR	7666	3879	3787	No
Godfrey et al. <sup>38</sup>	No imaging within the preceding 12 months ( $\pm 2$ months) of the surveillance visits	179	50	129	No
Dias et al. <sup>15</sup>	Non-attendance at yearly CT scans	279	5	274	No
Aburahma et al. <sup>39</sup>	No follow-up imaging (CT and/or DUS) for 2 y at any time during follow-up and/or missed first post-EVAR imaging over 6 mo	565	323	242	No
Cohen et al. <sup>43</sup>	Not compliant to follow-up protocol: 30 d, 1 y, and annual follow-up	517	92	425	No
De Mestral et al. <sup>17</sup>	Defined as “minimum appropriate imaging follow-up”: CT scan or an ultrasound of the abdomen within 90 d of EVAR and every 15 mo thereafter	4902	2043	2859	No
Garg et al. <sup>19</sup>	Defined as “incomplete surveillance” if surveillance gaps between images was longer than 15 mo	9695	5526	4169	Yes
Jones et al. <sup>40</sup>	Any patient who missed $\geq 2$ consecutive follow-up office visits	302	99	203	Yes
Leurs et al. <sup>41</sup>	Patients missing $\geq 1$ follow-up appointments	4433	2895	1538	Yes
Sarangarm et al. <sup>42</sup>	Any patient who missed $> 2$ consecutive follow-up office visits	126	19	107	Yes
Waduud et al. <sup>20</sup>	Patients who underwent no imaging in the first 12 mo after EVAR or who missed any subsequent annual imaging appointments thereafter	569	268	301	Yes
Wu et al. <sup>21</sup>	Defined as “moderately compliant” if patients missed appointments or surveillance imaging (either one appointment or multiple ones) but continued to follow-up thereafter	188	99	89	Yes
Hicks et al. <sup>16</sup>	Follow-up is an independent variable in the VQI registry. Patients are recorded as having only a single follow-up. If a patient had multiple follow-up visits, the latest recorded follow up status was used. (Assumption: in person follow up as compliant and telephone call follow-up as non-compliant.)	11,309	2461	8848	Yes

Note. EVAR = endovascular aneurysm repair; CT = computed tomography; DUS = duplex ultrasound; VQI = vascular quality initiative.

**Table 2.** Overall GRADE quality assessment for each outcome.<sup>12</sup>


Studies (n)	Study design	Risk of bias	Inconsistency	Indirectness	Imprecision	Publication bias	Quality
Overall compliance							
13	Observational studies	Not serious	Serious	Not serious	Not serious	Undetected/not tested	⊕⊕⊕○ MODERATE
ACM							
7	Observational studies	Not serious	Not serious	Serious	Not serious	Undetected/not tested	⊕⊕⊕○ MODERATE
Re-interventions							
3	Observational studies	Not serious	Not serious	Serious	Serious	Undetected/not tested	⊕⊕⊕○ LOW
Aneurysm related mortality							
4	Observational studies	Serious	Not serious	Serious	Not serious	Undetected/not tested	⊕○○○ VERY LOW

Note. GRADE = Grading of Recommendations, Assessment, Development and Evaluations; ACM = all cause mortality.

**Table 3.** Comparative meta-analyses of all cause mortality.

Time point (y)	OR	95% CI lower	95% CI upper	p-value	I <sup>2</sup>	τ <sup>2</sup>	Q	p value (Q)
1	5.77	0.74	45.14	0.10	92.95	6.48	36.71	<0.01
2	3.20	0.86	11.96	0.08	97.16	2.75	279.09	<0.01
3	2.28	0.92	5.66	0.08	95.87	1.27	218.82	<0.01
4	1.99	0.89	4.45	0.10	94.43	0.95	133.21	<0.01
5	1.81	0.88	3.74	0.11	90.60	0.71	89.77	<0.01

Note. Odds ratio (OR) > 1 indicates that the mortality rate is higher in compliant patients.



**Figure 3.** Summary of estimated average all cause mortality rates from five meta-analyses. (odds ratio [OR] > 1 indicates that the average mortality rate is higher in compliant patients). Forest plots for each of the five meta-analyses that contribute to this figure are shown in the [Supplementary Material](#).

number of issues, one being whether the increased re-intervention rate in compliant patients is reflected in the observed trend of increased overall mortality and ARM in compliant patients. As a result, more questions arise as to

whether EVAR surveillance is potentially causing unnecessary treatment with the possibility of iatrogenic harm. Another issue is whether the aim of EVAR surveillance—that of detecting and treating asymptomatic sac size increase before causing ARM—is being reached. However, the authors acknowledge that this paper did not study the causality between these outcomes.

This meta-analysis indicates that re-intervention rates for both groups during the first 2 years after EVAR, is very low. Similarly, death events at 1 year after EVAR are low. Hicks et al. also showed that non-compliance with surveillance during the first year after EVAR had a detrimental effect on survival and explains why inferences are sensitive to the inclusion or exclusion of this study.<sup>16</sup>

Apart from this observation, as there is no statistically significant difference in mortality between the two groups during the first 5 years after EVAR, and given that some published studies show that few asymptomatic complications are detected at surveillance,<sup>4,9</sup> this study potentially highlights the benefit of risk stratification of EVAR surveillance after the first year of surveillance. Although it is not the remit of this study to discuss ways of stratification, a model developed by this institution (St George's Vascular Institute score<sup>44</sup>) stratified patients into high and low risk for developing complications using their pre-operative aortic sac size and iliac diameter. Bastos Gonçalves et al. showed that patients with early sac shrinkage, adequate

**Table 4.** Laxity index criteria and explanation.

Study	Laxity index	Surveillance protocol	Definition of non-compliance
Leurs et al. <sup>41</sup>	0	Surveillance 1, 3, 6, 12, 18, and 24 mo, then annually	Incomplete = missing ≥ 1
Wu et al. <sup>21</sup>	0.125	Surveillance 1 and 12 months then annually (but can vary by surgeon or case)	Moderate compliance = missed ≥ 1
Jones et al. <sup>40</sup>	0.25	Surveillance: 1 wk, 1 mo, every 6 mo for 2 y, then annually	Incomplete = missed ≥ 2 follow-up appointments
Sarangarm et al. <sup>42</sup>	0.5	Surveillance: 1 mo, every 6 mo for 2 y, then annually	Incomplete = missed > 2 consecutive appointments
Garg et al. <sup>19</sup>	0.63	Surveillance: 1 mo (6 mo if abnormal 1 mo scan), 12 mo, then annually	Incomplete = gaps of 15 mo without surveillance
Hicks et al. <sup>16</sup>	1	Last recorded visit in 1 y follow-up. SVS guidelines: 30 d, 1 y, and annually after EVAR	Did not attend any in person follow-up after EVAR
Waduud et al. <sup>20</sup>	1	Surveillance varies	Incomplete = no imaging in the first 12 mo or missed any subsequent annual imaging surveillance

Note. Non-compliance in terms of months without scan (every 0.5 = 12 month gap). SVS = Society for Vascular Surgery; EVAR = endovascular aneurysm repair.

**Table 5.** Meta-regression analysis of all cause mortality using the laxity index.

Time point (y)	Coefficient	95% CI lower	95% CI upper	p value
1	1.63	-4.27	7.54	0.59
2	0.61	-3.17	4.39	0.75
3	-0.09	-2.72	2.55	0.95
4	0.23	-2.09	2.56	0.84
5	0.98	-0.88	2.84	0.30

Note. A positive regression coefficient indicates that larger laxity values are associated with larger odds ratios. CI = confidence interval.

EVAR seal and no endoleak during the first year after EVAR have low risk of late complications.<sup>45,46</sup> As a result, resources could be directed to improving and encouraging compliance in high risk patients while subjecting asymptomatic and low risk patients to lower risks of radiation exposure and nephrotoxic effects without the increased risk of re-intervention and mortality.<sup>47,48</sup>


**Limitations**

Different studies had different definitions for non-compliance, whereas some studies grouped non-compliant and lost to follow-up patients together.<sup>40</sup> An attempt to reduce this limitation effect was carried out through the introduction of the laxity index for non-compliance. Although meta-regression analyses perform better if the number of studies analysed is  $\geq 10$ ,<sup>49</sup> meta-regression analysis was carried out to further assess whether compliance is associated with ACM. The meta-regressions using the laxity index, despite their limitations, generally support the findings of the meta-analysis.

Another limitation is the way compliance data were collected by different authors. The studies by Schanzer et al. and Garg et al.,<sup>8,19</sup> which had the largest cohort of patients, included any imaging modality which captured EVAR. Thus, one may question whether the imaging of some patients was actually surveillance imaging. This may infer that overall non-compliance with surveillance may actually be higher than 42%.

Four studies analysed ARM. This event was rare and only a small number of studies provided data, so methods for meta-analysis cannot be expected to be very accurate in this situation.

Another limitation is that definitions used for complications and re-intervention varied between studies. From the three studies that provided Kaplan–Meier curves for analysis of re-intervention, two used the term re-intervention



**Figure 4.** Summary of all estimated average re-intervention rates from five meta-analyses. (odds ratio [OR] > 1 indicates that the average re-intervention rate is higher in compliant patients). Forest plots for each of the five meta-analyses that contribute to this figure are shown in the [Supplementary Material](#).

for AAA complications,<sup>19,42</sup> whereas the study by Jones et al. tried to differentiate between the two terms.<sup>40</sup> Despite difficulty with this definition, the authors feel that this should not bias the result of the meta-analysis as the study by Garg et al. had a very large patient cohort compared with the study by Jones et al.<sup>27,30</sup> This was further confirmed by the EUROSTAR registry,<sup>41</sup> which showed significantly fewer endoleaks, graft migrations, and transfemoral secondary interventions in non-compliant patients.

A further limitation is the lack of adjustment for significant confounders across all outcomes. Elective and emergency cases of AAA repair could not be distinguished to statistically assess the impact on survival and re-intervention rate. However, this may not have biased the results as the population based study by Schanzer et al. and the Medicare population study by Garg et al. showed that patients who had emergency rupture were more likely to be non-compliant.<sup>8,10</sup> Although the authors are unaware of whether emergency AAA repairs were included in two studies,<sup>20,40</sup> the study by Leurs et al. and Hicks et al., with large cohorts of patients, had only elective cases,<sup>16,41</sup> whereas the compliant and non-compliant groups in the study by Wu et al. had equal proportions of emergency AAA

**Table 6.** Comparative meta-analyses of re-intervention.

Time point (y)	OR	95% CI lower	95% CI upper	p-value	I <sup>2</sup>	τ <sup>2</sup>	Q	p value (Q)
1	6.36	0.23	172.73	0.27	88.23	7.38	13.75	<0.01
2	8.08	0.25	262.57	0.24	91.66	8.48	14.61	<0.01
3	3.94	1.46	10.69	0.01	67.96	0.51	7.00	0.03
4	4.37	1.31	14.57	0.02	74.22	0.80	9.25	0.01
5	5.34	1.87	15.29	<0.01	57.80	0.50	4.84	0.09

Note. Odds ratio (OR) > 1 indicates that the average intervention rate is higher in the compliant patients. CI = confidence interval.



**Table 7.** Aneurysm related mortality (ARM), overall data, and definitions of ARM.

Study	Compliant patients		Non-compliant patients		Definition	Length of follow-up (y)
	Total number of patients (n)	Mortality (n)	Total number of patients (n)	Mortality (n)		
Garg et al. <sup>19</sup>	3944	24	3944	13	"Aneurysm related in hospital mortality"	10
Sarangarm et al. <sup>42</sup>	107	3	19	0	"[D]ied from post-EVAR open conversion complications"	7
Leurs et al. <sup>41</sup>	1538	21	2895	26	"[D]eaths due to aneurysm rupture, a primary or secondary procedure, or surgical conversion"	7
Waduud et al. <sup>20</sup>	301	8	247	8	"AAA related death"	5

Note. EVAR = endovascular aneurysm repair; AAA = abdominal aortic aneurysm.

**Table 8.** Random effects meta-analysis of aneurysm related mortality.

Overall OR	95% CI lower	95% CI upper	p-value	I <sup>2</sup>	τ <sup>2</sup>	Q	p-value (Q)
1.47	0.99	2.19	0.06	0.00	0	1.82	0.61

Note. Odds ratio (OR) > 1 indicates that mortality rate is higher in the compliant patients. CI = confidence interval.

repairs.<sup>21</sup> Furthermore, Garg et al. and Sarangarm et al. excluded patients who had died within 30 days and those who died within the first year, respectively.<sup>19,42</sup> Thus, the trend effect of improved survival in the non-compliant group because of fewer emergency AAAs is potentially negligible. Further to the inability to adjust for emergency procedures, it was not possible to adjust the results for other significant confounders across the outcomes. None of the studies, except the studies by Garg et al. and Hicks et al.,<sup>16,19</sup> provided analysis after propensity matching.

## CONCLUSION

This study suggests that although surveillance is associated with an increased rate of re-interventions, it does not appear to be associated with improved survival. Thus, improved evidence based surveillance programmes are urgently required.

## CONFLICT OF INTEREST

I.L. has received research grants from Medtronic Endovascular and Endologix. K.S. is a consultant advisor for Endologix. M.M.T. is the Chief Medical Officer for Endologix.

## FUNDING

Circulation Foundation Surgeon Scientist Award to A.K. P.H. is a Clinician Scientist supported financially by the National Institute for Health Research (NIHR; NIHR-CS-011-008). K.S. and B.P. are NIHR funded clinical lecturers. The NIHR had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health.

## APPENDIX A. SUPPLEMENTARY DATA

Supplementary data related to this article can be found at <https://doi.org/10.1016/j.ejvs.2017.11.030>.

## REFERENCES

- Karthikesalingam A, Vidal-Diez A, Holt PJ, Loftus IM, Schermerhorn ML, Soden PA, et al. Thresholds for abdominal aortic aneurysm repair in England and the United States. *N Engl J Med* 2016;**375**:2051–9.
- Reise JA, Sheldon H, Earnshaw J, Naylor AR, Dick F, Powell JT, et al. Patient preference for surgical method of abdominal aortic aneurysm repair: postal survey. *Eur J Vasc Endovasc Surg* 2010;**39**:55–61.
- Nordon IM, Karthikesalingam A, Hinchliffe RJ, Holt PJ, Loftus IM, Thompson MM. Secondary interventions following endovascular aneurysm repair (EVAR) and the enduring value of graft surveillance. *Eur J Vasc Endovasc Surg* 2010;**39**:547–54.
- Karthikesalingam A, Holt PJE, Hinchliffe RJ, Nordon IM, Loftus IM, Thompson MM. Risk of reintervention after endovascular aortic aneurysm repair. *Br J Surg* 2010;**97**:657–63.
- Chaikof EL, Brewster DC, Dalman RL, Makaroun MS, Illig KA, Sicard GA, et al. SVS practice guidelines for the care of patients with an abdominal aortic aneurysm: executive summary. *J Vasc Surg* 2009;**50**:880–96.
- Wyss TR, Brown LC, Powell JT, Greenhalgh RM. Rate and predictability of graft rupture after endovascular and open abdominal aortic aneurysm repair: data from the EVAR trials. *Ann Surg* 2010;**252**:805–12.
- Walker TG, Kalva SP, Yeddula K, Wicky S, Kundu S, Drescher P, et al. Clinical practice guidelines for endovascular abdominal aortic aneurysm repair: written by the standards of practice committee for the society of interventional radiology and endorsed by the cardiovascular and interventional radiological society of europe. *J Vasc Interv Radiol* 2010;**21**:1632–55.
- Schanzer A, Messina LM, Ghosh K, Simons JP, Robinson 3rd WP, Aiello FA, et al. Follow-up compliance after endovascular abdominal aortic aneurysm repair in Medicare beneficiaries. *J Vasc Surg* 2015;**61**:16–22.
- Kret MR, Azarbal AF, Mitchell EL, Liem TK, Landry GJ, Moneta GL. Compliance with long-term surveillance recommendations following endovascular aneurysm repair or type B aortic dissection. *J Vasc Surg* 2013;**58**:25–32.
- Garg T, Baker LC, Mell MW. Adherence to postoperative surveillance guidelines after endovascular aortic aneurysm repair among medicare beneficiaries. *J Vasc Surg* 2015;**61**:23–7.

- 11 Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *BMJ* 2009;**339**. b2535–b2535.
- 12 Atkins D, Best D, Briss PA, Eccles M, Falck-Ytter Y, Flottorp S, et al. Grading quality of evidence and strength of recommendations. *BMJ* 2004;**328**. 1490–1490.
- 13 Chaikof EL, Blankensteijn JD, Harris PL, White GH, Zarins CK, Bernhard VM, et al. Reporting standards for endovascular aortic aneurysm repair. *J Vasc Surg* 2002;**35**:1048–60.
- 14 Veroniki AA, Jackson D, Viechtbauer W, Bender R, Bowden J, Knapp G, et al. Methods to estimate the between-study variance and its uncertainty in meta-analysis. *Res Synth Meth* 2016;**7**:55–79.
- 15 Dias NV, Riva L, Ivancev K, Resch T, Sonesson B, Malina M. Is there a benefit of frequent CT follow-up after EVAR? *Eur J Vasc Endovasc Surg* 2009;**37**:425–30.
- 16 Hicks CW, Zarkowsky DS, Bostock IC, Stone DH, Black III JH, Eldrup-Jorgensen J, et al. Endovascular aneurysm repair patients who are lost to follow-up have worse outcomes. *J Vasc Surg* 2017;**65**:1625–35.
- 17 de Mestral C, Croxford R, Eisenberg N, Roche-Nagle G. The impact of compliance with imaging follow-up on mortality after endovascular abdominal aortic aneurysm repair: a population based cohort study. *Eur J Vasc Endovasc Surg* 2017;**54**: 315–23.
- 18 Jackson D, Turner R. Power analysis for random-effects meta-analysis. *Res Synth Meth* 2017;**8**:290–302.
- 19 Garg T, Baker LC, Mell MW. Postoperative surveillance and long-term outcomes after endovascular aneurysm repair among medicare beneficiaries. *JAMA Surg* 2015;**150**:957–63.
- 20 Waduud MA, Choong WL, Ritchie M, Williams C, Yadavali R, Lim S, et al. Endovascular aneurysm repair: is imaging surveillance robust, and does it influence long-term mortality? *Cardiovasc Intervent Radiol* 2015;**38**:33–9.
- 21 Wu CY, Chen H, Gallagher KA, Eliason JL, Rectenwald JE, Coleman DM, et al. Predictors of compliance with surveillance after endovascular aneurysm repair and comparative survival outcomes. *J Vasc Surg* 2015;**62**:27–35.
- 22 Ohki T, Veith FJ, Shaw P, Lipsitz E, Suggs WD, Wain RA, et al. Increasing incidence of midterm and long-term complications after endovascular graft repair of abdominal aortic aneurysms: a note of caution based on a 9-year experience. *Ann Surg* 2001;**234**:323–35.
- 23 Alric P, Hinchliffe RJ, Wenham PW, Whitaker SC, Chuter TAM, Hopkinson BR. Lessons learned from the long-term follow-up of a first-generation aortic stent graft. *J Vasc Surg* 2003;**37**: 367–73.
- 24 Chang RW, Goodney P, Tucker L-Y, Okuhn S, Hua H, Rhoades A, et al. Ten-year results of endovascular abdominal aortic aneurysm repair from a large multicenter registry. *J Vasc Surg* 2013;**58**:324–32.
- 25 Matsagkas M, Kouvelos G, Peroulis M, Avgos S, Arnaoutoglou E, Papa N, et al. Standard endovascular treatment of abdominal aortic aneurysms in patients with very short proximal necks using the enduring stent graft. *J Vasc Surg* 2015;**61**:9–15.
- 26 Harrison GJ, Oshin OA, Vallabhaneni SR, Brennan JA, Fisher RK, McWilliams RG, et al. Surveillance after EVAR based on duplex ultrasound and abdominal radiography. *Eur J Vasc Endovasc Surg* 2011;**42**:187–92.
- 27 Noll Jr RE, Tonnessen BH, Kim J, Money SR, Sternbergh 3rd WC. Long-term postplacement cost comparison of aneur and zenith endografts. *Ann Vasc Surg* 2008;**22**:710–5.
- 28 Mehta M, Paty PSK, Roddy SP, Taggart JB, Sternbach Y, Kreienberg PB, et al. Treatment options for delayed aaa rupture following endovascular repair. *J Vasc Surg* 2011;**53**:14–20.
- 29 Antoniou GA, Georgiadis GS, Glancz L, Delbridge M, Murray D, Smyth JV, et al. Outcomes of endovascular aneurysm repair with 2 different endograft systems with suprarenal fixation in patients with hostile infrarenal aortic anatomy. *Vasc Endovasc Surg* 2013;**47**:9–18.
- 30 Antoniou GA, Georgiadis GS, Antoniou SA, Neequaye S, Brennan JA, Torella F, et al. Late rupture of abdominal aortic aneurysm after previous endovascular repair: a systematic review and meta-analysis. *J Endovasc Ther* 2015;**22**:734–44.
- 31 Beckerman WE, Tadros RO, Faries PL, Torres M, Wengerter SP, Vouyouka AG, et al. No major difference in outcomes for endovascular aneurysm repair stent grafts placed outside of instructions for use. *J Vasc Surg* 2016;**64**:63–74.e62.
- 32 Hwang D, Park S, Kim HK, Lee JM, Huh S. Reintervention rate after open surgery and endovascular repair for nonruptured abdominal aortic aneurysms. *Ann Vasc Surg* 2017;**43**:134–43.
- 33 Mahajan A, Barber M, Cumbie T, Filardo G, Shutze Jr WP, Sass DM, et al. The impact of aneurysm morphology and anatomic characteristics on long-term survival after endovascular abdominal aortic aneurysm repair. *Ann Vasc Surg* 2016;**34**:75–83.
- 34 Png CYM, Tadros RO, Faries PL, Torres MR, Kim SY, Lookstein R, et al. The effect of age on post-evar outcomes. *Ann Vasc Surg* 2016;**35**:156–62.
- 35 Zacharias N, Warner CJ, Taggart JB, Roddy SP, Kreienberg PB, Ozsvath KJ, et al. Anatomic characteristics of abdominal aortic aneurysms presenting with delayed rupture after endovascular aneurysm repair. *J Vasc Surg* 2016;**64**:1629–32.
- 36 Roy IN, Vallabhaneni SR. Profile of secondary interventions and triggering surveillance imaging after evar. *Cardiovasc Intervent Radiol* 2016;**39**:S153.
- 37 Spanos K, Karathanos C, Athanasoulas A, Sapelstis V, Giannoukas AD. Systematic review of follow up compliance after endovascular abdominal aortic aneurysm repair. *J Cardiovasc Surg (Torino)* 2016 Nov 23 [Epub ahead of print], PMID: 27879950.
- 38 Godfrey AD, Morbi AHM, Nordon IM. Patient compliance with surveillance following elective endovascular aneurysm repair. *Cardiovasc Intervent Radiol* 2015;**38**:1130–6.
- 39 AbuRahma AF, Yacoub M, Hass SM, AbuRahma J, Mousa AY, Dean LS, et al. Compliance of postendovascular aortic aneurysm repair imaging surveillance. *J Vasc Surg* 2016;**63**:589–95.
- 40 Jones WB, Taylor SM, Kalbaugh CA, Joels CS, Blackhurst DW, Langan lii EM, et al. Lost to follow-up: a potential underappreciated limitation of endovascular aneurysm repair. *J Vasc Surg* 2007;**46**:434–40.
- 41 Leurs LJ, Laheij RJ, Buth J. What determines and are the consequences of surveillance intensity after endovascular abdominal aortic aneurysm repair? *Ann Vasc Surg* 2005;**19**: 868–75.
- 42 Sarangarm D, Knepper J, Marek J, Biggs KL, Robertson D, Langsfeld M. Post-endovascular aneurysm repair patient outcomes and follow-up are not adversely impacted by long travel distance to tertiary vascular surgery centers. *Ann Vasc Surg* 2010;**24**:1075–81.
- 43 Cohen J, Pai A, Sullivan TM, Alden P, Alexander JQ, Cragg A, et al. A dedicated surveillance program improves compliance with endovascular aortic aneurysm repair follow-up. *Ann Vasc Surg* 2017;**44**:59–66.
- 44 Karthikesalingam A, Holt PJ, Vidal-Diez A, Choke EC, Patterson BO, Thompson LJ, et al. Predicting aortic

complications after endovascular aneurysm repair. *Br J Surg* 2013;**100**:1302–11.

- 45 Bastos Gonçalves F, Baderkhan H, Verhagen HJM, Wanhainen A, Björck M, Stolker RJ, et al. Early sac shrinkage predicts a low risk of late complications after endovascular aortic aneurysm repair. *Br J Surg* 2014;**101**:802–10.
- 46 Bastos Gonçalves F, van de Luitgaarden KM, Hoeks SE, Hendriks JM, ten Raa S, Rouwet EV, et al. Adequate seal and no endoleak on the first postoperative computed tomography

angiography as criteria for no additional imaging up to 5 years after endovascular aneurysm repair. *J Vasc Surg* 2013;**57**:1503–11.

- 47 Brenner DJ, Hall EJ. Computed tomography—an increasing source of radiation exposure. *N Engl J Med* 2007;**357**:2277–84.
- 48 Solomon R, DuMouchel W. Contrast media and nephropathy. *Invest Radiol* 2006;**41**:651–60.
- 49 Jackson D, Riley RD. A refined method for multivariate meta-analysis and meta-regression. *Stat Med* 2014;**33**:541–54.

Eur J Vasc Endovasc Surg (2018) 55, 502


## COUP D'OEIL

### Asymmetrical Arterial Diameters

Wolfgang G. Mouton <sup>a,\*</sup>, André Wyss <sup>b</sup>

<sup>a</sup> Department of Surgery, Spital Thun STS AG, Thun, Switzerland

<sup>b</sup> Department of Radiology, Spital Thun STS AG, Thun, Switzerland



An 88 year old male patient presented with rest pain in his right foot. Cardiovascular risk factors included diabetes mellitus and arterial hypertension. Poliomyelitis at the age of 15 had resulted in left leg paralysis without affecting sensation or leg length. MR angiography showed that arteries of the left leg had smaller diameters than those of the right, starting at the left common iliac artery (panel A). This may be an acquired feature as the still developing artery subsequently had less muscle mass to supply. Stenoses in the right non-paralysed leg (panel B) were successfully dilated.

\* Corresponding author.

E-mail address: [wolfgang.mouton@spitalstsag.ch](mailto:wolfgang.mouton@spitalstsag.ch) (Wolfgang G. Mouton).

1078-5884/© 2017 European Society for Vascular Surgery. Published by Elsevier Ltd. All rights reserved.

<https://doi.org/10.1016/j.ejvs.2017.11.009>