

Capturing complexity in clinician case-mix: classification system development using GP and physician associate data

Mary Halter, MSc, PhD^{1*}, Louise Joly, MSc, PhD²,
Simon de Lusignan, MSc, MD, FRCGP, FBCS CITP, FHEA³, Robert L Grant, MSc⁴,
Heather Gage, MSc, PhD⁵, Vari M Drennan, MSc, PhD⁶

¹Associate Professor, Faculty of Health, Social Care & Education, Kingston University & St George's, University of London, London, UK; ²Research Fellow, Social Care Workforce Unit, King's College London, London, UK; ³Professor of Primary Care and Clinical Informatics, Department of Clinical and Experimental Medicine, University of Surrey, Guildford, UK; ⁴Honorary Research Fellow, Faculty of Health, Social Care & Education, Kingston University & St George's, University of London, London, UK; ⁵Professor of Health Economics, School of Economics, University of Surrey, Guildford, UK; ⁶Professor of Health Care and Policy Research, Faculty of Health, Social Care & Education, Kingston University & St George's, University of London, London, UK

Abstract

Background: There are limited case-mix classification systems for primary care settings which are applicable when considering the optimal clinical skill mix to provide services.

Aim: To develop a case-mix classification system (CMCS) and test its impact on analyses of patient outcomes by clinician type, using example data from physician associates' (PAs) and GPs' consultations with same-day appointment patients.

Design & setting: Secondary analysis of controlled observational data from six general practices employing PAs and six matched practices not employing PAs in England.

Method: Routinely-collected patient consultation records (PA $n = 932$, GP $n = 1154$) were used to design the CMCS (combining problem codes, disease register data, and free text); to describe the case-mix; and to assess impact of statistical adjustment for the CMCS on comparison of outcomes of consultations with PAs and with GPs.

Results: A CMCS was developed by extending a system that only classified 18.6% (213/1147) of the presenting problems in this study's data. The CMCS differentiated the presenting patient's level of need or complexity as: acute, chronic, minor problem or symptom, prevention, or process of care, applied hierarchically. Combination of patient and consultation-level measures resulted in a higher classification of acuity and complexity for 639 (30.6%) of patient cases in this sample than if using consultation level alone. The CMCS was a key adjustment in modelling the study's main outcome measure, that is rate of repeat consultation.

Conclusion: This CMCS assisted in classifying the differences in case-mix between professions, thereby allowing fairer assessment of the potential for role substitution and task shifting in primary care, but it requires further validation.

*For correspondence:

maryhalter@sgul.kingston.ac.uk

Received: 18 April 2017

Accepted: 14 August 2017

Published: 10 January 2018

© This article is Open Access: CC BY-NC 4.0 license (<https://creativecommons.org/licenses/by-nc/4.0/>)

Author Keywords: classification, methods, case-mix, general practice, physician assistants, physician associate

Copyright © 2018, The Authors;

DOI:10.3399/

bjgpopen18X101277

Box 1. Definitions of terms used within the case-mix classification system (CMCS)

Term	Definition used in the CMCS
Acuity	Severity, intensity, and immediacy of care required for a presenting condition in a patient.
Complexity	Interaction between existing conditions and the condition the patient presents with.
Hierarchy	Ranking by relative status of acuity and complexity.
Categorisation	Assigning each presenting condition to a category defined by acuity.
Classification	Assigning each person into a hierarchical system according to their condition category or categories.
Case-mix	The mix of patients, according to their classification.

Introduction

Primary health care has a pivotal role in the delivery of health care internationally¹ and understanding who consults, for what reasons, is important for epidemiology, research, planning, reporting, and quality improvement, as well as for potential use in funding models,² including decisions about optimal professional skill mix. Increasingly, health professionals other than doctors are being employed to address problems of medical staff shortages and rising patient demand.^{3–5} In the UK, this shift is exemplified in the deployment of physician associates (PAs, known outside the UK as physician assistants) in general practice to provide consultations, mainly seeing patients booked into same-day or urgent appointments,⁶ a role similar that of PAs in the US.⁷ Health service planners and other decision makers require evidence as to the effectiveness, safety, efficiency, and acceptability of substituting for doctors, and any valid comparison requires understanding of case-mix.

The International Classification of Primary Care (ICPC) was developed to address the perceived inadequacies of the International Classification of Diseases (ICD)⁸ in describing primary care practice.⁹ The ICPC is well-established in a minority of countries,^{9–11} its unique feature being its biaxial structure, which includes reason for encounter as well as morbidity. Like other classification systems,¹² the purpose is to describe patient complexity of morbidity (and, in the case of adjusted clinical groups, predicted health resource use¹²) over a period of time.⁹ Complexity in patients¹³ is a contested concept,¹⁴ and there are some arguments that certain classification systems, such as the ambulatory care groups (ACGs),¹⁵ are limited by defining chronicity as the likelihood of recurrence of the diagnosis¹⁶ and do not translate fully into other cultural settings.¹⁷ Classification methodologies that identify acuity (the level of severity of an illness and immediacy of requirements for care¹⁸) and the complexity of the patient as a whole, are rare. Where they are used,^{19,20} greater transparency in the reporting of case definitions is argued for.²¹

De Jong *et al*²² presented a general practice classification system, developed using ICPC coding data extracted from routinely-recorded electronic medical records, clustered into chronic and oncological diseases (using Knottnerus *et al*²³), or minor illnesses and acute diseases (using the judgments of five experienced GPs). This approach met the needs of this study conceptually (investigating the substitution of doctors by PAs in same-day or urgent consultations in primary care in England²⁴); however, only 18.6% of the condition descriptors in the study data could be mapped to those listed in De Jong's study as acute or minor.²² In addition, those listed by De Jong *et al* as chronic²² included conditions that the authors would not have expected to receive such a classification (for example, acute myocardial infarction or suicide attempt). This study was also intended to capture the complexity of the patient in primary care, albeit complexity lacking a single definition,^{14,25} as this has been considered an explanation for high intra-GP variability in patient management.²⁶ Although full complexity of clinical practice cannot be reflected in patient-centred risk adjustments,¹⁶ this study sought to address complexity (defined as elements being entwined or interwoven into one system)²⁷ by combining the acuity of the patient condition at the urgent appointment with any comorbid chronic conditions into one patient classification descriptor.

Box 2. Primary care case-mix and complexity index

Case-mix and complexity index classification	Definition of the classification	Definition of the classification for each patient
Acute	1. Recent or rapid onset and of short duration (<4 weeks), such as acute pain. 2. Serious, sometimes requiring immediate intervention, such as acute abdomen. ⁵²	≥1 problem the patient presents with is classified as acute by de Jong et al ²² or by current authors.
Chronic	(synonym: longterm) As used in ICPC relating to an illness or disability of ≥6 months duration. ⁵²	Patient has no acute problems; AND patient has a chronic condition, as recorded on a disease register, (with any presenting problem) OR ≥1 problem the patient presents with is classified as chronic.
Minor problem or symptom	Minor acute illnesses include some of the commonest problems presented in general practice, such as upper respiratory tract infections or skin rashes. ⁵³	Patient has no acute problems, no record on a disease register, no chronic problems, AND ≥1 problem the patient presents with is classified as a minor problem or symptom by de Jong et al ²² or the current authors.
Prevention	Action to avoid occurrence or development of a health problem and/or its complications. Can be divided into four categories: <ul style="list-style-type: none"> • Primary prevention: action taken to avoid or remove the cause of a health problem in an individual or a population before it arises. Includes health promotion and specific protection, such as immunisation. • Secondary prevention: action taken to detect a health problem at an early stage in an individual or a population, thereby facilitating cure, or reducing or preventing it spreading, or reducing or preventing its long-term effects (for example, methods, screening, case finding, and early diagnosis). • Tertiary prevention: action taken to reduce the chronic effects of a health problem in an individual or a population by minimising the functional impairment consequent to the acute or chronic health problem (for example, prevent complications) diabetes). Includes rehabilitation. • Quaternary prevention: action taken to identify patient at risk of overmedicalisation, to protect them from new medical invasion, and to suggest to them interventions, which are ethically acceptable.⁵² 	Patient has no acute problems, no record on a disease register, no chronic problems, no minor problems or symptoms, AND ≥1 problem the patient presents with is classified as prevention.
Process of care	(synonym: procedure) In medical care, constitutes the actions undertaken by a physician. ⁵²	Patient has no acute problems, no record on a disease register, no chronic problems, no minor symptoms or problems, and no prevention problems, AND ≥1 problem the patient presents with is classified as a process of care.

Table 1. Listing of presenting condition classification and disease register combinations forming patients' case-mix classification

Patient case-mix classification	Condition category	Disease register	Total, n	GP, n	PA, n	Change in hierarchy from condition category or categories to patient-level classification	
Acute	Acute	No	31	19	12	No	
	Acute	Yes	27	17	10		
	Acute + chronic	No	1	0	1		
	Acute + chronic	Yes	3	2	1		
	Acute + chronic + minor	No	0	0	0		
	Acute + chronic + minor	Yes	5	3	2		
	Acute + minor	No	13	9	4		
	Acute + minor	Yes	15	11	4		
Subtotal acute			95	61	34		
Chronic	Chronic	No	23	14	9	No	
	Chronic	Yes	69	62	7		
	Chronic + minor	No	19	10	9		
	Chronic + minor	Yes	56	43	13		
	Chronic + process	Yes	1	1	0		
	Chronic + minor + process	Yes	2	2	0		
	Minor	Yes	608	355	253		Yes (n = 639)
	Minor + process	Yes	9	4	9		
	Prevention	Yes	3	3	0		
	Process	Yes	19	10	9		
Subtotal chronic			809				
Minor	Minor	No	1146	571	575	No	
	Minor + process	No	14	5	9		
	Minor + prevention	No	3	0	3		
Subtotal minor			1165				
Prevention	Prevention	No	4	3	1	No	
Process	Process	No	13	7	6	No	

In the US, PAs have a long history and there are large numbers working in primary care.²⁸ The role has been more recently introduced into the UK²⁹ and other countries.³⁰⁻³³ While this introduction has largely been in isolated developments,³⁰ this is changing rapidly.³⁴ Numerous studies describe the types of patient consultations that PAs undertake in primary care but presentation of clearly comparative data on case-mix is less common.³⁵ In this study, a picture emerges of either selected consultations for PAs,³⁶ or of doctors attending more chronic^{37,38} or more serious problems³⁹ while PAs attend more unspecified conditions.^{40,41} In this research^{24,42} it was therefore anticipated that there would be difference in the case-mix of PAs and GPs,⁶ and that this could confound the analyses.⁴³

This study reports how a revised classification system was built that addressed complexity in the general practice patient case data, and how the impact of this on the analyses of outcomes of care delivered by GPs and PAs was tested (reported elsewhere^{24,42}). The overall aim was to develop a method to describe what clinicians do in a way that would allow different types of clinicians to be meaningfully compared.

Method

- **Design:** CMCS development and testing, using example patient data from a controlled observational study.
- **Setting:** Data were collected from 12 volunteer general practices in England, six employing a PA and six not, matched by health economy and patient population size, during two study reference periods in 2011–2012.^{24,42}
- **Participant data:** The study included 2086 routinely-collected electronic consultation records of patients who attended a same-day PA ($n = 932$) or GP ($n = 1154$) appointment (the predominant appointment type for PAs at the time²⁵) in the study periods. Data were extracted on condition, as documented in free text and/or in a coding system (the majority of the practices used Read Codes⁴⁴). Data were also extracted on the number and type of chronic conditions the patient had, as recorded on a disease register (that is, the UK primary care pay-for-performance [P4P] Quality and Outcomes Framework⁴⁵).
- **Intervention:** The intervention involved the development of the primary care CMCS and the testing of its impact on the analysis of outcomes of consultation with different types of clinicians, in this case PAs and GPs. The definition of terms used is shown in **Box 1**.

Development of a primary care CMCS

Concepts and their definitions were discussed by the research team, and the concept of hierarchy evolved during in-person and email discussion, continued until consensus was reached. Following this, each patient condition in the dataset was searched for by LJ and MH in the lists De Jong *et al* classified as acute or minor,²² and those with a direct match to a diagnosis were assigned De Jong *et al*'s categorisation. All remaining conditions in the dataset were assigned to one of De Jong *et al*'s categories, or to categories further defined by the GP author (SdeL); these decisions were based on the range of patient conditions in the data, and SdeL's experience in chairing international primary care medical informatics groups and multiple publications on this issue.⁴⁶⁻⁵⁰ As a second stage, each patient consultation — frequently containing more than one condition in one category, or conditions in different categories — was placed into a hierarchy that allowed the patient to move up the hierarchy of complexity in the presence of one presenting condition in a higher (more acute) category, or the registration of a patient on a P4P⁴⁵ chronic conditions register.

Testing the impact of the CMCS on analyses of PA and GP outcomes

The amount of movement was described from the categorisation of a first condition to the overall classification of that patient. Analyses of the odds or rate ratio for the primary outcome (rate of reconsultation for the same problem within 14 days) and process outcomes (advice given, and numbers of diagnostic tests, referrals, prescriptions, and procedures) for PAs against the reference of GPs were carried out under three alternative adjustment models: case-mix only; other predictor or confounding variables only; and case-mix with other predictors or confounders (the fully adjusted model). The impact of each of these adjustments was assessed using generalised estimating equation models (GEE).⁵¹ The correlation matrix was set to be exchangeable; that is, patients were assumed to have some shared characteristics within each practice, whether from demographics or organisation of the practice. Cases with missing data on any one of the variables entered into each of the GEEs were excluded for analysis of that outcome. Analyses were carried out using IBM SPSS (version 22).

The study's anticipated outcomes were a developed primary care case-mix classification system and an assessment of the impact of this on analysis of processes and outcomes of care of different clinicians, using the example of PAs and GPs.

Results

The resultant structure of the primary care CMCS

The classification developed contained five categories: acute, chronic, minor problem or symptom, prevention, and process of care. These were considered in a hierarchy of medical acuity, starting with acute through to a process of care, as shown in **Box 2**.

Placing the example presenting condition data into a category

The 2086 patients in the study presented with 1147 condition descriptors. Examples of the categories applied are as follows: an 'alcohol withdrawal-induced seizure' or 'hyperemesis gravidarum' was classified as acute; an 'asthma review' or 'multiple sclerosis' was classified as chronic; 'productive cough' or 'diarrhoea and vomiting' as a minor problem or symptom; 'HIV screening' or 'smoking cessation' as prevention; and 'removal of suture from skin' or 'solicitor's report' as a process of care. A full listing of condition descriptors and their categorisation is available from the authors on request.

Placing the example patients' data into the CMCS: changes to classification level associated with the measure of patient complexity

Three-quarters ($n = 1548$) of the patients in the study presented with only one condition; 38.3% of these ($n = 593$) had a condition co-terminous with one described as chronic. Of the quarter ($n = 538$) of the patients presenting with >1 condition, 397 (73.8%) of these presented with ≥ 2 conditions that all received the same individual category (for example, four minor problems or symptoms presentations), but 133 (33.5%) of these patients also presented with a chronic condition. The remainder ($n = 141$) of those presenting with >1 condition did so with problems that received differing categorisations; of these 91 (64.5%) also presented with a chronic condition. The classification afforded, according to the category or categories of the patient's urgent and chronic conditions in combination, is shown in *Table 1*.

To exemplify the categories in combination, the patient who presented for 'stiff neck' (categorised as a minor problem or symptom) and 'anxiety states' (categorised as acute) was classified in complexity as acute. The patient presenting with 'dyspepsia' (categorised as a minor problem or symptom) and also recorded as being on the chronic kidney disease register was classified as chronic. This patient-level classification resulted in a higher classification of acuity and complexity for 639 (30.6%) of the patient cases in this sample; of these cases, 58.2% ($n = 372$) had consulted a GP and 41.8% ($n = 267$) a PA. Data for each patient case are available from the authors on request.

Testing the impact of the classification system on analyses of the processes and outcomes of care by different types of clinician, using the PA and GP example

A small number of cases had missing data on variables added to the analysis; the sample size used for each analysis was therefore $n = 929$ for PA consultation and $n = 1148$ for GP consultation.

The case-mix of the two clinician types differed in this study, and the CMCS was therefore used in all analyses to ensure that case-mix differences were adjusted for in any comparison.^{24,42} The results presented in *Table 2* show three sets of analyses: unadjusted; adjusted solely for CMCS; and adjusted for CMCS and other relevant variables (age, Index of Multiple Deprivation [IMD], and number of visits in the previous 3 months). The analyses use the three main classifications: acute, chronic and minor, excluding process and procedure classifications due to small numbers involved.

Adjusting for CMCS changed differences between the two groups of clinicians notably from the crude estimates, as in the case of 'advice on medication management'. Other measures, such as the 'number of referrals', were changed little by CMCS but notably by other demographic variables. For the primary outcome of re-consultation rate, the adjustment for CMCS alone resulted in only a small change in the estimate, while the larger change was associated with a combination of the CMCS and other variables (in this case: age, IMD, and number of visits in the previous 3 months).

Discussion

Summary

This study developed a classification aiming to address complexity in general practice by combining severity and medical immediacy of the urgent condition with information about any chronic conditions. A greater proportion of cases for one clinician type (GP rather than PA cases here) were classified as complex as a result of this, and making allowance for this complexity (rather than for urgent condition alone) resulted in some changes in the estimates of difference between the two groups.^{52,53} The classification system is replicable, using the detailed list of conditions included in each category, the hierarchy, and examples from the full dataset.

Table 2. Crude, CMCS-adjusted, other variable-adjusted and CMCS and other variable-adjusted odds ratios or rate ratios of process and outcome measure differences between PAs and GPs²⁴

Process or outcome measure (as defined elsewhere ²⁴)	Crude/univariate finding		OR or RR (95% CI)				
	GP (reference group)	PA	OR or RR	Unadjusted	Adjusted for CMCS only	Adjusted for other variables of relevance ^a , but not CMCS	Fully adjusted: CMCS and other variables of relevance
Processes	% cases						
General advice	22.9	51.4	OR	3.56 (2.58 to 4.29)	3.58 (1.82 to 7.01)	3.30 (1.68 to 6.47)	3.30 (1.69 to 6.46)
Advice on medication management	12.6	17.1	OR	1.43 (1.12 to 1.82)	^b	1.62 (1.05 to 2.49)	1.72 (1.08 to 2.72)
Advice on over the counter medication	9.5	20.5	OR	2.45 (1.92 to 3.18)	^b	6.63 (0.56 to 4.69)	1.74 (0.62 to 4.89)
	Mean per case						
Number of diagnostic tests	0.34	0.36	RR	1.06 (0.82 to 1.38)	1.11 (0.86 to 1.43)	1.07 (0.90 to 1.29)	1.08 (0.89 to 1.30)
Number of referrals	0.11	0.9	RR	0.84 (0.57 to 1.22)	^b	0.94 (0.63 to 1.41)	0.95 (0.63 to 1.43)
Number of prescriptions	0.78	0.89	RR	1.18 (0.86 to 1.58)	1.17 (0.88 to 1.55)	1.16 (0.86 to 1.56)	1.16 (0.87 to 1.53)
Number of procedures	0.1	0.1	RR	0.76 (0.32 to 1.84)	^c	^c	^c
Outcome							
Reconsultation for the same or a linked problem at the practice or urgent care facility within 14 days of the index consultation	0.29	0.32	RR	1.03 (0.75 to 1.43)	1.09 (0.77 to 1.54)	1.12 (0.81 to 1.54)	1.25 (0.91 to 1.72)

^aWith an independent association with outcome (see full report for detail).²⁴ ^bCould not be estimated in SPSS. ^cNumbers too small for further adjustment. CMCS = case-mix classification system. OR = odds ratio. RR = risk ratio.

Strengths and limitations

The disease register measure selected for this study as a measure of comorbidity was part of a P4P scheme.⁴⁵ While P4P is potentially effective in changing physician behaviour, it may lead to some gaming and focus on some activities at the expense of others.^{54,55} It also takes a single disease-based approach.⁵⁶ This study may therefore have failed to fully represent complexity. It also may not have been able to distinguish levels of acuity within, for example, conditions such as abdominal pain. However, a measure of complexity has been achieved beyond that previously used in research on this topic, and this has been done using not only system codes but also clinicians' free text descriptors. The authors acknowledge that, in using the judgments of one family physician (as opposed to a panel of five by De Jong *et al*²²) issues of potential differences of clinical opinion, or of bias, have not been addressed.

Comparison with existing literature

This study set out to both develop a classification system and test its impact on analysis, and therefore has two sets of interpretations. It is recognised that any classification system is reductionist and use of the biopsychosocial model including the patient's reason for encounter may be preferable.⁵⁶ Classification systems are also critiqued for failing to capture complexity and the undifferentiated symptoms encountered commonly in primary care;⁵⁶ to have struggled with the concept of symptoms at all;⁵⁷ and to change identity through the fact of classification.⁵⁸ Research using any coded information from primary care databases, if used without free text, can miss cases of relevance.⁵⁹ In using free text to inform this classification system, alongside coded information, this study aims to lay a foundation for more accurate comparison of the workload among the primary care workforce.

In terms of the application of the classification system to analyses of groups of clinicians, similar PA and GP case-mix difference has been reported elsewhere.³⁷⁻⁴⁰ However, comparison is limited by the implicit rather than explicit use of terms such as 'minor' or 'chronic' in previously published studies, and the lack of adjustment for these case-mix differences in analyses of outcome.

Implications for research

Task-shifting includes the creation of new professional groups, whereby tasks are shifted from workers with more general training to workers with specific training for a particular task.⁶⁰ However, PAs are trained as generalists rather than specialists,⁶¹ and may be seen as an example of where task shifting can be a promising policy option to increase the productive efficiency of the delivery of healthcare services.⁶⁰ However, a records-based analysis is required to measure the impact of skill-mix on task-shifting in primary care, and such analyses should include careful statistical adjustment based on explicit rather than implicit definitions of case-mix classification. The system proposed in this study requires further testing, refinement, and validation of the classification in other studies.

In conclusion, this study demonstrated the development of a classification of complexity as well as the difference that applying such classification to analyses makes to results of comparisons of different professionals working in the same system. In particular, the classification allows analyses to account for greater complexity in the patients seen by the GPs. Policymakers, workforce planners, and employers of PAs, or any other clinician substitutes, will need to consider these differences both when considering task allocation and when evaluating efficiency (including cost) and effectiveness. Researchers should be encouraged to consider further developing and evaluating the classification in other studies.

Funding

This project was funded by the National Institute for Health Research (NIHR) Health Services and Delivery Research (HS&DR) programme (project number 09/1801/1066). The views and opinions expressed herein are those of the authors and do not necessarily reflect those of the HS&DR Programme, NIHR, the NHS, or the Department of Health.

Service support costs, enabling general practices to allocate staff time to the collection of data from patient consultation records, were received from the UK Comprehensive Research Networks covering the four study regions.

Ethics approval

The study was approved by the Surrey NHS Research Ethics Committee (reference 10/H1109/28).

Provenance

Freely submitted; externally peer reviewed.

Acknowledgements

The authors acknowledge the contribution of the article's reviewers.

References

1. World Health Organization. Primary care: putting people first In. *World Health Report 2008: Primary Health Care — Now More than Ever*. Geneva: World Health Organization, 2008.
2. National Casemix and Classification Centre, University of Woolongong. Comments on the document "Activity based funding for Australian public hospitals: towards a pricing framework". <http://ahsri.uow.edu.au/content/groups/public/@web/@chsd/documents/doc/uow119700.pdf> (accessed 6 Dec 2017).
3. The Joint Commission, with support from Aramark. *Health care at the crossroads: guiding principles for the development of the hospital of the future*. https://www.jointcommission.org/assets/1/18/Hospital_Future.pdf. (accessed 6 Dec 2017).
4. Health and Education National Strategic Exchange. Review of Medical and Dental School Intakes in England. https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/213236/medical-and-dental-school-intakes.pdf (accessed 6 Dec 2017).
5. The Connexion. Stats reveal truth on GP shortage. <http://www.connexionfrance.com/statistics-reveal-truth-on-shortage-of-french-doctors-11594-news-article.html>. (accessed 6 Dec 2017).
6. Drennan V, Levenson R, Halter M, et al. Physician assistants in English general practice: a qualitative study of employers' viewpoints. *J Health Serv Res Policy*. doi: 10.1258/jhsrp.2010.010061

7. Morgan PA, Abbott DH, McNeil RB, et al. Characteristics of primary care office visits to nurse practitioners, physician assistants and physicians in United States Veterans Health Administration facilities, 2005 to 2010: a retrospective cross-sectional analysis. *Hum Resour Health* 2012; **10**: 1. doi: 10.1186/1478-4491-10-42
8. World Health Organization. International Classification of Primary Care (ICPC-2). Geneva. World Health Organization. 2003.
9. Soler JK, Okkes I, Wood M, et al. The coming of age of ICPC: celebrating the 21st birthday of the International Classification of Primary Care. *Fam Pract* 2008; **25**(4): 312–317. doi: 10.1093/fampra/cmn028
10. Australian Institute of Health and Welfare. *Australia's Health 2004*. Canberra: Australian Institute of Health and Welfare, 2004.
11. de Lusignan S. Codes, classifications, terminologies and nomenclatures: definition, development and application in practice. *Inform Prim Care* 2005; **13**(1): 65–69. doi: 10.14236/jhi.v13i1.580
12. Starfield B, Lemke K. Complexity of patient illness versus complexity of patient visits. http://www.annfamned.org/content/8/4/291.long/reply#annalsfm_el_13780 (accessed 19 Dec 2017).
13. Safford MM, Allison JJ, Kiefe CI. Patient complexity: more than comorbidity. the vector model of complexity. *J Gen Intern Med* 2007; **22**(3): 382–390. doi: 10.1007/s11606-007-0307-0
14. Green LA. The implications of measuring complexity. *Ann Fam Med* 2010; **8**(4): 291–292. doi: 10.1370/afm.1147
15. Weiner JP, Starfield BH, Lieberman RN. Johns Hopkins Ambulatory Care Groups (ACGs). A case-mix system for UR, QA and capitation adjustment. *HMO Pract* 1992; **6**(1): 13–19.
16. Katerndahl DA, Wood R, Jaén CR. A method for estimating relative complexity of ambulatory care. *Ann Fam Med* 2010; **8**(4): 341–347. doi: 10.1370/afm.1157
17. Halling A, Fridh G, Ovhed I. Validating the Johns Hopkins ACG Case-Mix System of the elderly in Swedish primary health care. *BMC Public Health* 2006; **6**(1): 1. doi: 10.1186/1471-2458-6-171
18. Brennan CW, Daly BJ. Patient acuity: a concept analysis. *J Adv Nurs* 2009; **65**(5): 1114–1126. doi: 10.1111/j.1365-2648.2008.04920.x
19. Everett CM, Thorpe CT, Palta M, et al. Division of primary care services between physicians, physician assistants, and nurse practitioners for older patients with diabetes. *Med Care Res Rev* 2013; **70**(5): 531–541. doi: 10.1177/1077558713495453
20. Parkerson GR, Michener JL, Yarnall KS, et al. Duke Case-Mix System (DUMIX) for ambulatory health care. *J Clin Epidemiol* 1997; **50**(12): 1385–1394. doi: 10.1016/S0895-4356(97)00217-5
21. Bhattarai N, Charlton J, Rudisill C, et al. Coding, recording and incidence of different forms of coronary heart disease in primary care. *PLoS ONE* 2012; **7**(1): e29776. doi: 10.1371/journal.pone.0029776
22. De Jong J, Visser MR, Wieringa-de Waard M. Exploring differences in patient mix in a cohort of GP trainees and their trainers. *BMJ Open* 2011; **1**(2): e000318. doi: 10.1136/bmjopen-2011-000318
23. Knottnerus JA, Metsemakers J, Höppener P, et al. Chronic illness in the community and the concept of 'social prevalence'. *Fam Pract* 1992; **9**(1): 15–21. doi: 10.1093/fampra/9.1.15
24. Drennan VM, Halter M, Brearley S, et al. Investigating the contribution of physician assistants to primary care in England: a mixed-methods study. *Health Services and Delivery Research* 2014; **2**(16): 1–136. doi: 10.3310/hsdr02160
25. Peek CJ, Baird MA, Coleman E. Primary care for patient complexity, not only disease. *Fam Syst Health* 2009; **27**(4): 287–302. doi: 10.1037/a0018048
26. Rethans JJ, Saebu L. Do general practitioners act consistently in real practice when they meet the same patient twice? Examination of intradoctor variation using standardised (simulated) patients. *BMJ* 1997; **314** (7088): 1170–1173. doi: 10.1136/bmj.314.7088.1170
27. Sturmberg JP, Martin CM, Katerndahl DA. It is complicated! Misunderstanding the complexities of 'complex'. *J Eval Clin Pract* 2017; **23**(2): 426–429. doi: 10.1111/jep.12579
28. American Academy of Physician Assistants. Physician assistant census report: results from the 2010 AAPA Census. <http://www.aapa.org/workarea/downloadasset.aspx?id=1454> (accessed 6 Dec 2017).
29. Drennan VM, Chattopadhyay K, Halter M, et al. Physician assistants in English primary care teams: a survey. *J Interprof Care* 2012; **26**(5): 416–418. doi: 10.3109/13561820.2012.686538
30. Frossard LA, Liebich G, Hooker RS, et al. Introducing physician assistants into new roles: international experiences. *Med J Aust* 2008; **188**(4): 199–201.
31. Jones IW, Hooker RS. Physician assistants in Canada: update on health policy initiatives. *Can Fam Physician* 2011; **57**(3): e83–e88.
32. Nederlandse Associatie Physician Assistants [Dutch Association of Physician Assistants]. English Information. <http://www.napa.nl/kennisbank/english/> (accessed 14 Dec 2017)
33. O'Connor TM, Hooker RS. Extending rural and remote medicine with a new type of health worker: physician assistants. *Aust J Rural Health* 2007; **15**(6): 346–351. doi: 10.1111/j.1440-1584.2007.00926.x
34. Gallagher J, BBC News. NHS plans rapid expansion of 'doctor's assistant' jobs. <http://www.bbc.co.uk/news/health-28896625> (accessed 6 Dec 2017).
35. Halter M, Drennan V, Chattopadhyay K, et al. The contribution of physician assistants in primary care: a systematic review. *BMC Health Serv Res* 2013; **13**: 223. doi: 10.1186/1472-6963-13-223
36. Fethke C, Ekwo E, Daniels M. Management practices: task allocation between physicians and physician assistants. *J Ambul Care Manage* 1979; **2**(2): 31–45.
37. Nelson EC, Johnson KG, Jacobs AR. Impact of medex on physician activities: redistribution of physician time after incorporating a medex into the practice. *J Fam Pract* 1977; **5**(4): 607–612.

38. Grzybicki DM, Sullivan PJ, Oppy JM, et al. The economic benefit for family/general medicine practices employing physician assistants. *Am J Manag Care* 2002; **8(7)**: 613–662.
39. Duttera MJ, Harlan WR. Evaluation of physician assistants in rural primary care. *Arch Intern Med* 1978; **138(2)**: 224–22. doi: 10.1001/archinte.1978.03630260036013
40. Parle JV, Ross NM, Doe WF. The medical care practitioner: developing a physician assistant equivalent for the United Kingdom. *Med J Aust* 2006; **185(1)**: 13–17.
41. Simkens ABM, van Baar ME, van Balen FAM, et al. The Physician Assistant in General Practice in the Netherlands. *J Physician Assist Educ* 2009; **20(1)**: 30–38. doi: 10.1097/01367895-200920010-00005
42. Drennan V, Halter M, Joly L, et al. Physicians Assistants and General Practitioners: a comparison of same day appointment consultations. *Br J Gen Pract* 2015; **65(634)**: e344–e350. doi: 10.3399/bjgp15X684877
43. Dean BB, Lam J, Natoli JL, et al. Review: use of electronic medical records for health outcomes research: a literature review. *Med Care Res Rev* 2009; **66(6)**: 611–638. doi: 10.1177/1077558709332440
44. NHS Digital. Read Codes. <https://digital.nhs.uk/article/1104/Read-Codes> (accessed 21 Dec 2017).
45. NHS Employers. 2014/15 General Medical Services (GMS) Contract Quality and Outcomes Framework (QOF): guidance for GMS contract 2014/15. <http://www.nhsemployers.org/~media/Employers/Publications/2014%2015%20QOF%20guidance%20stakeholders.pdf> (accessed 6 Dec 2017)
46. de Lusignan S. Codes, classifications, terminologies and nomenclatures: definition, development and application in practice. *Inform Prim Care* 2005; **13(1)**: 65–69. doi: 10.14236/jhi.v13i1.580
47. de Lusignan S, van Weel C. The use of routinely collected computer data for research in primary care: opportunities and challenges. *Fam Pract* 2006; **23(2)**: 253–263. doi: 10.1093/fampra/cmi106
48. de Lusignan S, Liaw ST, Michalakidis G, et al. Defining datasets and creating data dictionaries for quality improvement and research in chronic disease using routinely collected data: an ontology-driven approach. *Inform Prim Care* 2011; **19(3)**: 127–134. doi: 10.14236/jhi.v19i3.805
49. de Lusignan S, Chan T, Jones S. Large complex terminologies: more coding choice, but harder to find data — reflections on introduction of SNOMED CT (systematized nomenclature of medicine — clinical terms) as an NHS standard. *Inform Prim Care* 2011; **19(1)**: 3–5. doi: 10.14236/jhi.v19i1.787
50. de Lusignan S. In this issue: ontologies a key concept in informatics and key for open definitions of cases, exposures, and outcome measures. *J Innov Health Inform* 2015; **22(2)**: 170170. doi: 10.14236/jhi.v22i2.170
51. Snijders TAB, Bosker RJ. *Multilevel analysis: an introduction to basic and advanced multilevel modeling*. 2nd ed. London: Sage Publications Ltd, 2012.
52. Bentzen N. Preface in *Wonca Dictionary for General/Family Practice*. Copenhagen: WONCA International Classification Committee, 2003. http://www.ph3c.org/4daction/w3_CatVisu/en/wonca-dictionary-of-general-family-practice.-2003.html?wDocID=92 (accessed 6 Dec 2017).
53. Jones R, White P, Armstrong D. et al. Managing acute illness. http://www.kingsfund.org.uk/sites/files/kf/field/field_document/managing-acute-illness-gp-inquiry-research-paper-mar11.pdf (accessed 6 Dec 2017).
54. Dixon A, Khachatryan A, Wallace A, et al. *The Quality and Outcomes Framework (QOF): does it reduce health inequalities?* Southampton: National Institute of Health Research: Queen's Printer and Controller of HMSO, 2011. http://www.netscc.ac.uk/hsdr/files/project/SDO_FR_08-1716-207_V01.pdf (accessed 22 Dec 2017).
55. Smith PC, Mossialos E, Papanicolas I. et al. Performance measurement for health system improvement. Experiences, challenges and prospects. Copenhagen: World Health Organization 2008 and World Health Organization, on behalf of the European Observatory on Health Systems and Policies. <http://www.who.int/management/district/performance/PerformanceMeasurementHealthSystemImprovement2.pdf> (accessed 6 Dec 2017).
56. Gillam S. Should the quality and outcomes framework be abolished? Yes. *BMJ* 2010; **340**: c2710. doi: 10.1136/bmj.c2710
57. Soler JK, Okkes I. Reasons for encounter and symptom diagnoses: a superior description of patients' problems in contrast to medically unexplained symptoms (MUS). *Fam Pract* 2012; **29(3)**: 272–282. doi: 10.1093/fampra/cmr101
58. Armstrong D. Diagnosis and nosology in primary care. *Soc Sci Med* 2011; **73(6)**: 801–807. doi: 10.1016/j.socscimed.2011.05.017
59. Ford E, Nicholson A, Koeling R, et al. Optimising the use of electronic health records to estimate the incidence of rheumatoid arthritis in primary care: what information is hidden in free text? *BMC Med Res Methodol* 2013; **13**: 105. doi: 10.1186/1471-2288-13-105
60. Fulton BD, Scheffler RM, Sparkes SP, et al. Health workforce skill mix and task shifting in low income countries: a review of recent evidence. *Hum Resour Health* 2011; **9**: 1. doi: 10.1186/1478-4491-9-1
61. Miles DL, Rushing WA. A study of physicians' assistants in a rural setting. *Med Care* 1976; **15(12)**: 987–995. doi: 10.1097/00005650-197612000-00002