

Selection and T-cell antigenicity of synthetic long peptides derived from SARS-CoV-2

Katarzyna Piadel, Amin Haybatollahi, Angus George Dagleish and Peter Lawrence Smith*

Abstract

The pandemic caused by SARS-CoV-2 has led to the successful development of effective vaccines however the prospect of variants of SARS-CoV-2 and future coronavirus outbreaks necessitates the investigation of other vaccine strategies capable of broadening vaccine mediated T-cell responses and potentially providing cross-immunity. In this study the SARS-CoV-2 proteome was assessed for clusters of immunogenic epitopes restricted to diverse human leucocyte antigen. These regions were then assessed for their conservation amongst other coronaviruses representative of different alpha and beta coronavirus genera. Sixteen highly conserved peptides containing numerous HLA class I and II restricted epitopes were synthesized from these regions and assessed *in vitro* for their antigenicity against T-cells from individuals with previous SARS-CoV-2 infection. Monocyte derived dendritic cells were generated from these peripheral blood mononuclear cells (PBMC), loaded with SARS-CoV-2 peptides, and used to induce autologous CD4+ and CD8+ T cell activation. The SARS-CoV-2 peptides demonstrated antigenicity against the T-cells from individuals with previous SARS-CoV-2 infection indicating that this approach holds promise as a method to activate anti-SARS-CoV-2 T-cell responses from conserved regions of the virus which are not included in vaccines utilising the Spike protein.

DATA SUMMARY

NC_045512.2, MN996532.2, MT072864.1, MT040333.1, KC881005.1, KY417144.1, KT444582.1, KF367457.1, KC881006.1, KY417152.1, MK211376.1, AY572034.1, AY572035.1, AY572038.1, AY686864.1, AY686863.1, NC_004718.3, AY390556.1, KY352407.1, NC_014470.1, GU190215.1, MG772933.1, MG772934.1, MK211374.1, JX993987.1, MK211377.1, DQ084199.1, GQ153539.1, DQ084200.1, DQ022305.2, KJ473815.1, KJ473813.1, KJ473812.1, KP886808.1, DQ648857.1, DQ412043.1, KY417147.1, DQ648856.1, JX993988.1, FJ588686.1, MK211378.1, KJ473816.1, KJ473811.1, MG596802.1, MG596803.1, NC_009020.1, MG021452.1, MK129253.1, JX869059.2, MT576585.1, NC_038294.1, DQ648794.1, KJ473821.1, KJ473822.1, KX442564.1, EF065505.1, NC_009019.1, KC869678.4, NC_039207.1, KC545386.1, KJ473806.1, NC_026011.1, NC_017083.1, FJ647223.1, KF686343.1, AY391777.1, NC_030886.1, NC_009021.1, HM211098.1, MK211375.1, KY770860.1, KU182964.1, MT430884.1, LC469308.1, NC_048212.1, MF593268.1, NC_034440.1, NC_028814.1, JQ989273.1, NC_018871.1, AF304460.1, JQ410000.1, NC_048216.1, AY567487.2, NC_046964.1, NC_028833.1, KJ473820.1, NC_028811.1, NC_022103.1, EF203064.1, NC_028824.1, DQ648858.1, EU420138.1, KJ473795.1, EU420137.1.

INTRODUCTION

The pandemic caused by the novel coronavirus, SARS-CoV-2, has prompted a global response to produce effective vaccines. A number of vaccines have been approved for use having demonstrated varying levels of efficacy in clinical trials [1]. These vaccines, often based upon targeting the viral Spike protein, responsible for facilitating entry of SARS-CoV-2 into cells [2], have demonstrated an ability to limit infection, transmission and the onset of serious disease. However, a number of issues remain. The recent identification of SARS-CoV-2 variants to which vaccine elicited immune protection may be reduced [3] raises the prospect of continued susceptibility to serious infection and the need for repeated vaccination to raise immunity to new variants

Received 10 August 2021; Accepted 17 October 2021; Published 11 January 2022

Author affiliations: ¹Institute of Infection and Immunity, St Georges University of London, London UK.

***Correspondence:** Peter Lawrence Smith, plsmith@sgul.ac.uk

Keywords: SARS-CoV-2; T-cells; peptide; vaccine.

Abbreviations: ELISPOT, enzyme-linked immunospot; mDDC, monocyte-derived dendritic cells; PBMC, peripheral blood mononuclear cells.

Three supplementary tables are available with the online version of this article.

001698 © 2022 The Authors



This is an open-access article distributed under the terms of the Creative Commons Attribution License. This article was made open access via a Publish and Read agreement between the Microbiology Society and the corresponding author's institution.

or novel coronavirus outbreaks. Additional problems related to vaccination include cost, logistics and the duration of protection afforded by neutralising antibody [4].

T-cell responses to SARS-CoV-1 demonstrate greater durability than those of neutralising antibody [5] and are associated with protection against SARS-CoV-2 [6], particularly in the context of waning antibody titres [7], indicating that T-cell mediated immunity may offer durable immune protection which may limit the severity of disease and potentially offer immune responses that are cross reactive to variant SARS-CoV-2 and other coronaviruses [6, 8, 9] similar to those have observed for different influenza viruses [10]. T-cell immune responses are generated by vaccination with the SARS-CoV-2 spike protein however these responses are thought to represent approximately 50% of the total anti-SARS-CoV-2 CD4+ T cell response and 25% of CD8+ T cell responses [11]. Therefore, spike-based vaccines will likely induce sub-optimal anti-SARS-CoV-2 T-cell responses and alternative methods of inducing T-cell immunity need to be explored. Here we provide a rationale for the selection of antigenic regions from SARS-CoV-2 proteins including the nucleoprotein, membrane protein, envelope protein, ORF3, ORF7a and the non-structural proteins intended to provide broad T-cell activation and assess these synthetic long peptides for immunogenicity *in vitro*.

METHODS

Coronavirus sequence conservation

Analysis of SARS-CoV-2 and other coronaviruses protein sequences from proteins harbouring epitope rich regions was performed using FASTA sequences deposited in the NCBI database (<https://www.ncbi.nlm.nih.gov/labs/virus/vssi/>), alignment using Clustal Omega with default settings (<https://www.ebi.ac.uk/Tools/msa/clustalo/>) and analysis of conservation using Microsoft Excel. Accession numbers of coronaviruses strains can be found in the supplementary material file.

In silico prediction of T-cell epitopes

Identification of SARS-CoV-2 HLA restricted epitopes using prediction algorithms and experimentally validated epitopes deposited in the Immune epitope database and by using the NetMHCpan EL 4.1 (2020.09) prediction algorithm against an HLA allele reference set (<https://www.iedb.org/>).

Selection and synthesis of synthetic long peptides

Selection of conserved immunogenic regions between 15–30 amino acids in length (synthetic long peptides) was determined by assessing their suitability for synthesis based upon the physiochemical properties of the amino acids in the sequence and potential as CPP (cell penetrating peptides) defined by a net positive charge. Selected peptides were synthesized (Genscript, Netherlands).

Peripheral blood mononuclear cells

Peripheral blood mononuclear cells (PBMC) were purchased from the national blood service (NHS, UK) prior to the distribution of SARS-CoV-2 vaccines. PBMC demonstrating responses to a SARS-CoV-2 consensus peptide pool and serum antibody responses to the SARS-CoV-2 spike protein were defined as having been previously infected with SARS-CoV-2 whilst PBMC and sera lacking detectable responses were defined as SARS-CoV-2 naïve.

Isolation of PBMC and generation of monocyte-derived DC

The generation of mDDC (monocyte-derived dendritic cells) was performed using established protocols. CD14+ cells were isolated by positive selection using anti-CD14 conjugated magnetic beads (Miltenyi, Germany). The CD14+ cells were cultured for 6 days in RPMI (Sigma, UK) 10% foetal calf serum (sigma, UK) and 5% streptomycin/penicillin (Sigma, UK) of 10 ng ml⁻¹ IL-4 and 50 ng ml⁻¹ GM-CSF (Miltenyi, Germany). These mDDC were subsequently co-cultured with the T-cells enriched from the CD14- fraction of PBMC using anti-CD3 magnetic beads (Miltenyi, Germany) in the presence of SARS-CoV-2 peptides individual and in groups, and control peptides including CEF (Cytomegalovirus/Epstein Barr virus and influenza), CEFT (Cytomegalovirus/Epstein Barr virus/influenza/tetanus) SARS-CoV-2 reference group (including overlapping peptides from the spike, nucleocapsid and membrane) (1 µg ml⁻¹ for each peptide) for antigenicity assays.

ELISPOT

IFN-γ ELISPOT (enzyme-linked immunosorbent spot) assays using peptide pulsed mDDC-T-cell co-cultures (2×10⁴ mDDC: 2×10⁵ T-cells) were incubated in IFN-γ ELISPOT plates (CTL Ltd, Germany) for 48 h in order to assess antigen specific T-cell activation. After 48 h ELISPOT plates were processed according to the manufacturers protocol. Briefly, plates were washed in PBS ×3 prior to the incubation with detection anti-IFN-γ antibody for 2 h. Plates were washed ×3 in PBS wash buffer and Strep-Biotin reagent was added for 1 h. Plates were washed ×3 prior to the addition of substrate solution. Spot formation was observed and the plates were washed once in distilled H₂O and left to dry before enumeration using a CTL Immunospot entry ELISPOT plate reader. Positive responses compared to the no peptide control were defined as >1.5-fold change and statistical significance using Student's t-test.

Flow cytometry

Multiparameter flow cytometry was used to measure CD8⁺ and CD4⁺ T cell activation using the AIM assay. A total of 1×10^5 MDDC were incubated with SARS-CoV-2 or control peptides and co cultured with 1×10^6 of the T-cells enriched from the CD14⁺ fraction of PBMC for 24 h in flat bottomed 96 well plates prior to staining. V450 anti-CD3, AF488 anti-CD8, APC-fire-anti-CD4, V670 anti-CD45RA and V710 anti-CCR7 were used as T-cell subset markers. APC anti-CD69, Percp anti-CD137 and PE-dazzle anti-OX40, Alexa Fluor 700-OX were used as activation induced markers. Responses to the peptides were defined as a 1.5-fold increase to the no peptide control.

RESULTS

Identification of antigenic regions within the SARS-CoV-2 proteome

In order to identify amino acid sequences within the SARS-CoV-2 proteome that contain multiple class I and II restricted epitopes, peptides from conserved regions from SARS-CoV-2 proteins were assessed using the IEDB epitope prediction tool [12]. Identification of clusters of epitopes, previously validated for HLA binding or T-cell activation and deposited within the IEDB database, were also used. This resulted in the identification of 25 peptide regions harbouring multiple predicted or experimentally validated epitopes. Five of these peptides were identified within the Spike protein and were not investigated further since T-cell responses to these regions may be raised by existing vaccines. The remaining 20 peptides were derived from the nucleoprotein, envelop protein, membrane protein, ORF3a, ORF7a and the ORF1ab polyprotein (Table 1). These peptides were assessed for conservation between 500–2500 SARS-CoV-2 protein sequences deposited on the NCBI virus database and including sequences derived from different geographic locations and belonging to variants of concern. The amino acids in each peptide were highly conserved with typically between 98–100% conservation for each aa residue within each peptide (Table S1). Some peptides demonstrated 100% conservation whilst the average conservation across all 20 peptides was 99.4 % (Table 1). The limited variation was often between similar amino acids (Table S1). Early analysis of available protein sequences from the Omicron variant of SARS-CoV-2 also demonstrated 99% conservation with the peptides identified here.

Peptides intended to induce broad antigen specific T-cell responses need to contain epitopes to the most common HLA alleles in human populations. The collective HLA restriction of the experimentally validated epitopes identified in the IEDB database in each of the 20 peptides was determined (Table 2). The 20 peptides included a total of 144 experimentally validated epitopes, 125 restricted to HLA class I and 19 restricted to HLA-class II. Next the presence of predicted, but untested, epitopes was determined (Table 3) defined as being within the top 0.1% of predicted binders for each HLA allele. Ninety-five predicted epitopes were identified, 93 restricted to HLA class I and 2 to HLA-class II. In total the epitopes identified within the peptides were restricted to the most common HLA-class I alleles including but not limited to HLA*A01 : 01, HLA*A02 : 01, HLA*A03 : 01, HLA*A11 : 01, HLA*A24 : 02, HLA*A68 : 01, HLA*A68 : 02, HLA*B07 : 02, HLA*08 : 01, HLA-B*15 : 02, HLA*B35 : 01, HLA*B40 : 01.

Significant variation exists between bat coronaviruses related to SARS-CoV-2 and between other coronaviruses known to infect humans. Conservation of the 20 SARS-CoV-2 peptides with 93 other coronaviruses was assessed (Table 4). High sequence conservation between SARS-CoV-2 and other Serbecoviruses, including SARS-CoV-1 and bat derived SARS-like viruses, was demonstrated. Peptides derived from ORF1ab demonstrated greater conservation between viruses compared to peptides derived from the structural proteins such as the nucleoprotein. High conservation was also observed within the peptides between SARS-CoV-2 and Marbecoviruses, including MERS-CoV, responsible for pathogenic human infection. Again, higher levels of conservation were observed for the ORF1ab peptides. Some conservation was seen for coronaviruses more distantly related to SARS-CoV-2 such as Embecovirus, Duvinacovirus and Setracovirus genera including the viruses OC43, NL63, 229E, HKU1 which infect humans, consistent with recent studies detecting T-cell responses against SARS-CoV-2 in uninfected individuals [13–17]. The conservation in the peptides between SARS-CoV-2 and 93 other coronaviruses was then compared to conservation within regions of the Spike protein known to be targets for neutralising antibody (Table 4). The receptor binding domain (RBD) and the N-terminal domain (NTD) of the spike protein demonstrated greater variation between SARS-CoV-2 and the 93 other coronaviruses relative to the SARS-CoV-2 peptides. For example, the majority of Serbecoviruses demonstrated 100% homology with SARS-CoV-2 in the NSP16⁶⁸²¹⁻⁴⁵² peptide. In contrast the same viruses demonstrated approximately 47% homology to the SARS-CoV-2 NTD and between 60–70% to the SARS-CoV-2 RBD. These data indicate that immune responses raised against the SARS-CoV-2 peptides identified here could mediate cross immunity against diverse coronavirus strains, including those containing spike proteins with limited homology to SARS-CoV-2.

Most HLA class I restricted epitopes consist of 8-10mer amino acid sequences. Whilst high levels of conservation in the peptides was demonstrated between Coronaviruses (Table 4) a relatively small amount of variation can significantly alter recognition by either T-cell or antibody-based immune responses, demonstrated by observations that amino acid substitutions allow immune escape from neutralising antibody, however variation within one or two amino acids within the epitope sequence may still allow for T-cell recognition, albeit sometimes with altered TCR avidity for the peptide/MHC complex. This is particularly true for conservative amino acid substitution such as isoleucine and leucine. For this reason, the conservation of the 125 validated, HLA-class I restricted epitopes identified in SARS-CoV-2 peptides were determined in each of the 20 peptides from each of the

Table 1. Immunogenic SARS-CoV-2 peptides. The peptides were identified based upon the presence of HLA class I and II restricted epitopes, sequence conservation amongst SARS coronaviruses, suitability for synthesis and positive charge at neutral pH

No	Region	Sequence	Sequence length	Hydrophobicity	Gravy	MW average G mol ⁻¹	Theoretical pI	Average aa conservation (%)
1	NP ⁴¹⁻⁵⁷	RRPQLPNNITASWFTALTQHGKEDLK	26	50.44	-0.88	2864.2423	10.1	99.0
2	NP ⁶⁵⁻⁹⁴	FPRGQVPINSGPDDQIYYRRATRRI	27	31.61	-1.04	3090.4843	11.1	99.3
3	NP ¹⁰⁰⁻²⁸	KMKDLSPRWFYFYLGTGPEAGLPGANK	28	43.16	-0.75	3223.7303	9.7	99.4
4	NP ²¹⁵⁻²²⁹	GDAALALLLDRLNQLLESKMSGK	23	44.86	-0.29	2769.2873	10.6	99.3
5	NP ³¹¹⁻³³⁵	SASAFGMSRIGMEVTPSGTWLTYT	25	46.39	0.26	2698.0933	6.9	98.9
6	NP ³³⁰⁻³⁷⁴	VILLNKHIDAYKTFPPTEPKDKKKK	25	33.36	-1.06	2952.5603	10.3	99.2
7	M ⁹⁴⁻¹¹¹	SYFIASFRLFARTRSMWSEFNPETNIL	26	59.53	0.03	3155.6593	11.1	100.0
8	M ¹⁵⁶⁻¹⁵⁸	SELVIGAVILRGHLRIAGHHLGR	23	41.21	0.51	2474.9643	12.2	100.0
9	E ⁵⁰⁻⁶⁹	SLVKPSFYVYSRVKNLNSSRV	21	36.83	-0.23	2443.8583	10.8	100.0
10	ORF3 ²⁰²⁻¹⁹	VLHSYFTSDYYQLYSTQLRR	20	41.13	-0.67	2540.8443	9.4	99.9
11	ORF7a ⁷⁶⁻⁹⁷	QLRARSVSPKLFIRQEEVQELYR	23	37.01	-0.8	2846.3133	10.4	97.3
12	NSP3 ³²³⁻⁴⁰	RRIVAEWFLAVILFTRFFVYRR	22	73.42	0.15	2987.5793	11.2	98.2
13	NSP6 ⁵⁶²³⁻⁶¹⁷	SAFAMMFLVKHKHAFLLPLSLATVA	25	65.11	1.22	2854.5453	10.6	98.2
14	NSP8 ⁴⁰⁸²⁻¹⁰²	RRNTCDGNTFTYASALWEIQVYVRR	25	45	-0.76	2985.3633	9.9	99.3
15	NSP12 ⁴⁸⁹⁴⁻⁹¹⁵	DALFAYTKRNVIPTTIQMNLKY	22	42.89	-0.16	2601.0893	10	99.9
16	NSP12 ⁵²⁴⁶⁻⁶³	LMIERFYSLAIDAYPLTKRR	20	49.67	0.2	2392.9163	10.4	100.0
17	NSP12 ⁵⁶⁷⁻⁵⁶	QEYADVFFHLYLQYIRKRLTGHMLDMYR	27	53.9	-0.57	3461.0613	9.3	100.0
18	NSP13 ⁵⁴⁵⁵⁻⁷⁹	KLFAAETLKATEETFKLSYGIATVR	25	43.63	0.04	2788.2603	9.6	100.0
19	NSP13 ⁵⁷⁴⁵⁻⁵⁸	RRCPAEIVDTVSALVY	18	35.09	-0.13	1792.1063	9.9	100.0
20	NSP16 ⁶⁸²¹⁻⁴⁵	KYTQLCQYLNTLTLAVPYNMRRVIHF	25	53.14	0.16	3030.6383	9.5	100.0

Table 2. Experimentally proven SARS-CoV-2 epitopes present in the selected peptides. SARS-CoV-2 epitopes and their HLA restriction present in each peptide from epitopes deposited based upon T-cell activation data and/or MHC binding data deposited on the IEDB database

HLA	No	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	
		NP117	NP154	NP158	NP112-19	NP1105	NP1074	NP111	M10-68	E100	ORF3b191	ORF3a8	NSP1210-88	NSP10-67	NSP100-82	NSP12100-67	NSP1210-65	NSP1310-79	NSP1310-58	NSP1610-47	
A*01:01	5			ILSRWQY GTGFGAGLY		VPSGTWLY					FESDYQLY				NTGCGNTFY						
A*02:01	17			YLGTGREAL	LALLLRL RNLQLESKY	GRSRGIV	ILLNKHDA LLNKHD	SAWSNRETHL		SIVRSFY YVSRVNL		KLRQEV	ILTFHYV		FTYSAMW		LMEFNSL SLADAVFL	KLSVGIATV	DDFTVSALV	YNTITLAV	
A*02:03	7									SIVRSFY YVSRVNL		ILTFHYV		ADWHRQVY		LMEFNSL	KLSVGIATV		YNTITLAV		
A*02:06	9									SIVRSFY YVSRVNL		ILTFHYV		ADWHRQVY		LMEFNSL	KLSVGIATV	DDFTVSALV	YNTITLAV		
A*11:01	6					ASAFGASR	KTFPTEPK				SAGAMHYK					ADAVPEK	KELAAETLK		TLAPPNSMR		
A*23:02	9		YTRATRI					SIVRSFY YVSRVNL			YVQISTQL		AVLETRF		TYASALWEI						
A*23:02								SIVRSFY YVSRVNL													
A*28:01	3			ILSRWQY		VPSGTWLY															
A*30:01	13			AGLPGANK	QLKSNK ALALLER	ASAFGASR	AVLTPTEPK KTFPTEPK		RAGHILGR			QLKSNK					ADAVPEK	KLSVGIATV KELAAETLK		TLAPPNSMR	
A*30:01	3					VPSGTWLY															
A*31:01	5					ASAFGASR			AVLQGHK												
A*32:01	1					ASAFGASR	REAFRTSMW														
A*33:01	2					ASAFGASR			AVLQGHK												
A*98:01	6					ASAFGASR	AVLQGHK ILRAGHILGR														
A*98:02	1																				
B*07:02	6	LPNNTASWF	PRGQCPH	SPRWYTYL																	
B*08:01	5																				
B*14:02	2	NTASWITAL																			
B*15:01																					
B*16:01	3							REAFRTSMW													
B*27:05	0																				
B*35:01	2																				
B*40:01	4					MEVPSGTWL			SELVGVIL												
B*44:02/03	4					MEVPSGTWL PSGTWLYH			SELVGVIL												AEVDYNSAL
B*51:01	4	LPNNTASWF																			CPAFVDTY

Continued

Table 2. Continued

HLA	No	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	
		NP107	NP104	NP105	NP103	NP106	NP102	NP101	M100	E100	ORF3 ¹⁰¹⁻¹¹¹	ORF7 _A ¹¹²⁻¹²⁰	NSP13 ¹²¹⁻¹³⁰	NSP6 ¹³¹⁻¹⁴⁰	NSP8 ¹⁴¹⁻¹⁵⁰	NSP12 ¹⁵¹⁻¹⁶⁰	NSP12 ¹⁶¹⁻¹⁷⁰	NSP13 ¹⁷¹⁻¹⁸⁰	NSP13 ¹⁸¹⁻¹⁹⁰	NSP16 ¹⁹¹⁻²⁰⁰	
PS3-01	4	LPNSTASWF			AFSASAFGGM																
PS7-01	3			KMDLSRW	AFSASAFGGM			LHGRSMW													
PS8-01	1							LHRRSMW													
CLASS II	19	TASWFAIQHKEE, NSGPDQIGYTRAT LPNSTASWFAIQH, QGYTRATRVRGDDK	MIDLSRWYFYLTG, NSGPDQIGYTRAT LPNSTASWFAIQH, QGYTRATRVRGDDK	MDLSRWYFYLTG, NSGPDQIGYTRAT LPNSTASWFAIQH, QGYTRATRVRGDDK	GDAALLLLELNQ, SASAFGSRGMV FGSRGMVYTSGT, LLLRLNQLSMM	SASAFGSRGMV, FGSRGMVYTSGT GSRGMVYTSGT, YKTPTEKDKKK	VLLNKHDAKTFP, LLNKHDAKTFP YKTPTEKDKKK	GAVLRLGRLRAGHILGR, LLNKHDAKTFP YKTPTEKDKKK	FYYSRVNLSNSV, VLSHTTSYQYQ												
Total	144	6	8	12	17	17	8	8	9	9	3	5	6	5	5	8	8	9	9	6	7

Table 3. Predicted SARS-CoV-2 epitopes present in the selected peptides. Predicted SARS-CoV-2 epitopes and their HLA restriction present in each peptide based upon the NetMHCpan EL4.1 HLA binding prediction (Reynisson et al., 2020). Positive binding was defined as the top 0.1% scoring epitopes

HLA	No	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19			
		NP1457	NP664	NP9534	NP15235	NP15355	NP15373	M ¹⁴⁵ 115	M ¹⁴⁵ 114	E ¹⁴⁵ 48	ORF3 ²⁵¹⁻¹⁹	ORF7 ²⁶⁻⁶⁷	NSP3 ³²¹⁻⁴⁵	NSP6 ³²⁵⁻⁴⁴⁷	NSP8 ³⁸⁵⁻⁵⁰³	NSP12 ⁵⁸⁴⁻⁶¹⁰	NSP13 ⁶¹⁰⁻⁷⁹	NSP13 ⁷¹⁵⁻⁷⁸	NSP16 ⁶⁰¹⁻⁶				
A*01:01	8	NSPDDQIGYY VLTGTPEAGLPY			LINKHIDAY					HSYFISDY YFISDYQYLY		IVAEWFAY					ATEEFPKLSY	IVDTVSAIYV					
A*02:01	4		ILLDRLNQL											FLPFLATV	SALWEIQQVY ALWEIQQVY								
A*02:03	2													FLPFLATV							TLAVFPNMRV		
A*02:06	2													FLPFLATV	SALWEIQQY								
A*11:01	4	FFALTQHKG		AGLPYGANK		AYKTPPPEPK					SVSPKLER												
A*24:02/A*23:02	7		YYLGTGPEAGL							LVKISFYVY	SYFTSDYQL		EWHAVILF	MFVKHKHAF							VPYNNRVIHE KYTQLCQYL		
A*26:01	2												IVAEWFAY								EIVDTYSAL		
A*03:01	2	FFALTQHKG								LVKISFYVY													
A*30:01	1									SFYYSRVK													
A*31:01	2									SIVKPSFYVY			IVAEWFAY										
A*32:01	3								RIAGHHLGR			RSVSPKLER											
A*33:01	3								LVIGAVILR				FLAVILFR										
A*68:01	9	FFALTQHKG		EAGLPYGANK		YKTPPPEPK			ELVIGAVILR			SVSPKLER	FLAVILFR	DALEAYTKR			LSYGLATVR				ITLAVPNNMR		
A*68:02	4																NFTTYSAL	LMERFVSL	ETFKLSYGI	EIVDTYSAL			
B*07:02	0																						
B*08:01	5			DISPRWYF	ILLDRLNQL																		
B*14:02	0																						
B*15:01	9			DISPRWYFY KMKDISRWY		LINKHIDAY				LVKISFYVY	VLHSYFTSDY	RQEEVQELY	IVAEWFAY	MMFVKHKHAF							TLKATEETF		
B*18:01	1			KDLSRWYF																			
B*27:05	1									SRVKNLNSR													
B*35:01	6	SPDDQIGYY		GPEAGLPY		TPSGTWIY							IVAEWFAY				TCDGNIFY				FVSLADAY		
B*40:01	2			KDLSRWYF																	AEMDTYSAL		
B*44:02/03	4			KDLSRWYF		MEVTPSGTW															ATEEFPKLSY TEEFPKLSY		
B*51:01	5				DAALALILL	SAFFGMSR															SALWEIQQY	FAYTKRNYI	LAVFPNMRV

Continued

Table 3. Continued

HLA	No	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19		
			NP ⁶⁰⁻⁶⁴	NP ⁶⁸⁻⁷⁴	NP ⁷⁵⁻⁷⁹	NP ⁸⁰⁻⁸⁴	NP ⁸⁵⁻⁸⁹	M ⁹⁰⁻⁹⁴	M ⁹⁵⁻⁹⁹	E ¹⁰⁰	ORF3 ¹⁰¹⁻¹¹⁹	ORF7 ¹²⁰⁻¹²⁷	NSP3 ¹²⁸⁻¹⁴⁰	NSP6 ¹⁴¹⁻¹⁴⁷	NSP8 ¹⁴⁸⁻¹⁶²	NSP12 ¹⁶³⁻¹⁶⁹	NSP12 ¹⁷⁰⁻¹⁷⁶	NSP13 ¹⁷⁷⁻¹⁷⁹	NSP13 ¹⁸⁰⁻¹⁸⁴	NSP16 ¹⁸⁵⁻¹⁸⁹		
P*53:01	0																					
P*57:01	4											RARSYSKLF	LAVLETRF		NIFTYASALW							LTLAVPTNM
P*58:01	3		NSSPDDQIGY												NIFTYASALW							LTLAVPTNM
CLASS II	2						SFRLPRTSMAWSE							LCLLEPLSLATVAYF								
Total	95	3	3	11	3	3	4	1	3	7	4	5	9	8	8	4	2	6	4	6	4	6

Table 4. Conservation of the candidate peptides and regions of the Spike protein between different Coronaviruses. Amino acid conservation was assessed for the peptides selected from the membrane, envelop, ORF3a, ORF7a, nucleoprotein and non-structural proteins between SARS-CoV-2 reference strain (YP_009724389.1) and 94 other coronaviruses. Conservation between reference strain (YP_009724389.1) and 94 other coronaviruses for the receptor binding domain (RBD) and N-terminal, domain (NTD) both antigenic sites for neutralising antibody, was also performed

Genus	Strain	Envelope				Membrane				Nucleocapsid				ORF1ab						Spike							
		ORF7a	ORF3a	Envelope	Membrane	M94-111	M136-158	NP41-57	NP41-158	NP100-128	NP215-229	NP311-335	NP350-374	NSP323233-45	NSP6622-647	NSP84082-102	NSP124894-915	NSP125246-63	NSP125247-95	NSP135455-79	NSP135768-76	NTD	RBD	RBD439-474	RBD550-59		
Betacoronavirus	SARS-CoV-2 Wuhan-Hu-1	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
	Bat CoV RaTG13	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	95.7	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	98.5	91.5	81.8	100.0	100.0	
	PCoV_GXC-P2V	4.5	77.8	100.0	100.0	100.0	100.0	100.0	100.0	100.0	95.7	100.0	100.0	83.3	100.0	90.5	100.0	100.0	100.0	100.0	100.0	86.7	87.3	81.8	92.0	100.0	
	PCoV PCoV_GX-P4E	100.0	77.8	100.0	100.0	100.0	100.0	100.0	100.0	95.7	100.0	100.0	83.3	100.0	90.5	100.0	100.0	100.0	100.0	100.0	100.0	86.7	87.6	81.8	92.0	100.0	
	Bat SARS-like CoV RsSHC014	100.0	66.7	85.7	100.0	87.0	93.3	95.7	92.9	87.0	96.0	100.0	72.2	85.2	95.2	100.0	100.0	100.0	100.0	72.0	100.0	47.2	75.7	52.3	78.0	78.0	
	Bat SARS-like CoV Rs4084	100.0	66.7	85.7	100.0	87.0	93.3	95.7	92.9	87.0	92.0	100.0	77.8	85.2	95.2	100.0	100.0	100.0	100.0	100.0	100.0	47.2	75.3	52.3	78.0	78.0	
	SARS-like CoV WIV16	90.9	66.7	85.7	100.0	87.0	93.3	95.7	92.9	87.0	96.0	100.0	77.8	85.2	95.2	100.0	100.0	100.0	64.0	100.0	100.0	47.7	74.9	45.5	78.0	78.0	
	Bat SARS-like CoV WIV1	90.9	66.7	85.7	100.0	87.0	93.3	95.7	92.9	87.0	96.0	100.0	77.8	85.2	95.2	100.0	100.0	100.0	72.0	100.0	100.0	47.2	74.9	45.5	78.0	78.0	
	Bat SARS-like CoV Rs3367	100.0	66.7	85.7	100.0	87.0	93.3	95.7	92.9	87.0	96.0	100.0	77.8	85.2	95.2	100.0	100.0	100.0	72.0	100.0	100.0	47.2	74.9	45.5	78.0	78.0	
	Bat SARS-like CoV Rs9401	95.5	66.7	85.7	96.2	87.0	93.3	95.7	92.9	87.0	96.0	100.0	77.8	85.2	95.2	100.0	100.0	100.0	100.0	100.0	100.0	47.2	74.5	45.5	78.0	78.0	
	CoV BRs-BetaCoV/YN2018B	85.7	100.0	87.0	100.0	87.0	93.3	95.7	92.9	87.0	96.0	100.0	77.8	85.2	95.2	100.0	100.0	100.0	100.0	100.0	100.0	47.2	74.5	45.5	78.0	78.0	
	SARS-related-CoV Chret CoV 007/2004	0.0	66.7	85.7	96.2	87.0	93.3	95.7	92.9	87.0	96.0	100.0	72.2	85.2	95.2	100.0	100.0	100.0	60.0	100.0	100.0	46.2	73.7	45.5	78.0	78.0	
	SARS-related CoV civet010	0.0	66.7	85.7	96.2	87.0	93.3	95.7	92.9	87.0	96.0	100.0	72.2	85.2	95.2	100.0	100.0	100.0	60.0	100.0	100.0	46.2	73.7	45.5	78.0	78.0	
	SARS CoV civet020	0.0	66.7	85.7	96.2	87.0	93.3	95.7	92.9	87.0	96.0	100.0	72.2	85.2	95.2	100.0	100.0	100.0	60.0	100.0	100.0	46.2	73.7	45.5	78.0	78.0	
	SARS CoV-1 B039	0.0	66.7	85.7	96.2	87.0	93.3	95.7	92.9	87.0	96.0	100.0	72.2	85.2	95.2	100.0	100.0	100.0	60.0	100.0	100.0	46.2	73.7	45.5	78.0	78.0	
SARS CoV A022	100.0	66.7	85.7	96.2	87.0	93.3	95.7	92.9	87.0	96.0	100.0	72.2	85.2	95.2	100.0	100.0	100.0	60.0	100.0	100.0	46.2	73.7	45.5	78.0	78.0		
SARS-CoV-1 Tor2	66.7	85.7	96.2	87.0	93.3	95.7	92.9	87.0	96.0	96.0	100.0	72.2	85.2	95.2	100.0	100.0	100.0	100.0	100.0	100.0	46.2	73.4	45.5	78.0	78.0		
SARS-CoV-1 GZ02	0.0	66.7	9.5	96.2	87.0	93.3	95.7	92.9	87.0	96.0	100.0	72.2	85.2	95.2	100.0	100.0	100.0	60.0	100.0	100.0	45.6	73.0	45.5	76.0	76.0		
SARS-related CoV BIKY72	100.0	50.0	85.7	92.3	82.6	86.7	91.3	92.9	91.3	92.0	100.0	83.3	77.8	90.5	100.0	100.0	100.0	80.0	92.9	96.0	41.0	73.0	45.5	78.0	78.0		
Bat-CoV BM48-31/BGR/2008	0.0	55.6	85.7	92.3	87.0	86.7	91.3	92.9	91.3	92.0	100.0	72.2	81.5	95.2	100.0	100.0	100.0	100.0	100.0	100.0	38.5	70.7	31.8	80.0	80.0		
Sarbecovirus Bat CoV BM48-31/BGR/2008	0.0	55.6	85.7	92.3	87.0	91.3	86.7	92.9	91.3	92.0	80.8	72.2	81.5	95.2	100.0	100.0	100.0	72.0	100.0	100.0	38.5	70.7	31.8	80.0	80.0		

Continued

Table 4. Continued

Genus	Strain	ORF7a ORF3a Envelope				Membrane				Nucleocapsid				ORF1ab						Spike						
		ORF7a	ORF3a	E50-69	ORF7a ORF3a	M84-111	M136-158	NP41-57	NP69-94	NP100-128	NP215-229	NP311-335	NP350-374	NSP32323-45	NSP63622-647	NSP84082-102	NSP124894-915	NSP125246-63	NSP125267-95	NSP13 NSP13(5768-76)	NSP166821-452	NTD	RBD	RBD439-474	RBD590-59	
Betacoronavirus	Bat SARS-like CoV-SL-CoVZC45	0.0	94.4	100.0	100.0	100.0	100.0	96.7	95.7	96.4	87.0	96.0	100.0	100.0	100.0	95.2	100.0	100.0	100.0	100.0	100.0	61.0	69.9	31.8	78.0	
	Bat-SL-CoVZXC21	100.0	100.0	100.0	100.0	100.0	100.0	96.7	95.7	96.4	87.0	96.0	100.0	100.0	100.0	95.2	100.0	100.0	100.0	100.0	100.0	59.0	69.5	29.5	78.0	
Betacoronavirus	BRRF-BetaCoV/SG2018		85.7	92.3	87.0	90.0	95.7	92.9	92.9	92.9	87.0	96.0	92.3	77.8	85.2	95.2	100.0	100.0	100.0	100.0	100.0	42.1	66.0	31.8	78.0	
	Bat-CoV Rp/Shamsh2011	66.7	85.7	85.7	87.0	90.0	95.7	92.9	92.9	92.9	87.0	96.0	100.0	77.8	85.2	95.2	95.2					41.0	66.0	34.1	82.0	
Betacoronavirus	CoV BRRS-BetaCoV/YN2018C		85.7	96.2	87.0	93.3	95.7	92.9	92.9	92.9	87.0	96.0	100.0	77.8	85.2	95.2	100.0	100.0	100.0	100.0	100.0	47.2	65.6	31.8	78.0	
	BatSARS-CoV HKU3-2	0.0	66.7	85.7	87.0	90.0	95.7	92.9	92.9	92.9	87.0	96.0	100.0	72.2	85.2	95.2	100.0	100.0	100.0	100.0	100.0	47.2	65.3	34.1	80.0	
Betacoronavirus	BatSARS CoV HKU3-4	45.5	5.6	85.7	87.0	90.0	95.7	92.9	92.9	92.9	87.0	96.0	100.0	72.2	85.2	95.2	100.0	100.0	100.0	100.0	100.0	47.2	65.3	34.1	80.0	
	BatSARS CoV HKU3-3	90.9	5.6	85.7	87.0	90.0	95.7	92.9	92.9	92.9	87.0	96.0	100.0	72.2	85.2	95.2	100.0	100.0	100.0	100.0	100.0	47.2	65.3	34.1	80.0	
Betacoronavirus	BatSARS CoV HKU3-1	100.0	66.7	85.7	87.0	90.0	95.7	92.9	92.9	92.9	87.0	96.0	100.0	72.2	85.2	95.2	100.0	100.0	100.0	100.0	100.0	47.2	65.3	34.1	80.0	
	BRRS-Beta CoV/GX2013	100.0	72.2	85.7	87.0	90.0	95.7	92.9	92.9	92.9	87.0	96.0	100.0	77.8	85.2	95.2	100.0	100.0	100.0	100.0	100.0	46.7	65.3	34.1	80.0	
Betacoronavirus	BRRF-BetaCoV/SX2013	95.5	61.1	81.0	87.0	86.7	95.7	92.9	92.9	92.9	87.0	96.0	100.0	77.8	85.2	95.2	100.0	100.0	100.0	100.0	100.0	47.7	64.9	34.1	78.0	
	BRRF-BetaCoV/HeB2013	95.5	61.1	85.7	87.0	86.7	95.7	92.9	92.9	92.9	87.0	96.0	100.0	77.8	85.2	95.2	100.0	100.0	100.0	100.0	100.0	47.7	64.9	34.1	78.0	
Betacoronavirus	Bat SARS-like CoV/YNIF_31C	100.0	61.1	85.7	87.0	90.0	95.7	92.9	92.9	92.9	87.0	96.0	100.0	72.2	85.2	95.2	100.0	100.0	100.0	100.0	100.0	47.2	64.9	34.1	76.0	
	Bat CoV 279/2005	0.0	72.2	85.7	87.0	90.0	95.7	92.9	92.9	92.9	87.0	96.0	100.0	77.8	85.2	95.2	100.0	100.0	100.0	100.0	100.0	48.2	64.5	31.8	80.0	
Betacoronavirus	Bat SARS-like CoV/R4237	100.0	66.7	85.7	87.0	93.3	95.7	92.9	92.9	92.9	87.0	96.0	100.0	77.8	85.2	95.2	100.0	100.0	100.0	100.0	100.0	48.2	64.5	31.8	80.0	
	Bat CoV 273/2005	95.5	61.1	85.7	87.0	86.7	95.7	92.9	92.9	92.9	87.0	96.0	100.0	77.8	85.2	95.2	100.0	100.0	100.0	100.0	100.0	48.2	64.1	31.8	78.0	
Betacoronavirus	Bat-CoV Cp/Yunnan2011		66.7	85.7	87.0	90.0	95.7	92.9	92.9	92.9	87.0	96.0	100.0	83.3	85.2	95.2	95.2					47.7	64.1	34.1	80.0	
	SARS CoV Rs_672/2006	100.0	66.7	85.7	87.0	90.0	95.7	92.9	92.9	92.9	87.0	96.0	100.0	72.2	85.2	95.2	100.0	100.0	100.0	100.0	100.0	43.6	64.1	29.5	80.0	
Betacoronavirus	BRRS-Beta-CoV/YN2018D		85.7	100.0	87.0	93.3	95.7	92.9	92.9	92.9	87.0	96.0	100.0	77.8	85.2	95.2	100.0	100.0	100.0	100.0	100.0	43.6	64.1	29.5	82.0	
	BRRS-BetaCoV/YN2013	100.0	61.1	85.7	87.0	95.7	93.3	92.9	92.9	92.9	87.0	96.0	80.8	72.2	85.2	95.2	100.0	100.0	100.0	100.0	100.0	46.7	63.3	31.8	82.0	
Betacoronavirus	BRRF-Beta-CoV/JL2012	95.5	55.6	81.0	87.0	86.7	95.7	92.9	92.9	92.9	87.0	96.0	100.0	77.8	85.2	95.2	100.0	100.0	100.0	100.0	100.0	35.9	63.3	34.1	78.0	
	MERS-related CoV H.savii/Italy/206645-40/2011	0.0	23.8	73.1	34.8	60.0	60.9	67.9	67.9	67.9	47.8	40.0	46.2	38.9	51.9	23.8	86.4	94.4	65.5	100.0	85.7	16.4	24.7	13.6	26.0	

Continued

Table 4. Continued

Genus	Strain	Envelope				Membrane				Nucleocapsid				ORF1ab						Spike								
		ORF7a	ORF3a	ORF3a	ORF7a	M94-111	M136-158	NP41-57	NP69-94	NP100-128	NP215-229	NP311-335	NP350-374	NSP32323-45	NSP65622-647	NSP84082-102	NSP124894-915	NSP125246-63	NSP135267-95	NSP135455-79	NSP135768-76	NSP166821-452	NTD	RBD	RBD439-474	RBD590-59		
Merescovirus	MERS-related CoV PKuhui/Italy/2006645-63/2011	0.0	23.8	73.1	34.8	60.0	60.9	67.9	47.8	40.0	46.2	38.9	51.9	23.8	86.4	94.4	65.5	100.0	85.7	80.0	164	24.7	13.6	26.0				
	Pipistrellus bat CoV HKU5	16.7	9.5	69.2	34.8	60.0	65.2	64.3	30.4	32.0	38.5	50.0	51.9	28.6	86.4	94.4	65.5	100.0	85.7	80.0	16.9	23.6	13.6	26.0				
	MERS-CoV NL140422	5.6	28.6	76.9	39.1	60.0	60.9	64.3	34.8	36.0	46.2	27.8	55.6	28.6	86.4	94.4	65.5	100.0	85.7	80.0	16.4	23.6	11.4	20.0				
	MERS-related CoV KOR/KCDC/001_2018-TSV1	5.6	23.8	73.1	39.1	60.0	65.2	67.9	47.8	36.0	50.0	38.9	55.6	28.6	81.8	94.4	65.5	100.0	85.7	88.0	16.4	23.2	13.6	20.0				
	Hu beta CoV 2 c EMC/2012	5.6	23.8	73.1	39.1	60.0	65.2	67.9	47.8	36.0	50.0	38.9	55.6	28.6	86.4	94.4	65.5	72.0	85.7	88.0	16.4	23.2	13.6	20.0				
	MERS-related CoV MA_30	5.6	23.8	73.1	39.1	60.0	65.2	67.9	47.8	36.0	50.0	38.9	55.6	28.6	86.4	94.4	65.5	100.0	85.7	88.0	16.4	23.2	13.6	20.0				
	BetaCoV England 1	5.6	23.8	73.1	39.1	60.0	65.2	67.9	47.8	36.0	50.0	38.9	55.6	28.6	86.4	94.4	65.5	100.0	85.7	88.0	16.4	23.2	13.6	20.0				
	BetaCoV/133/2005	0.0	28.6	65.4	43.5	56.7	65.2	60.7	39.1	36.0	46.2	50.0	51.9	19.0	90.9	94.4	65.5	72.0	85.7	76.0	12.8	23.2	13.6	22.0				
	MERS-related CoV BRYs-BetaCoV/SC2013	5.6	28.6	73.1	39.1	60.0	65.2	64.3	39.1	40.0	42.3	38.9	51.9	28.6	86.4	94.4	65.5	72.0	85.7	88.0	14.9	22.8	11.4	20.0				
	Tylymsteris bat CoV HKU4 BHP-BetaCoV/GX2012	0.0	28.6	65.4	43.5	56.7	65.2	60.7	43.5	36.0	23.1	44.4	51.9	19.0	86.4	83.3	65.5	64.0	85.7	88.0	15.9	22.8	13.6	24.0				
Hypugo bat CoV HKU25	28.6	28.6	65.4	43.5	56.7	65.2	64.3	39.1	36.0	46.2	38.9	51.9	33.3	86.4	94.4	65.5	100.0	85.7	80.0	14.9	22.8	11.4	20.0					
Bat-CoV HKU4-1	0.0	28.6	65.4	43.5	56.7	65.2	60.7	39.1	36.0	46.2	50.0	48.1	23.8	86.4	83.3	65.5	60.0	85.7	88.0	15.9	22.4	11.4	24.0					
Tylymsteris bat CoV HKU4	28.6	28.6	65.4	43.5	56.7	65.2	60.7	39.1	36.0	46.2	50.0	48.1	23.8	86.4	83.3	65.5	100.0	85.7	88.0	15.9	22.4	11.4	24.0					
CoV Neoromicial/PML-PHE1/RSNA/2011	5.6	23.8	73.1	34.8	56.7	60.9	67.9	47.8	40.0	46.2	44.4	51.9	28.6	86.4	94.4	65.5	72.0	85.7	84.0	14.9	21.2	20.5	26.0					
BetaCoV Erinnacus/VMC/DEU/2012	5.6	28.6	73.1	34.8	56.7	56.5	64.3	39.1	36.0	38.5	27.8	55.6	28.6	86.4	94.4	65.5	100.0	85.7	76.0	12.3	20.5	18.2	24.0					
Myotis ricketti alpha-CoV Sax-2011	11.1	14.3	61.5	21.7	23.3	17.4	32.1	21.7	12.0	11.5	5.6	44.4	28.6	68.2	72.2	31.0	72.0	85.7	72.0	10.8	9.7	6.8	22.0					
Beta CoV HKU24	33.3	33.3	69.2	39.1	36.7	26.1	57.1	26.1	28.0	19.2	44.4	48.1	28.6	77.3	83.3	48.3	100.0	92.9	84.0	12.8	24.7	11.4	24.0					
Rabbit CoV HKU14	28.6	28.6	69.2	39.1	36.7	26.1	57.1	21.7	32.0	19.2	38.9	40.7	33.3	77.3	83.3	48.3	100.0	92.9	84.0	10.3	22.8	6.8	24.0					
Hu-CoV HKU1	38.1	38.1	69.2	39.1	30.4	40.0	60.7	21.7	24.0	19.2	38.9	40.7	38.1	72.7	83.3	48.3	72.0	85.7	84.0	11.8	21.2	6.8	26.0					
Hu-CoV OC43	85.7	85.7	69.2	39.1	36.7	26.1	57.1	21.7	32.0	23.1	38.9	40.7	33.3	77.3	83.3	48.3	60.0	92.9	84.0	10.3	20.8	6.8	24.0					

Continued

Table 4. Continued

Genus	Strain	Envelope					Membrane					Nucleocapsid					ORF1ab					Spike				
		ORF7a	ORF3a	E50-69	M94-111	M136-111	NP41-158	NP49-94	NP100-128	NP215-229	NP311-355	NP350-374	NP32323-43	NSP64622-647	NSP84082-102	NSP124894-915	NSP125246-63	NSP125267-95	NSP1315768-76	NSP166821-452	NTD	RBD	RBD439-474	RD50-59		
Nobecovirus	Rousettus bat CoV GCCDC1	100.0	16.7	28.6	73.1	39.1	36.7	47.8	50.0	34.8	36.0	50.0	33.3	48.1	38.1	86.4	88.9	69.0	100.0	92.9	80.0	154	24.3	11.4	26.0	
	Rousettus bat CoV HKU9	45.5	22.2	23.8	69.2	39.1	40.0	43.5	46.4	30.4	28.0	65.4	33.3	37.0	38.1	90.9	88.9	69.0	100.0	92.9	80.0	159	22.4	11.4	24.0	
	Bat CoV HKU9-5-1	16.7	23.8	23.8	69.2	39.1	50.0	47.8	46.4	26.1	24.0	65.4	38.9	40.7	38.1	90.9	88.9	69.0	72.0	92.9	84.0	164	22.0	11.4	28.0	
Unclassified	CoV FRS-BetaCoV/YN2018A	85.7	85.7	85.7	96.2	87.0	90.0	95.7	92.9	87.0	96.0	100.0	72.2	85.2	95.2	100.0	100.0	100.0	100.0	100.0	48.2	64.9	31.8	82.0		
	Bat CoV Jnyuan-84	85.7	85.7	85.7	96.2	87.0	86.7	95.7	92.9	87.0	96.0	100.0	77.8	85.2	95.2	95.5	100.0	100.0	100.0	47.7	64.9	34.1	78.0			
	Bat CoV JTMCI15	40.9	55.6	81.0	96.2	87.0	86.7	95.7	92.9	87.0	96.0	100.0	77.8	85.2	95.2	95.5	100.0	100.0	80.0	36.4	63.3	34.1	78.0			
Decacovirus	Apodemus peninsulae CoV	33.3	69.2	39.1	40.0	26.1	57.1	26.1	28.0	19.2	44.4	48.1	28.6	48.1	28.6	77.3	83.3	48.3	100.0	92.9	84.0	12.8	23.9	11.4	24.0	
	Bat V s-CoV-1	5.6	28.6	73.1	39.1	60.0	65.2	64.3	39.1	40.0	42.3	38.9	51.9	28.6	28.6	90.9	94.4	65.5	72.0	85.7	76.0	164	22.4	9.1	26.0	
	Bat-CoV CMR704-P12	16.7	19.0	19.0	61.5	30.4	40.0	43.5	53.6	30.4	32.0	65.4	38.9	44.4	28.6	90.9	88.9	69.0	96.0	85.7	80.0	13.3	22.4	11.4	18.0	
Dinacovirus	MERS-related CoV BIV s-BetaCoV/SC2013	5.6	23.8	73.1	34.8	56.7	60.9	67.9	47.8	40.0	46.2	44.4	51.9	28.6	86.4	94.4	65.5	100.0	85.7	84.0	149	21.2	20.5	26.0		
	Bat coronavirus PDF-2180	5.6	23.8	73.1	34.8	60.0	65.2	67.9	43.5	40.0	42.3	44.4	51.9	28.6	86.4	94.4	65.5	100.0	85.7	84.0	149	19.7	18.2	28.0		
	BtRE-AlphaCoV/Hub2013	13.6	11.1	14.3	53.8	21.7	30.0	8.7	46.4	8.7	16.0	7.7	16.7	40.7	19.0	77.3	77.8	31.0	60.0	78.6	68.0	11.3	10.0	4.5	18.0	
Setracovirus	bat-CoV HKU10	100.0	16.7	19.0	53.8	21.7	26.7	8.7	42.9	13.0	8.0	7.7	22.2	48.1	19.0	77.3	72.2	31.0	72.0	78.6	76.0	9.2	9.3	4.5	14.0	
	Rousettus bat CoV HKU10	4.5	16.7	19.0	53.8	21.7	26.7	8.7	42.9	13.0	8.0	7.7	22.2	48.1	19.0	77.3	72.2	31.0	100.0	78.6	76.0	13.3	8.5	2.3	16.0	
	Hu-CoV 229E	14.3	53.8	30.4	30.0	8.7	46.4	13.0	28.0	3.8	5.6	48.1	23.8	48.1	23.8	68.2	72.2	27.6	60.0	85.7	72.0	4.1	9.7	6.8	26.0	
Nyctacovirus	Human CoV 229E	14.3	53.8	30.4	30.0	8.7	46.4	13.0	32.0	3.8	5.6	51.9	23.8	48.1	23.8	72.7	72.2	27.6	72.0	85.7	76.0	4.1	8.5	4.5	26.0	
	Ni63-related bat CoV BKYN163-9b	5.6	19.0	19.0	65.4	26.1	30.0	8.7	39.3	13.0	28.0	19.2	5.6	48.1	23.8	81.8	72.2	31.0	96.0	85.7	72.0	6.7	9.7	4.5	18.0	
	Hu-CoV NL63	5.6	9.5	53.8	21.7	33.3	8.7	42.9	17.4	24.0	7.7	5.6	48.1	23.8	81.8	72.2	27.6	60.0	85.7	72.0	9.2	9.3	13.6	16.0		
Alphacoronavirus	Bat-CoV/Pkuh11	16.7	14.3	14.3	50.0	21.7	26.7	4.3	57.1	13.0	20.0	11.5	22.2	44.4	19.0	68.2	72.2	31.0	96.0	85.7	72.0	7.2	11.2	4.5	22.0	
	BtNs-AlphaCoV/SC2013	16.7	14.3	14.3	50.0	21.7	30.0	4.3	50.0	13.0	28.0	16.7	37.0	44.4	19.0	72.7	72.2	31.0	100.0	85.7	72.0	6.2	8.9	2.3	16.0	
	Pipistrellus bat CoV HKU15 BPp-BetaCoV/GD2013	5.6	9.5	69.2	34.8	60.0	65.2	64.3	30.4	32.0	38.5	50.0	51.9	28.6	28.6	86.4	94.4	65.5	72.0	85.7	80.0	174	22.8	13.6	26.0	
BtMf-AlphaCoV/SAX2011	11.1	14.3	61.5	21.7	23.3	17.4	32.1	21.7	12.0	11.5	5.6	44.4	28.6	28.6	68.2	72.2	31.0	100.0	85.7	72.0	10.8	9.7	6.8	22.0		

Continued

Table 4. Continued

Genus	Strain	Envelope					Membrane					Nucleocapsid					ORF1ab					Spike				
		ORF7a	ORF7a	ORF3a	E50-	E50-	M94-	M136-	NP41-	NP69-	NP100-	NP215-	NP311-	NP350-	NSP3233-	NSP63622-	NSP84082-	NSP124894-	NSP125246-	NSP125267-	NSP13	NSP13(5768-	NSP166821-	NTD	RBD	RBD439-
		69	111	158	57	94	128	229	335	374	45	647	102	915	63	95	79	5455-	76)	452	8.7	6.6	2.3	474	474	59
	BatCoV CDP/HE15/ USA/2006	33.3	14.3	46.2	17.4	30.0	13.0	39.3	21.7	16.0	11.5	16.7	40.7	14.3	68.2	72.2	31.0	100.0	85.7	64.0	8.7	6.6	2.3	20.0	20.0	20.0
Rhinacovirus	bat-CoV HKU2	9.1	14.3	53.8	26.1	36.7	13.0	46.4	8.7	20.0	7.7	11.1	48.1	28.6	72.7	72.2	31.0	60.0	78.6	72.0	8.2	3.1	9.1	20.0	20.0	20.0
	BRRF-AlphaCoV/ YN2012	11.1	14.3	53.8	26.1	30.0	13.0	46.4	8.7	20.0	7.7	11.1	48.1	28.6	72.7	72.2	31.0	100.0	78.6	72.0	7.2	2.7	4.5	20.0	20.0	20.0
Pedacovirus	bat-CoV 512	22.2	23.8	53.8	17.4	30.0	13.0	42.9	26.1	24.0	19.2	16.7	55.6	14.3	77.3	72.2	31.0	60.0	78.6	68.0	9.2	8.1	2.3	22.0	22.0	22.0
	Miniopterus bat-CoV LAFCD62	100.0	11.1	61.5	21.7	33.3	13.0	50.0	17.4	20.0	19.2	16.7	37.0	19.0	72.7	72.2	27.6	72.0	85.7	64.0	8.7	11.2	9.1	22.0	22.0	22.0
Minusacovirus	BtMF-AlphaCoV/ AH2011	11.1	14.3	61.5	21.7	33.3	8.7	50.0	17.4	20.0	19.2	16.7	37.0	19.0	72.7	72.2	27.6	72.0	85.7	64.0	7.7	11.2	9.1	22.0	22.0	22.0
	Bat CoV IB	11.1	9.5	61.5	21.7	33.3	13.0	50.0	17.4	20.0	15.4	16.7	37.0	19.0	72.7	72.2	27.6	72.0	85.7	64.0	8.7	10.8	6.8	22.0	22.0	22.0




Fig. 1. IFN- γ responses to the SARS-CoV-2 synthetic long peptides. PBMC from SARS-CoV-2 seronegative and seropositive donors were used to generate monocyte derived dendritic cells (mDDC) (a) and incubated with SARS-CoV-2 consensus peptides, a group of 11 SLP or CEF peptides prior to co culture with autologous T-cells in triplicate wells of ELISPOT plates to measure T-cell responses (b). MDDC from seven SARS-CoV-2 seropositive individuals were loaded with groups of SLP, SARS-CoV-2 consensus peptides or CEF and CEFT positive controls peptide pools and incubated with autologous T-cells for 48 h in triplicate wells of IFN- γ ELISPOT plates (c) Example of ELISPOT responses to SARS and positive control peptide pools (d). MDDC from the seropositive donors were also loaded with Individual SLP and co cultured with autologous T-cells in order to measure T-cell responses to each peptide (e). Positive responses were defined as a fold increase of 1.5 and statistical significance using a Student's *t*-test.

b) and CD8⁺ (Fig. 3a, b) T-cell responses indicative of HLA restricted T-cell activation were observed with at least one response observed for each peptide with the exception of NSP⁴⁸⁹⁵⁻⁹¹⁵ which demonstrated a CD4⁺ T cell response from the PBMC of one individual but no CD8⁺ T cell responses. This pattern of responses was consistent with other reports studying anti-SARS T-cell responses [13, 17]. Interestingly the greatest responses were observed amongst CD45RA⁺CCR7⁻ TEMRA CD4⁺ T cells (Fig. 4a, b) effector and central memory CD8⁺ T cells (Fig. 4c, d) consistent with recent studies investigating the phenotype of SARS-CoV-2 reactive T-cells [18, 19].

Taken together these data demonstrate the *in vitro* T-cell antigenicity of SARS-CoV-2 derived SLP containing epitopes restricted to multiple HLA and conserved between SARS-CoV-2 variants and other coronaviruses.

DISCUSSION

SARS-CoV-2 vaccines based upon the Spike protein have demonstrated between 60–95% efficacy in phase three trials and are now in widespread use. Although these vaccines are highly efficacious numerous issues remain unresolved. These include supply, cost and the requirement of some vaccines for a cold chain. From an immunological perspective there remains concern that




Fig. 3. CD8+T cell responses to SARS-CoV-2 derived peptides. Peptide loaded MDDC and T-cell co-cultures from six seropositive donors were assessed by flow cytometry for the expression of activation induced markers CD137, CD69 and OX40 (a). Individual SARS-CoV-2 derived peptides increase the expression of activation induced markers from CD8+T cells (b). Groups of SARS-CoV-2 derived or a class II restricted peptide pool increase the expression of activation induced markers from CD8+T cells. Positive responses were defined as a >1.5-fold increase over the no peptide controls.




Fig. 4. SARS-CoV-2 derived peptides preferentially activate CD4⁺ and CD8⁺ T cell subsets. CD4⁺ T cells subsets were defined based upon their expression of CCR7 and CD45RA and assessed for their expression of activation induced markers from the PBMC of individuals with previous SARS-CoV-2 infection (a). Percentage expression of CD137 and CD40L is shown for CD4⁺ T cell subsets from three donors. CD8⁺ T cells subsets were defined based upon their expression of CCR7 and CD45RA and assessed for their expression of activation induced markers from the PBMC of individuals with previous SARS-CoV-2 infection (b). Percentage expression of CD137 and OX40 is shown for CD8⁺ T cell subsets from three donors.

also targets other SARS-CoV-2 proteins including the nucleoprotein, membrane protein and non-structural proteins of ORF1ab [17, 19, 31–33]. These studies show that significant variation exists in the T-cell antigenic targets of SARS-CoV-2 which may lower the efficacy of spike-based vaccines in patients who demonstrate limited anti-spike T-cell activation. Although infection offers an opportunity to gain immune protection to diverse SARS-CoV-2 antigen, and some studies have identified strong T-cell responses from individuals with asymptomatic or mild COVID-19, other studies suggest that asymptomatic infection may not provide sufficient antigenic stimulation to activate protective, long-lasting anti-SARS-T-cell response [29, 33], supported by observations that CD8+ T cell responses could not be detected in 30% of convalescent individuals [12].

In this study, immunogenic regions from SARS-CoV-2 proteins other than the spike were identified and their conservation amongst selected alpha and beta coronaviruses was assessed. The selected peptides contain multiple epitopes restricted to the most common HLA class I molecules and which have previously demonstrated induction of T-cell activation in response to SARS-CoV-2 (Table 2). Importantly, these peptides were highly conserved between different coronaviruses, particularly of the SARS-like Serbacoviruses, compared to the receptor binding domain of the spike protein, the major antigenic site of neutralising antibodies (Table 4). Each of the peptides induced T-cell responses from the T-cells of at least one individual with previous SARS-CoV-2 infection (Figs 1–3) however future work is warranted in order to extend the limited observations made here. This could define the nature of T-cell responses raised to the peptides in greater detail along with their ability to contribute to protection from SARS-CoV-2 challenge and induce responses from naïve donors, which were not observed in the present study, likely due to the small number of experiments performed using cells from naïve donors. Nevertheless, these data indicate that T-cell responses raised against these peptides may cross react with future SARS-CoV-2 variants, which may evolve to escape neutralising antibody responses, and against future emerging coronaviruses. This is supported by studies screening SARS-CoV-2 epitopes in COVID-19 and uninfected patients which have observed SARS-CoV-2 epitopes specific CD4+ and CD8+ T cell responses in SARS-CoV-2 uninfected individuals, which share homology with epitopes in other human coronaviruses [6, 8–10, 31]. The SARS-CoV-2 peptides studied here include 125 epitopes identified by these epitope screening studies of SARS-CoV-2 patients [6, 8, 9, 19, 31, 32]. A recent study, screening epitopes in 16 COVID-19 patients identified 122 epitopes reactive to T-cells in these individuals [33]. The peptides detailed in the present study share 17 epitopes with this study. Future epitope screening studies may reveal further SARS-CoV-2 specific epitopes which, if part of epitope rich clusters, may identify new regions suitable for the generation of synthetic long peptides of the kind studied here.

The peptides are derived from the immunodominant viral proteins other than the spike (Table 1) so could complement existing spike-based vaccination and contribute to the induction of broad T-cell reactivity associated with improved anti-viral immunity. For example, antibody titres in COVID-19 patients correlate with CD4+ T cell immune responses not just to the spike protein, but also to the nucleoprotein and membrane protein [14, 34] and the peptides studied here include multiple class II restricted epitopes from the nucleoprotein and membrane protein not present in existing spike-based vaccines. Harnessing CD4+ T cell epitopes from other SARS-CoV-2 antigen represents a strategy for improving the response or longevity of protection afforded by existing spike-based vaccines, particularly given observations that the diversity [6], functionality [18, 35] and quality [36] of CD4+ T cell activation supports the generation of cellular and humoral immune responses associated with protection. These observations are supported by the reports that hospitalised patients with robust B-cell responses yet suffering from severe COVID-19 infection demonstrate limited activation of circulating CD4+ follicular T-cells, indicative of the importance of these cells to effective humoral immunity [37]. Vaccine approaches including the envelop and nucleoprotein are under investigation [38, 39], consistent with this approach.

A recent study showed that CD8+ T cell responses against SARS-CoV-2 were raised against approximately 17 epitopes derived from between 1–6 viral proteins (average 2.7) [17]. In our *in vitro* experiments DC-T-cell co-cultures generated from the PBMC of individuals with previous SARS-CoV-2 infection responded with IFN- γ expression to an average of five peptides derived from an average of four proteins. Previous studies have demonstrated that broad T-cell responses against multiple epitopes are more effective than narrow responses targeting fewer epitopes [40–42] and less susceptible to exhaustion [43] indicating that broadening the anti-SARS-CoV-2 T-cell response from vaccination is desirable.

Analysis of T-cell responses to the ChAdOx1 spike-based vaccine showed that nearly 30% of unique TCRs raised by the vaccine mapped to a single region of the spike protein which is mutated in the Beta variant of SARS-CoV-2 [44]. This may contribute to the failure of ChAdOx1 to protect against mild-to-moderate COVID-19 [20]. These studies suggest that variation in SARS-CoV-2 has the potential to reduce vaccine efficacy and support the use of SARS-CoV-2 antigen derived from non-spike proteins.

Synthetic long peptides of the kind studied here have been used in numerous therapeutic vaccines for both infectious disease and cancer and demonstrated an ability to induce efficacious T-cell responses [45–47]. Peptide based vaccines are inherently safe, can be easily manufactured, combined with different adjuvants, including those selected for therapeutic properties such as trained innate immunity, and do not have the same requirements for cold chains as other vaccines. They may be useful alternatives to other vaccine designs for the generation of broad T-cell responses since they exclude non-immunogenic regions and avoid the generation of non-neutralising antibody responses which may be induced by whole virus vaccines [48] and are associated with ADE or toxicity. Alternatively, the peptide regions identified here could also be incorporated into mRNA-based vaccines.

Currently vaccination with whole, killed SARS-CoV-2 virions, which have the potential to induce T-cell responses against each viral protein, have demonstrated lower efficacy compared to mRNA or viral vector-based vaccines indicating that other methods of broadening the antigenic repertoire of SARS-CoV-2 vaccines are needed. The peptides studied here are candidate SARS-CoV-2 immunogens with the potential to increase the breadth and cross reactivity of T-cell activation to existing SARS-CoV-2 vaccines.

Funding information

This work received no specific grant from any funding agency.

Acknowledgements

The authors would like to acknowledge the Institute of Cancer Vaccines and Immunotherapy for supporting this work.

Author contributions

K.P., Investigation, Writing – Review and Editing; A.H., Investigation, Methodology; A.G.D., Conceptualisation Supervision, funding; P.L.S., Investigation, Methodology, Formal Analysis, Conceptualisation, Writing – Original Draft Preparation, Visualisation, Funding.

Conflicts of interest

The authors declare that there are no conflicts of interest.

References

1. The Lancet Respiratory Medicine. Realising the potential of SARS-CoV-2 vaccines—a long shot. *Lancet Respir Med* 2021; 9:117.
2. Arashkia A, Jalilvand S, Mohajel N, *et al.* SARS CoV-2 Rapid Response Team of Pasteur Institute of Iran (PII). Severe acute respiratory syndrome-coronavirus-2 spike (S) protein based vaccine candidates: State of the art and future prospects. *Rev Med Virol* 2021;31:e2183.
3. Williams TC, Burgers WA. SARS-CoV-2 evolution and vaccines: cause for concern? *Lancet Respir Med* 2021;9:333–335.
4. Wu L-P, Wang N-C, Chang Y-H, Tian X-Y, Na D-Y, *et al.* Duration of antibody responses after severe acute respiratory syndrome. *Emerg Infect Dis* 2007;13:1562–1564.
5. Ng O-W, Chia A, Tan AT, Jadi RS, Leong HN, *et al.* Memory T cell responses targeting the SARS coronavirus persist up to 11 years post-infection. *Vaccine* 2016;34:2008–2014.
6. Nelde A, Bilich T, Heitmann JS, Maringer Y, Salih HR, *et al.* SARS-CoV-2-derived peptides define heterologous and COVID-19-induced T cell recognition. *Nat Immunol* 2021;22:74–85.
7. McMahan K, Yu J, Mercado NB, Loos C, Tostanoski LH, *et al.* Correlates of protection against SARS-CoV-2 in rhesus macaques. *Nature* 2021;590:630–634.
8. Lineburg KE, Grant EJ, Swaminathan S, Chatzileontiadou DSM, Szeto C, *et al.* CD8+ T cells specific for an immunodominant SARS-CoV-2 nucleocapsid epitope cross-react with selective seasonal coronaviruses. *Immunity* 2021;54:1055–1065.
9. Schmidt KG, Nganou-Makamdop K, Tenbusch M, El Kenz B, Maier C, *et al.* SARS-CoV-2-seronegative subjects target CTL Epitopes in the SARS-CoV-2 nucleoprotein cross-reactive to common cold coronaviruses. *Front Immunol* 2021;12:627568
10. Koutsakos M, Illing PT, Nguyen THO, Mifsud NA, Crawford JC, *et al.* Human CD8+ T cell cross-reactivity across influenza A, B and C viruses. *Nat Immunol* 2019;20:613–625.
11. Grifoni A, Weiskopf D, Ramirez SI, Mateus J, Dan JM, *et al.* Targets of T cell responses to SARS-CoV-2 coronavirus in humans with COVID-19 disease and unexposed individuals. *Cell* 2020;181:1489–1501.
12. Reynisson B, Alvarez B, Paul S, Peters B, Nielsen M, *et al.* NetMHCpan-4.1 and NetMHCIpan-4.0: improved predictions of MHC antigen presentation by concurrent motif deconvolution and integration of MS MHC eluted ligand data. *Nucleic Acids Res* 2020;48:W449–W454.
13. Braun J, Loyal L, Frentsch M, Wendisch D, Georg P, *et al.* SARS-CoV-2-reactive T cells in healthy donors and patients with COVID-19. *Nature* 2020;587:270–274.
14. Mateus J, Grifoni A, Tarke A, Sidney J, Ramirez SI, *et al.* Selective and cross-reactive SARS-CoV-2 T cell epitopes in unexposed humans. *Science* 2020;370:89–94.
15. Bonifacius A, Tischer-Zimmermann S, Dragon AC, Gussarow D, Vogel A, *et al.* COVID-19 immune signatures reveal stable antiviral T cell function despite declining humoral responses. *Immunity* 2021;54:340–354.
16. Rydyznski Moderbacher C, Ramirez SI, Dan JM, Grifoni A, Hastie KM, *et al.* Antigen-specific adaptive immunity to SARS-CoV-2 in acute COVID-19 and associations with age and disease severity. *Cell* 2020;183:996–1012.
17. Tarke A, Sidney J, Kidd CK, Dan JM, Ramirez SI, *et al.* Comprehensive analysis of T cell immunodominance and immunoprevalence of SARS-CoV-2 epitopes in COVID-19 cases. *Cell Rep Med* 2021;2:100204.
18. Sekine T, Perez-Potti A, Rivera-Ballesteros O, Strålin K, Gorin J-B, *et al.* Robust T cell immunity in convalescent individuals with asymptomatic or mild COVID-19. *Cell* 2020;183:158–168.
19. Peng Y, Mentzer AJ, Liu G, Yao X, Yin Z, *et al.* Broad and strong memory CD4+ and CD8+ T cells induced by SARS-CoV-2 in UK convalescent individuals following COVID-19. *Nat Immunol* 2020;21:1336–1345.
20. Madhi SA, Baillie V, Cutland CL, Voysey M, Koen AL, *et al.* Efficacy of the ChAdOx1 nCoV-19 Covid-19 Vaccine against the B.1.351 Variant. *N Engl J Med* 2021;384:1885–1898.
21. Wall EC, Wu M, Harvey R, Kelly G, Warchal S, *et al.* Neutralising antibody activity against SARS-CoV-2 VOCs B.1.617.2 and B.1.351 by BNT162b2 vaccination. *Lancet* 2021;397:2331–2333.
22. Shrotri M, Navaratnam AMD, Nguyen V, Byrne T, Geismar C, *et al.* Spike-antibody waning after second dose of BNT162b2 or ChAdOx1. *Lancet* 2021;398:385–387.
23. Lee WS, Wheatley AK, Kent SJ, DeKosky BJ. Antibody-dependent enhancement and SARS-CoV-2 vaccines and therapies. *Nat Microbiol* 2020;5:1185–1191.
24. Wang Q, Zhang L, Kuwahara K, Li L, Liu Z, *et al.* Immunodominant SARS coronavirus epitopes in humans elicited both enhancing and neutralizing effects on infection in non-human primates. *ACS Infect Dis* 2016;2:361–376.
25. Soresina A, Moratto D, Chiarini M, Paolillo C, Baresi G, *et al.* Two X-linked agammaglobulinemia patients develop pneumonia as COVID-19 manifestation but recover. *Pediatr Allergy Immunol* 2020;31:565–569.
26. Tan AT, Linster M, Tan CW, Le Bert N, Chia WN, *et al.* Early induction of functional SARS-CoV-2-specific T cells associates with rapid viral clearance and mild disease in COVID-19 patients. *Cell Rep* 2021;34:108728.
27. Xu B, Fan C-Y, Wang A-L, Zou Y-L, Yu Y-H, *et al.* Suppressed T cell-mediated immunity in patients with COVID-19: A clinical retrospective study in Wuhan, China. *J Infect* 2020;81:e51–e60.
28. Mazzoni A, Salvati L, Maggi L, Capone M, Vanni A, *et al.* Impaired immune cell cytotoxicity in severe COVID-19 is IL-6 dependent. *J Clin Invest* 2020;130:4694–4703.

29. Bacher P, Rosati E, Esser D, Martini GR, Saggau C, et al. Low-avidity CD4+ T cell responses to SARS-CoV-2 in unexposed individuals and humans with severe COVID-19. *Immunity* 2020;53:1258–1271.
30. Diao B, Wang C, Tan Y, Chen X, Liu Y, et al. Reduction and functional exhaustion of T cells in patients with coronavirus disease 2019 (COVID-19). *Front Immunol* 2020;11:827.
31. Le Bert N, Tan AT, Kunasegaran K, Tham CYL, Hafezi M, et al. SARS-CoV-2-specific T cell immunity in cases of COVID-19 and SARS, and uninfected controls. *Nature* 2020;584:457–462.
32. Ferretti AP, Kula T, Wang Y, Nguyen DMV, Weinheimer A, et al. Unbiased screens show CD8+ T cells of COVID-19 patients recognize shared epitopes in SARS-CoV-2 that largely reside outside the spike protein. *Immunity* 2020;53:1095–1107.
33. Saini SK, Hersby DS, Tamhane T, Povlsen HR, Amaya Hernandez SP, et al. SARS-CoV-2 genome-wide T cell epitope mapping reveals immunodominance and substantial CD8+ T cell activation in COVID-19 patients. *Sci Immunol* 2021;6:58.
34. Kusnadi A, Ramírez-Suástegui C, Fajardo V, Chee SJ, Meckiff BJ, et al. Severely ill COVID-19 patients display impaired exhaustion features in SARS-CoV-2-reactive CD8+ T cells. *Sci Immunol* 2021;6:55.
35. Bertoletti A, Tan AT, Le Bert N. The T-cell response to SARS-CoV-2: kinetic and quantitative aspects and the case for their protective role. *Oxford Open Immunology* 2021;2:iqab006.
36. Havenar-Daughton C, Carnathan DG, Torrents de la Peña A, Pauthner M, Briney B, et al. Direct probing of germinal center responses reveals immunological features and bottlenecks for neutralizing antibody responses to HIV Env trimer. *Cell Rep* 2016;17:2195–2209.
37. Mathew D, Giles JR, Baxter AE, Oldridge DA, Greenplate AR, et al. Deep immune profiling of COVID-19 patients reveals distinct immunotypes with therapeutic implications. *Science* 2020;369:eabc8511.
38. Abdelmageed MI, Abdelmoneim AH, Mustafa MI, Elfadol NM, Murshed NS, et al. Design of a multiepitope-based peptide vaccine. Multiepitope-Based Peptide Vaccine against the E protein of huProtein of Human COVID-19: An immunoinformatics approach. *Biomed Res Int* 2020;26:83286.
39. Tazehkand MN, Hajipour O. Evaluating the vaccine potential of a tetravalent fusion protein against coronavirus (COVID-19). *J Vaccines Vaccin* 2020;11:411.
40. Meyer-Olson D, Shoukry NH, Brady KW, Kim H, Olson DP, et al. Limited T cell receptor diversity of HCV-specific T cell responses is associated with CTL escape. *J Exp Med* 2004;200:307–319.
41. Nikolich-Zugich J, Slifka MK, Messaoudi I. The many important facets of T-cell repertoire diversity. *Nat Rev Immunol* 2004;4:123–132.
42. Chang C-M, Feng P-H, Wu T-H, Alachkar H, Lee K-Y, et al. Profiling of T cell repertoire in SARS-CoV-2-infected COVID-19 patients between mild disease and pneumonia. *J Clin Immunol* 2021;41:1131–1145.
43. Fuller MJ, Zajac AJ. Ablation of CD8 and CD4 T cell responses by high viral loads. *J Immunol* 2003;170:477–486.
44. Swanson PA, Padilla M, Hoyland W, McGlinchey K, Fields PA, et al. T-cell mediated immunity after AZD1222 vaccination: A polyfunctional spike-specific Th1 response with a diverse TCR repertoire. *medRxiv* 2021.
45. Lind A, Sommerfelt M, Holmberg JO, Baksaas I, Sørensen B, et al. Intradermal vaccination of HIV-infected patients with short HIV Gag p24-like peptides induces CD4 + and CD8 + T cell responses lasting more than seven years. *Scand J Infect Dis* 2012;44:566–572.
46. Malonis RJ, Lai JR, Vergnolle O. Peptide-based vaccines: current progress and future challenges. *Chem Rev* 2020;120:3210–3229.
47. Pollard RB, Rockstroh JK, Pantaleo G, Asmuth DM, Peters B, et al. Safety and efficacy of the peptide-based therapeutic vaccine for HIV-1, Vacc-4x: a phase 2 randomised, double-blind, placebo-controlled trial. *Lancet Infect Dis* 2014;14:291–300.
48. Rostad CA, Anderson EJ. Optimism and caution for an inactivated COVID-19 vaccine. *Lancet Infect Dis* 2021;21:581–582.

Five reasons to publish your next article with a Microbiology Society journal

1. The Microbiology Society is a not-for-profit organization.
2. We offer fast and rigorous peer review – average time to first decision is 4–6 weeks.
3. Our journals have a global readership with subscriptions held in research institutions around the world.
4. 80% of our authors rate our submission process as 'excellent' or 'very good'.
5. Your article will be published on an interactive journal platform with advanced metrics.

Find out more and submit your article at microbiologyresearch.org.