

Zoonotic *Ancylostoma ceylanicum* Hookworm Infections, Ecuador

William J. Sears,¹ Jorge Cardenas,¹ Joseph Kubofcik, Thomas B. Nutman, Philip J. Cooper

Ancylostoma ceylanicum hookworms are zoonotic parasites that can infect humans. To detect autochthonous transmission, we analyzed human fecal samples collected in 2000. Multiparallel quantitative PCR detected infection in persons who had never traveled outside Ecuador. These data indicate human transmission of *A. ceylanicum* in the Americas, although endemicity remains unknown.

A*ncylostoma ceylanicum* hookworms can infect humans and have been increasingly recognized as endemic among humans living or traveling to the Asia-Pacific region (1). Although *A. ceylanicum* hookworms are known to infect dogs and cats globally, locally acquired human infections have not been reported by persons who have never traveled outside of North or South America. Previous evidence for the potential presence of *A. ceylanicum* hookworms in humans in the Americas is derived from 3 observations: 1) a 1922 autopsy study from northern Brazil identified zoonotic *Ancylostoma* spp. adult hookworms in human intestines when morphologic differentiation of *A. ceylanicum* from *A. braziliense* was still a matter of debate (2); 2) a molecular analysis of fecal samples from patients at hospitals in France identified *A. ceylanicum* hookworms in 2 samples, one in a migrant from Colombia and the other in a traveler returning from French Guiana (3); and 3) diffuse subacute unilateral neuroretinitis in a child adopted from Colombia and living in Germany was molecularly identified as *A. ceylanicum* infection (4). One report describes *A. ceylanicum* identified morphologically in domestic animal populations in South America that originated in Surinam, but no molecular methods were used (5).

Author affiliations: National Institutes of Health, Bethesda, Maryland, USA (W.J. Sears, J. Kubofcik, T.B. Nutman); University of Miami Miller School of Medicine, Miami, Florida, USA (J. Cardenas); Fundación Ecuatoriana Para Investigación en Salud, Quito, Ecuador (P.J. Cooper); Universidad Internacional del Ecuador, Quito (P.J. Cooper); St George's University of London, London, UK (P.J. Cooper)

Such reports, although suggestive, do not provide definitive evidence for autochthonous transmission of *A. ceylanicum* hookworms to humans in the Americas. We used molecular methods to identify autochthonous infections with *A. ceylanicum* in humans who had no history of travel outside Ecuador and who lived in a region of the country where soil-transmitted helminths are endemic. The study protocol was approved by the ethics committees of St George's Hospital Medical School, London, United Kingdom, and the Fundación Salud y Desarrollo Andino, Quito, Ecuador. Informed written consent was obtained from adult participants or from the parents or legal representatives of child participants.

The Study

In 2000, we collected fecal samples from a convenience subsample of preschool children, schoolchildren, and daycare staff in a larger cross-sectional study of the relationship between soil-transmitted helminth infections and allergy among urban and rural populations living in the Provinces of Pichincha and Esmeraldas, Ecuador. The subsamples represented settings considered to be at high risk for helminth infection (6). (Appendix Figure 1, <https://wwwnc.cdc.gov/EID/article/28/9/22-0248-App1.pdf>). We analyzed single frozen samples from 230 participants with a median age of 9 years (mean age 11, range <1 to 63 years); 40.4% were male and 69.6% were living in an urban setting (town of San Lorenzo). None of the study participants had a history of travel outside Ecuador.

The samples were thawed and ≈250 mg was transferred into collection microtubes in a 96-well format (QIAGEN, <https://www.qiagen.com>). We performed initial sample lysis by using a TissueLyzer II bead mill (QIAGEN) with 5-mm stainless steel beads. We performed high-throughput DNA extraction on a QiaSymphony (QIAGEN) instrument by using a PowerFecal Pro Kit (QIAGEN), following the manufacturer's instructions with minor modifications (i.e., samples

were pretreated with only the proprietary buffer CD1, and steps involving pretreatment with buffer CD2 were omitted to expedite the high-throughput protocol). All samples were extracted after an internal control was added as reported previously (7).

We prepared high-throughput, multiparallel quantitative PCR (qPCR) reactions by using 3.5 μ L of Taqman Fast Advanced Master Mix (QIAGEN), 2 μ L water, and 1 μ L of a PrimeTime qPCR Probe Assay (Integrated DNA Technologies, <https://www.idtdna.com>) with forward primer, reverse primer, and probe at a final reaction concentrations of 500 nmol/L, 500 nmol/L, and 250 nmol/L, respectively. A 96-well qPCR master plate was prepared for each respective target (*Ascaris lumbricoides/suum* [8], *A. ceylanicum* [9], *Strongyloides stercoralis* [10], *Trichuris trichiura* [11], *Necator americanus* [11], and *Ancylostoma duodenale* [11]), and 6.5 μ L of the qPCR solution was transferred to a 384-well plate with a Beckman Coulter liquid handler (Beckman Coulter, <https://www.beckmancoulter.com>) and subsequently frozen at -80°C until use. After DNA elution, we used the liquid handler to transfer 2 μ L of eluted DNA into a 384-well plate preloaded with the qPCR reagents. After brief centrifugation, the sample reactions were run on a ViiA7 Real Time PCR System (Thermo Fisher Scientific, <https://www.thermofisher.com>). Genomic DNA or plasmids served as positive reaction controls.

We first amplified the hookworm internal transcribed spacer (ITS) 1, 5.8s, and ITS2 regions by using primers and cycling conditions as described previously (12). After PCR amplification, we prepared the amplicon library for sequencing by using the

Nanopore Ligation Sequencing Kit (SQK-LSK109; Oxford Nanopore Technologies, <https://nanoporetech.com>) according to the manufacturer's instructions. The prepped library was loaded into an R9.4 flow cell and sequenced on a MinION device (Oxford Nanopore Technologies). We base called sequences by using Guppy Software, version 4.5.4 (Oxford Nanopore Technologies), and we determined consensus sequences by using NGSspeciesID with the following parameters: `-ont -sample size 300 -m 330 -s 50 -consensus -medaka (13)`. We performed alignment to reference analysis by using Geneious Prime 2021.02 (Biomatters, Ltd, <https://www.geneious.com>). In addition, the cyclooxygenase 1 region was PCR amplified by using previously reported primers and then Sanger sequenced (Quintara Biosciences, <https://quintarabio.com>) (14). We produced a phylogenetic tree of ITS and *cox1* sequences by using the maximum-likelihood method performed in Geneious Prime 2021.02 with bootstrapping $\times 1,000$ (15).

Of samples tested, hookworm infection was detected in 88 (38.3%) samples, of which 85 were infected with *N. americanus* (37.0%), 6 with *A. ceylanicum* (2.6%), and none with *A. duodenale* hookworms. Prevalence rates for other soil-transmitted helminth infections were *Ascaris lumbricoides* (63.9%) *T. trichiura* (69.7%), and *S. stercoralis* (15.2%). Cycle threshold values for *A. ceylanicum*-positive fecal samples detected by qPCR ranged from 25.2 to 37.6. We stratified distributions of sociodemographic factors and soil-transmitted helminth co-infections according to presence versus absence of *A. ceylanicum* or *N. americanus* hookworm infections (Table).

Table. Sociodemographic and soil-transmitted helminth co-infection characteristics of 230 persons for whom fecal samples were examined, Ecuador*

Variable	<i>Ancylostoma ceylanicum</i>		<i>Necator americanus</i>	
	Uninfected, n = 224	Infected, n = 6	Uninfected, n = 145	Infected, n = 85
Median age, y (range)	9 (<1–53)	4 (1–12)	10 (<1–63)	9 (<1–48)
Sex				
M, n = 93	89 (39.7)	4 (66.7)	54 (37.2)	46 (54.1)
F, n = 137	135 (60.3)	2 (33.3)	91 (62.8)	39 (45.9)
Residence				
Rural, n = 70	70 (31.2)	0	25 (17.2)	45 (52.9)
Urban, n = 160	154 (68.8)	6 (100)	120 (82.8)	40 (47.1)
STH co-infections				
<i>Ascaris lumbricoides</i> , n = 147	142 (63.4)	5 (83.3)	74 (51.0)	73 (85.8)
<i>Trichuris trichiura</i> , n = 160	154 (68.8)	6 (100)	94 (64.8)	66 (77.7)
<i>Strongyloides stercoralis</i> , n = 35	32 (14.3)	50.0 (3)	29 (20.0)	6 (7.1)
<i>N. americanus</i> , n = 85	82 (36.8)	50.0 (3)	NA	NA
<i>A. ceylanicum</i> , n = 6	NA	NA	3 (2.1)	6 (7.1)
No. co-infecting species				
0, n = 22	22 (9.8)	0	22 (15.2)	0
1, n = 62	62 (27.7)	0	59 (40.7)	3 (3.5)
≥ 2 , n = 146	140 (62.5)	6 (100)	64 (44.1)	82 (96.5)

*Values are no. (%) except as indicated. Data are stratified according to presence versus absence of *A. ceylanicum* or *N. americanus* hookworm infections. STH, soil-transmitted infection; NA, not applicable.

We successfully performed hookworm ITS1, 5.8S, ITS2 PCR amplification and amplicon sequencing on 2 of 6 *A. ceylanicum*-positive samples. The consensus sequences (GenBank accession nos. ON773142, ON773142) obtained from these 2 samples were >99% identical to the corresponding unique *A. ceylanicum* sequence targeted by the qPCR (Appendix Figure 2). We prepared a phylogenetic tree comparing known ITS1 sequences of *A. ceylanicum*, *A. duodenale*, *Ancylostoma braziliense*, and *Ancylostoma caninum* hookworms, including the consensus sequences from these 2 positive samples (referred to as being from subjects A and B; Appendix Figure 3). We also compared the phylogeny of *cox1* gene sequences for *A. ceylanicum* hookworms from this study (*A. ceylanicum* accession nos. ON773158) with isolates of the parasite from Asia (Appendix Figure 4).

Conclusions

We identified *A. ceylanicum* hookworm infection in 6 study participants who lived in an urban setting in Esmeraldas Province and had not traveled outside of Ecuador. All infections were detected in children in a marginalized urban setting where overcrowding is typical and hygiene standards are poor. Although the study was limited by small sample size and geographic restriction, the data provide evidence for autochthonous transmission of *A. ceylanicum* hookworm to humans in the Americas.

Funding for this study was provided by the National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, MD, USA, and Wellcome Trust, London, UK.

About the Author

Dr. Sears is an infectious disease physician-scientist in the Laboratory of Parasitic Disease at the US National Institute of Allergy and Infectious Disease, National Institutes of Health. His research interests include development of novel diagnostics and application of “omics” approaches to describe underlying mechanisms of parasitic phenomena.


References

1. Traub RJ, Zendejas-Heredia PA, Massetti L, Colella V. Zoonotic hookworms of dogs and cats - lessons from the past to inform current knowledge and future directions of research. *Int J Parasitol.* 2021;51:1233–41. <https://doi.org/10.1016/j.ijpara.2021.10.005>
2. Gordon RM. Ancylostomes recorded from sixty-seven post-mortems performed in Amazonas. *Ann Trop Parasitol.* 1922;16:223–8. <https://doi.org/10.1080/00034983.1922.11684313>
3. Gerber V, Le Govic Y, Ramade C, Chemla C, Hamane S, Desoubreux G, et al. *Ancylostoma ceylanicum* as the second most frequent hookworm species isolated in France in travellers returning from tropical areas. *J Travel Med.* 2021;28:taab014.
4. Poppert S, Heideking M, Agostini H, Fritzenwanker M, Wüppenhorst N, Muntau B, et al. Diffuse unilateral subacute neuroretinitis caused by *Ancylostoma* hookworm. *Emerg Infect Dis.* 2017;23:343–4. <https://doi.org/10.3201/eid2302.142064>
5. Rep BH, Heinemann DW. Changes in hookworm distribution in Surinam. *Trop Geogr Med.* 1976;28:104–10.
6. Chico ME, Vaca MG, Rodriguez A, Cooper PJ. Soil-transmitted helminth parasites and allergy: Observations from Ecuador. *Parasite Immunol.* 2019;41:e12590. <https://doi.org/10.1111/pim.12590>
7. Deer DM, Lampel KA, González-Escalona N. A versatile internal control for use as DNA in real-time PCR and as RNA in real-time reverse transcription PCR assays. *Lett Appl Microbiol.* 2010;50:366–72. <https://doi.org/10.1111/j.1472-765X.2010.02804.x>
8. Pilotte N, Maasch JRMA, Easton AV, Dahlstrom E, Nutman TB, Williams SA. Targeting a highly repeated germline DNA sequence for improved real-time PCR-based detection of *Ascaris* infection in human stool. *PLoS Negl Trop Dis.* 2019;13:e0007593. <https://doi.org/10.1371/journal.pntd.0007593>
9. Papaikakou M, Pilotte N, Grant JR, Traub RJ, Llewellyn S, McCarthy JS, et al. A novel, species-specific, real-time PCR assay for the detection of the emerging zoonotic parasite *Ancylostoma ceylanicum* in human stool. *PLoS Negl Trop Dis.* 2017;11:e0005734. <https://doi.org/10.1371/journal.pntd.0005734>
10. Basuni M, Muhi J, Othman N, Verweij JJ, Ahmad M, Miswan N, et al. A pentaplex real-time polymerase chain reaction assay for detection of four species of soil-transmitted helminths. *Am J Trop Med Hyg.* 2011;84:338–43. <https://doi.org/10.4269/ajtmh.2011.10-0499>
11. Pilotte N, Papaikakou M, Grant JR, Bierwert LA, Llewellyn S, McCarthy JS, et al. Improved PCR-based detection of soil transmitted helminth infections using a next-generation sequencing approach to assay design. *PLoS Negl Trop Dis.* 2016;10:e0004578. <https://doi.org/10.1371/journal.pntd.0004578>
12. Traub RJ, Inpankaew T, Sutthikornchai C, Sukthana Y, Thompson RC. PCR-based coprodiagnostic tools reveal dogs as reservoirs of zoonotic ancylostomiasis caused by *Ancylostoma ceylanicum* in temple communities in Bangkok. *Vet Parasitol.* 2008;155:67–73. <https://doi.org/10.1016/j.vetpar.2008.05.001>
13. Sahlin K, Lim MCW, Prost S. NGSspeciesID: DNA barcode and amplicon consensus generation from long-read sequencing data. *Ecol Evol.* 2021;11:1392–8. <https://doi.org/10.1002/ece3.7146>
14. Inpankaew T, Schär F, Dalsgaard A, Khieu V, Chimnoi W, Chhoun C, et al. High prevalence of *Ancylostoma ceylanicum* hookworm infections in humans, Cambodia, 2012. *Emerg Infect Dis.* 2014;20:976–82. <https://doi.org/10.3201/eid2006.131770>
15. Stamatakis A. RAxML-VI-HPC: maximum likelihood-based phylogenetic analyses with thousands of taxa and mixed models. *Bioinformatics.* 2006;22:2688–90. <https://doi.org/10.1093/bioinformatics/btl446>

Address for correspondence: William Sears, Bldg 4, Rm 211B, 4 Center Dr, National Institutes of Health, Bethesda, MD 20892-0425, USA; email: william.sears@nih.gov

Zoonotic *Ancylostoma ceylanicum* Hookworm Infections, Ecuador


Appendix




Appendix Figure 1. Locations of samples obtained from Ecuador, indicated by red and white dots.

DQ780009.1	CCTTACTGCTTGTGTTGGTGGTTGAGCATTAGGCTAACGCCTAGTGCGGCACCTGTCTGTCAGGAAACCTTAATG
Subject AC.....T.....
Subject BC.....T.....
DQ780009.1	ATCTGCTAACGCGGACGCCAGTACAGCAATAACTTTTTACGTTTAAATGTTTGAGAATCGTGACTTTATGTCACA
Subject A
Subject B
DQ780009.1	ATCGACTAGCTTCAGCGATGGATCGGTTCGATTGCGGTATCGATGAAAAACGCAGCTAGCTGCGTTATTTACCACG
Subject A
Subject B
DQ780009.1	AATTGCAGACGCTTAGAGTGGTGAAATTTTGAACGCATAGCGCCGTTGGGTTTTCCCTTCGGCACGTCTGGTTCA
Subject A
Subject B
DQ780009.1	GGGTTGTTTATATCTACTACAGTGTAGCTTGTGACA
Subject AC.....
Subject BC.....

Appendix Figure 2. The alignment of the *Ancylostoma ceylanicum* ITS reference sequence DQ780009.1 and the consensus sequence of the RTHW1 PCR amplicon. The sequences in the top, middle, and lower lane correspond to the amplified region of the reference (Accession #: DQ7800091), patient 5 sample (Acc #: ON773142), and patient 6 sample (Acc#: ON773143), respectively. Nucleotides discordant to the reference are indicated by their corresponding letter. Dots represent agreement with the reference.



Appendix Figure 3. A phylogenetic tree comparing sequences from *Ancylostoma ceylanicum* infections detected in subjects A and B samples with published ITS1 sequences of *A. ceylanicum*, *A. caninum*, *A. braziliensis*, and *A. duodenale*. The samples in blue are those of *A. ceylanicum* with origin. Sequences from subject samples A and B are indicated in red. Ex-Colombia refers to a sequence obtained from a Colombian migrant living in France.



Appendix Figure 4. A phylogenetic tree comparing Cox1 sequences of *Ancylostoma ceylanicum* from subjects A and B with various regions of Asia. The sequences from this study are in red.