DOI: 10.1289/EHP12141

**Note to readers with disabilities:** *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to 508 standards due to the complexity of the information being presented. If you need assistance accessing journal content, please contact <a href="ehp508@niehs.nih.gov">ehp508@niehs.nih.gov</a>. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

## **Supplemental Material**

Long-Term Exposure to Low-Level PM<sub>2.5</sub> and Mortality: Investigation of Heterogeneity by Harmonizing Analyses in Large Cohort Studies in Canada, United States, and Europe

Jie Chen, Danielle Braun, Tanya Christidis, Michael Cork, Sophia Rodopoulou, Evangelia Samoli, Massimo Stafoggia, Kathrin Wolf, Xiao Wu, Weiran Yuchi, Zorana J. Andersen, Richard Atkinson, Mariska Bauwelinck, Kees de Hoogh, Nicole A.H. Janssen, Klea Katsouyanni, Jochem O. Klompmaker, Doris Tove Kristoffersen, Youn-Hee Lim, Bente Oftedal, Maciej Strak, Danielle Vienneau, Jiawei Zhang, Richard T. Burnett, Gerard Hoek, Francesca Dominici, Michael Brauer, and Bert Brunekreef

## **Table of Contents**

- **Table S1.** Individual- and area-level covariates included in the main Model 4 in each cohort.
- **Table S2.** Characteristics of the full study populations in the stacked CanCHEC and ELAPSE cohorts.
- **Table S3.** Comparison of different exposure assessment methods applied in cohorts.
- Figure S1. Random-effects meta-analysis of eight cohorts.
- **Figure S2.** Cohort-specific concentration-response curve for PM<sub>2.5</sub> exposure and all-cause mortality in study populations aged 65 years and older.

## References