DOI: 10.1289/EHP14156

**Note to readers with disabilities:** *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to 508 standards due to the complexity of the information being presented. If you need assistance accessing journal content, please contact <a href="mailto:ehpsubmissions@niehs.nih.gov">ehpsubmissions@niehs.nih.gov</a>. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

## **Supplemental Material**

Associations between Aircraft Noise, Sleep, and Sleep–Wake Cycle: Actimetric Data from the UK Biobank Cohort near Four Major Airports

Xiangpu Gong, Katie Eminson, Glory O. Atilola, Calvin Jephcote, Kathryn Adams, Gabriella Captur, Andrew P. Hall, Marta Blangiardo, John Gulliver, Alex V. Rowlands, and Anna L. Hansell

## **Table of Contents**

**Table S1.** Sleep disturbance outcomes – name, definition, data range, and interpretation – as included in the UK Biobank questionnaire or actimetrically measured.

**Table S2.** Field IDs or return ID of the variables used in the study and their descriptions.

## Main models - STATA code

**Figure S1.** Directed Acyclic Graph (DAG) identifying potential confounders for the relationship between night-time aircraft noise exposure and sleep disturbance.

## **Sensitivity Analysis**

**Figure S2.** Cross-sectional association between night-time aircraft noise and average acceleration during the least active 8 hours while excluding subgroups susceptible to noise misclassification using UK Biobank cohort.

**Figure S3.** Cross-sectional association between night-time aircraft noise and proportion of time spent on sleep or in bed (7-day average) while excluding subgroups susceptible to noise misclassification using UK Biobank cohort.

**Figure S4.** Cross-sectional association between night-time aircraft noise and relative amplitude while excluding subgroups susceptible to noise misclassification using UK Biobank cohort.

- **Figure S5.** Cross-sectional association between night-time aircraft noise and intra-daily variability while excluding subgroups susceptible to noise misclassification using UK Biobank cohort.
- **Figure S6.** Cross-sectional association between night-time aircraft noise and inter-daily stability while excluding subgroups susceptible to noise misclassification using UK Biobank cohort.
- **Figure S7.** Repeated cross-sectional association between night-time aircraft noise and self-reported sleeplessness while excluding subgroups susceptible to noise misclassification using UK Biobank cohort.
- **Figure S8.** Repeated cross-sectional association between night-time aircraft noise and self-reported daytime dozing while excluding subgroups susceptible to noise misclassification using UK Biobank cohort.
- **Figure S9.** Repeated cross-sectional association between night-time aircraft noise and self-reported sleep duration while excluding subgroups susceptible to noise misclassification using UK Biobank cohort.
- **Figure S10.** Cross-sectional association between night-time aircraft noise and average acceleration during the least active 8 hours among some demographic or socioeconomic groups using UK Biobank cohort.
- **Figure S11.** Cross-sectional association between night-time aircraft noise and proportion of time spent on sleep or in bed (7-day average) among some demographic or socioeconomic groups using UK Biobank cohort.
- **Figure S12.** Cross-sectional association between night-time aircraft noise and relative amplitude among some demographic or socioeconomic groups using UK Biobank cohort.
- **Figure S13.** Cross-sectional association between night-time aircraft noise and intra-daily variability among some demographic or socioeconomic groups using UK Biobank cohort.
- **Figure S14.** Cross-sectional association between night-time aircraft noise and inter-daily stability among some demographic or socioeconomic groups using UK Biobank cohort.
- **Figure S15.** Repeated cross-sectional association between night-time aircraft noise and self-reported sleeplessness among some demographic or socioeconomic groups using UK Biobank cohort.
- **Figure S16.** Repeated cross-sectional association between night-time aircraft noise and self-reported daytime dozing among some demographic or socioeconomic groups using UK Biobank cohort.

- **Figure S17.** Repeated cross-sectional association between night-time aircraft noise and self-reported sleep duration among some demographic or socioeconomic groups using UK Biobank cohort.
- **Figure S18.** Cross-sectional association between night-time aircraft noise and average acceleration during the least active 8 hours among people with health conditions linked to sleep disturbance using UK Biobank cohort.
- **Figure S19.** Cross-sectional association between night-time aircraft noise and proportion of time spent on sleep or in bed (7-day average) among people with health conditions linked to sleep disturbance using UK Biobank cohort.
- **Figure S20.** Cross-sectional association between night-time aircraft noise and relative amplitude among people with health conditions linked to sleep disturbance using UK Biobank cohort.
- **Figure S21.** Cross-sectional association between night-time aircraft noise and intra-daily variability among people with health conditions linked to sleep disturbance using UK Biobank cohort.
- **Figure S22.** Cross-sectional association between night-time aircraft noise and inter-daily stability among people with health conditions linked to sleep disturbance using UK Biobank cohort.
- **Figure S23.** Repeated cross-sectional association between night-time aircraft noise and self-reported sleeplessness among people with health conditions linked to sleep disturbance using UK Biobank cohort.
- **Figure S24.** Repeated cross-sectional association between night-time aircraft noise and self-reported daytime dozing among people with health conditions linked to sleep disturbance using UK Biobank cohort.
- **Figure S25.** Repeated cross-sectional association between night-time aircraft noise and self-reported sleep duration among people with health conditions linked to sleep disturbance using UK Biobank cohort.

**Additional File-** Excel Document