

The Journal of Urology
Pinpointing the urological risk in open spina bifida
--Manuscript Draft--

Manuscript Number:	
Article Type:	Letter to the Editor
Corresponding Author:	Pippa Oakeshott, MD St George's University of London London, London UNITED KINGDOM
First Author:	Pippa Oakeshott, MD
Order of Authors:	Pippa Oakeshott, MD
	Alison Poulton, MD
	Gillian M Hunt, MD

Pinpointing the urological risk in open spina bifida

In a recent cross-sectional study from The Netherlands, Veenboer and colleagues report that adult patients with spina bifida who walk into the consulting room (as opposed to those who use a wheelchair) are unlikely to have unfavourable urodynamic findings¹. This useful observation may also apply to children. In the Cambridge cohort of 117 consecutive, unselected patients born with open spina bifida between 1963 and 1971², 75 survived to the mean age of 9 years at which time 15 (20% of survivors) walked independently. None of these walkers subsequently died of urological causes over the next 37 years compared with 20% (12/60) of the remainder.

However neurological level in infancy in terms of sensory loss to pinprick is a better predictor of an adverse urological outcome than walking, or bony level, cutaneous level or motor level³. Using 2013 data from the Cambridge cohort³ (which has no loss to follow up), Figure 1 shows that urological deaths occurred only in those born with a sensory level of T5-L1 ($p < 0.001$). This consistent relationship between a high sensory level in infancy (recorded by the neurosurgeon) and subsequent urological death persisted despite a range of urological management and operations carried out at many different hospitals over a period of nearly 50 years.

We also found clustering of urological deaths. Of 31 patients born with a sensory level of T9 or T10, 14 (45%) died of urological causes. This interesting finding deserves further consideration. Anaesthetic skin in the thoracic or first lumbar dermatomes might indicate a neuropathic renal tract, with both skin and kidneys vulnerable to disease mediated by the common risk factor of interruption of their nerve supply.

Finally, sensory level may be related to congenital renal abnormalities. In another study of 190 patients with open spina bifida⁴, renal agenesis was associated with a sensory level between T5-8, horseshoe kidney with T9-L1 and ureteral duplication with a sacral sensory level. Sensory loss and renal abnormalities may reflect the level at which neurulation failed⁵.

We agree with Veenboer and colleagues that urological review in patients with open spina bifida should focus on the ones who are most at risk¹. In practice this means non-walkers and those with a sensory loss to pin prick extending to (or above) the first lumbar dermatome.

Pippa Oakeshott¹ oakeshot@sgul.ac.uk

Alison Poulton² alison.poulton@sydney.edu.au

Gillian M Hunt³ jillhuntpoulton@btinternet.com

1. **Population Health Sciences, St George's, University of London, UK.**
2. **Department of Paediatrics, Sydney Medical School Nepean, University of Sydney, Australia**
3. **65 Grantchester St, Cambridge**

Correspondence to Professor Oakeshott oakeshot@sgul.ac.uk

Funding: Newlife Foundation for Disabled Children. SG/11-12/06

Reference List

- (1) Veenboer PW, Bosch JL, Rosier PF, Dik P, van Asbeck FW, de Jong TP et al. Cross-sectional study of determinants of upper and lower urinary tract outcomes in adults with spinal dysraphism--new recommendations for urodynamic followup guidelines? *J Urol* 2014; 192(2):477-482.
- (2) Hunt GM, Lewin L, Gleave J, Gairdner D. Predictive factors in open myelomeningocele with special reference to sensory level. *BMJ* 1973; 4:187-201.
- (3) Oakeshott P, Reid F, Poulton A, Markus H, Whitaker RH, Hunt GM. Neurological Deficit at Birth Predicts Survival to the Mid-40s and Urological Deaths in Open Spina Bifida: A Complete Prospective Cohort Study. *Dev Med Child Neurol*. 2015. DOI: 10.1111/dmcn.12698
- (4) Hunt GM, Whitaker RH. The pattern of congenital renal anomalies associated with neural-tube defects. *Dev Med Child Neurol* 1987; 29(1):91-95.
- (5) Copp AJ, Stanier P, Greene ND. Neural tube defects: recent advances, unsolved questions, and controversies. *Lancet Neurol* 2013; 12(8):799-810.

Figure 1. Deaths from urological causes and sensory level recorded in infancy in a complete cohort of 117 consecutive cases of open spina bifida

