

40. Gemert vL, Schuurmans M. The Neecham Confusion Scale and the Delirium Observation Screening Scale. *BMC Nursing* 2007; 6: 3.
41. Godden S, Pollock AM. The use of acute hospital services by elderly residents of nursing and residential care homes. *Health Soc Care Community* 2001; 9: 367–74.
42. Coast J, Peters TJ, Natarajan L, Sproston K, Flynn T. An assessment of the construct validity of the descriptive system for the ICECAP capability measure for older people. *Qual Life Res* 2008; 17: 967–76.
43. Bulamu NB, Kaambwa B, Ratcliffe J. A systematic review of instruments for measuring outcomes in economic evaluation within aged care. *Health Qual Life Outcomes* 2015; 13: 179.

Received 24 December 2015; accepted in revised form 1 April 2016

*Age and Ageing* 2016; **45**: 661–666  
doi: 10.1093/ageing/afw080  
Published electronically 4 May 2016

© The Author 2016. Published by Oxford University Press on behalf of the British Geriatrics Society.  
This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

## Hearing impairment and incident disability and all-cause mortality in older British community-dwelling men

ANN E. M. LIJAS<sup>1</sup>, S. GOYA WANNAMETHEE<sup>1</sup>, PETER H. WHINCUP<sup>2</sup>, OLIA PAPACOSTA<sup>1</sup>, KATE WALTERS<sup>1</sup>, STEVE ILIFFE<sup>1</sup>, LUCY T. LENNON<sup>1</sup>, LIVIA A. CARVALHO<sup>3</sup>, SHEENA E. RAMSAY<sup>1</sup>

<sup>1</sup>Department of Primary Care and Population Health, University College London, London NW3 2PF, UK

<sup>2</sup>Population Health Research Centre, Division of Population Health Sciences and Education, St George's, University of London, London, UK

<sup>3</sup>Research Department of Epidemiology and Public Health, University College London, London, UK

Address correspondence to: A. E. M. Liljas. Tel: (+44) 2077940500; Fax (+44) 2077941224. Email: [ann.liljas.13@ucl.ac.uk](mailto:ann.liljas.13@ucl.ac.uk)

### Abstract

**Background and objective:** hearing impairment is common in older adults and has been implicated in the risk of disability and mortality. We examined the association between hearing impairment and risk of incident disability and all-cause mortality.

**Design and setting:** prospective cohort of community-dwelling older men aged 63–85 followed up for disability over 2 years and for all-cause mortality for 10 years in the British Regional Heart Study.

**Methods:** data were collected on self-reported hearing impairment including hearing aid use, and disability assessed as mobility limitations (problems walking/taking stairs), difficulties with activities of daily living (ADL) and instrumental ADL (IADL). Mortality data were obtained from the National Health Service register.

**Results:** among 3,981 men, 1,074 (27%) reported hearing impairment. Compared with men with no hearing impairment, men who could hear and used a hearing aid, and men who could not hear despite a hearing aid had increased risks of IADL difficulties (age-adjusted OR 1.86, 95% CI 1.29–2.70; OR 2.74, 95% CI 1.53–4.93, respectively). The associations remained after further adjustment for covariates including social class, lifestyle factors, co-morbidities and social engagement. Associations of hearing impairment with incident mobility limitations, incident ADL difficulties and all-cause mortality were attenuated on adjustment for covariates.

**Conclusion:** this study suggests that hearing problems in later life could increase the risk of having difficulties performing IADLs, which include more complex everyday tasks such as shopping and light housework. However, further studies are needed to determine the associations observed including the underlying pathways.

**Keywords:** ageing, older adults, hearing impairment, disability, mortality

## Introduction

Hearing impairment increases with age and has been associated with chronic conditions including cardiovascular disease (CVD) and poor quality of life [1, 2]. Demographic changes mean that people are living longer with chronic diseases and associated physical limitations affecting independent living and overall well-being [3]. Disability in later life often occurs first as mobility limitations (for example, difficulties walking or climbing stairs) [4]. Other forms of disability refer to disablement in tasks essential to caring for oneself (basic activities of daily living [ADL], e.g. bathing, dressing) and more complex tasks that refer to living independently in the community (instrumental activities of daily living [IADL], e.g. shopping, telephoning) [4]. Earlier studies have shown an association between hearing impairment and mobility limitations [5] including increased risks of subsequent mobility limitations among older adults with hearing impairment [6, 7]. Previous research has also shown an association between hearing impairment and incident ADL deficits in hospital-based samples of older people [8, 9]. Hearing impairment has furthermore been associated with an increased risk of incident IADL; however, findings are inconsistent [6, 10–12]. In addition, it has been suggested that hearing impairment increases the risk of all-cause mortality [2], but some studies have shown no association after adjustment for demographic factors, physical functioning and cognition [12, 13].

It is important to understand the influence of hearing impairment on disability including activities of daily living to establish the impact of hearing impairment on functional independence in later life. Therefore, we investigated the association of self-reported hearing impairment with subsequent mobility limitations, ADL, IADL and all-cause mortality in a representative sample of older British men aged 63–85 followed up for 2 years for disability and 10 years for mortality. We also examined whether these associations were independent of age, social class, lifestyle factors and co-morbidities known to be associated with hearing impairment including CVD, hypertension and diabetes [1, 14].

## Methods

### Study design and participants

This study uses data from the British Regional Heart Study (BRHS), a prospective study in a socioeconomically and geographically representative sample of 7,735 middle-aged men drawn from 24 general practices representing all major British Regions [15]. The men were recruited in 1978–80 and have regularly been followed up since. For this study, baseline data on 3,981 men, then aged 63–85, were obtained through self-reported questionnaires in 2003. Ethical approval was obtained from relevant local research ethics committees.

### Hearing impairment

Questions on hearing impairment included ‘Do you use a hearing aid?’ and ‘Using a hearing aid if needed, is your hearing good enough to follow a TV programme at a volume others find acceptable?’ with answer options yes/no [16], and allowed for participants to be divided into four groups: could follow TV and used no hearing aid (could hear, no aid [no hearing impairment]) (reference group), could follow TV and used hearing aid (could hear, used aid), could not follow TV and did not use hearing aid (could not hear, no aid), and could not follow TV and used hearing aid (could not hear, used aid).

### Outcome measures

All men were followed up for mobility limitations and activities of daily living (ADL) and instrumental ADL (IADL) from 2003 to 2005 and for all-cause mortality from 2003 to 2013. Two questions asking whether they had problems taking the stairs and problems walking 400 yards with answer options yes/no were used to assess mobility limitation. Reporting problems with one or both was classified as having mobility limitations. ADL was classified as having some difficulty or in need of help undertaking one or more of the following activities: bathing, dressing, eating, getting in or out of bed or chair, toileting and/or walking across a room [17]. IADL was based on reporting some problem or in need of help undertaking cooking, shopping, using public transport, managing money and/or using the telephone [18]. Incidence was determined as having no previous mobility limitation, ADL and IADL, respectively. Mortality data were collected from the National Health Service register. Binary outcomes assessed in the current analyses were for incident mobility limitations, difficulties in ADL, IADL and all-cause mortality.

### Covariates

Covariates included socioeconomic and lifestyle factors including social class, social engagement, cigarette smoking, obesity and physical activity. Co-morbidity-related covariates included doctor-diagnosed CVD (coronary thrombosis, myocardial infarction, angina and/or stroke), hypertension and diabetes analysed dichotomously. Participants were divided into manual and non-manual social class based on the longest held occupation of subjects at study entry using the Registrar Generals’ Social Class Classification. The men were grouped into non-smokers, ex-smokers and current smokers. Being obese was defined as having a body mass index (BMI) of 30 kg/m<sup>2</sup> and over [19]. Physical activity scores were based on exercise type and frequency categorised as none, occasional, light, moderate, moderately vigorous and vigorous [20], where none or occasional activity was classified as being inactive. Other covariates included social engagement, doctor-diagnosed depression and difficulty keeping balance. Low social engagement was classified

as doing three or fewer activities part of a 9-item social engagement scale on a weekly basis: voluntary work, go to the pub or a club, attend religious services, play cards or games, visit the cinema, restaurants or sports events, attend a class or course of study, and, sometimes go on day or overnight trips, and been on a holiday in the last year [21]. Depression and reporting not being able to keep balance were analysed dichotomously. Data on all covariates were collected at baseline (2003).

**Statistical analyses**

Logistic regression was used to assess the associations of hearing impairment with incident mobility limitations and difficulties in ADLs and IADLs. Odds ratios (OR) with 95% confidence intervals (CI) were obtained using no hearing impairment (could hear, no aid) as reference group. Survival analysis was used to examine the association between hearing impairment and mortality, and Cox proportional hazards regression was used to calculate hazard ratios (HR) with 95% CIs. We also performed Cox regression using age as the time variable. The test confirmed that the proportionality hazards assumption was met. Participants who did not answer any of the hearing-related questions (*n* = 38) were excluded. Participants free from mobility limitations and difficulties in ADLs and IADLs at baseline were followed up for each of these types of disability. Models were adjusted for age, social class, lifestyle factors and co-morbidities. A Bonferroni correction was applied for multiple comparisons, and findings remained significant. All analyses were carried out using SAS version 9.3 software (SAS Institute, Inc., Cary, NC, USA).

**Results**

In 2003, 3,981 men aged 63–85 completed the questionnaire (82% response rate). Of these, 3,108 men had no previous mobility limitations, 3,346 men had no previous ADL

and 3,410 men had no previous IADL. At 2-year follow-up, there were 238 (8%) new cases of mobility limitations, 260 (8%) new cases of ADL and 207 (6%) new cases of IADL. All 3,981 men were also followed for all-cause mortality over 10 years during which 1,463 (37%) deaths occurred. Characteristics of participants by hearing impairment are shown in Table 1. Table 2 presents odds ratios (OR) with 95% CIs for incident mobility limitations, ADL and IADL for hearing impairment. Compared with men with no hearing impairment, men who could not hear and used a hearing aid had over a twofold greater risk of mobility limitations at 2-year follow-up (age-adjusted OR 2.24, 95% CI 1.29–3.89). The association remained after further adjustment for social class, lifestyle factors and co-morbidities (OR 1.89, 95% CI 1.04–3.41) but was attenuated upon adjustment for social engagement. Men who could not hear, irrespective of using hearing aid, had greater risks of developing problems performing ADL compared with men with no hearing impairment (OR 1.74, 95% CI 1.19–2.55; OR 2.01, 95% CI 1.16–3.46). The association was attenuated after further adjustment among men who used an aid but remained in those who could not hear and did not use hearing aid even after further adjustment for social engagement (OR 1.68, 95% CI 1.11–2.55). However, the association was attenuated after further adjustment for mobility limitations (OR 1.49, 95% CI 0.97–2.29). Compared with men with no hearing impairment, those who could hear and used a hearing aid and those who could not hear despite aid were more likely to develop IADL problems (OR 1.86, 95% CI 1.29–2.70; OR 2.74, 95% CI 1.53–4.93). These associations are of particular interest as they remained after further adjustment including social engagement (OR 2.00, 95% CI 1.34–2.99; OR 2.61, 95% CI 1.38–4.96) and also after further adjustment for mobility limitations, depression and poor balance (OR 2.03, 95% CI 1.35–3.07; OR 2.77, 95% CI 1.43–5.36). Further analyses of the associations between hearing impairment and individual components of IADL showed that men who

**Table 1.** Percentages and numbers for socioeconomic and lifestyle characteristics, co-morbidities and mean age by hearing impairment in a cohort of British men aged 63–85 in 2003

% ( <i>n</i> )	Overall	No hearing impairment		Hearing impairment groups		<i>P</i> -value
		Could hear, no aid	Could hear, used aid	Could not hear, no aid	Could not hear, used aid	
Totals	100 (3,981)	73 (2,851)	12 (482)	11 (424)	4 (168)	
Covariates						
Manual social class	51 (1,962)	48 (1,317)	53 (245)	63 (263)	60 (98)	<0.01
Current smokers	10 (389)	10 (284)	7 (33)	13 (54)	9 (15)	0.67
Ex-smokers	60 (2,385)	59 (1,681)	66 (314)	58 (244)	69 (115)	
Never smoked	30 (1,174)	31 (870)	27 (131)	29 (123)	22 (37)	
Physical inactivity	38 (1,430)	36 (971)	44 (196)	41 (157)	54 (87)	0.06
Obese	17 (639)	16 (445)	13 (61)	21 (85)	26 (42)	<0.01
CVD	27 (1,087)	26 (728)	32 (153)	26 (112)	43 (72)	<0.01
Hypertension	39 (1,547)	38 (1,092)	41 (196)	39 (165)	46 (78)	0.24
Diabetes	10 (393)	10 (281)	9 (45)	10 (41)	11 (19)	0.89
Age						
Mean age in years ± SD	72 (5.4)	72 (5.3)	75 (5.4)	72 (5.4)	74 (5.4)	

**Table 2.** Odds ratios (OR) with 95% CIs for associations between incidence of mobility limitations, ADL and IADL and hearing impairment in British men aged 63–85 in 2003 followed up for 2 years to 2005

		No hearing impairment		Hearing impairment	
		Could hear, no aid	Could hear, used aid	Could not hear, no aid	Could not hear, used aid
.....		.....		.....	
Limitations in mobility	<i>n</i> (%)	150 (7)	39 (11)	23 (8)	17 (16)
Model 1	OR (95% CI)	1.00	1.40 (0.95–2.05)	1.16 (0.73–1.83)	2.24 (1.29–3.89)
Model 2		1.00	1.40 (0.92–2.12)	1.26 (0.78–2.03)	1.89 (1.04–3.41)
Model 2 + social engagement <sup>a</sup>		1.00	1.41 (0.93–2.14)	1.24 (0.77–2.01)	1.79 (0.98–3.27)
ADL		161 (7)	41 (10)	37 (11)	17 (15)
Model 1		1.00	1.30 (0.90–1.88)	1.74 (1.19–2.55)	2.01 (1.16–3.46)
Model 2		1.00	1.23 (0.82–1.84)	1.76 (1.16–2.66)	1.62 (0.90–2.94)
Model 2 + social engagement <sup>a</sup>		1.00	1.25 (0.83–1.87)	1.68 (1.11–2.55)	1.59 (0.87–2.88)
IADL		126 (5)	44 (11)	19 (6)	15 (15)
Model 1		1.00	1.86 (1.29–2.70)	1.09 (0.66–1.79)	2.74 (1.53–4.93)
Model 2		1.00	2.03 (1.36–3.01)	1.01 (0.59–1.75)	2.56 (1.35–4.86)
Model 2 + social engagement <sup>a</sup>		1.00	2.00 (1.34–2.99)	0.95 (0.54–1.67)	2.61 (1.38–4.96)
IADL components					
Shopping		73 (3)	31 (7)	17 (5)	13 (9)
Model 1		1.00	2.05 (1.32–3.20)	1.63 (0.95–2.80)	2.80 (1.50–5.23)
Model 2		1.00	1.96 (1.20–3.19)	1.56 (0.87–2.82)	2.39 (1.22–4.68)
Model 2 + social engagement <sup>a</sup>		1.00	2.01 (1.23–3.28)	1.46 (0.80–2.68)	2.30 (1.15–4.60)
Light housework		66 (2)	25 (6)	12 (3)	12 (8)
Model 1		1.00	1.93 (1.19–3.12)	1.24 (0.66–2.32)	3.08 (1.61–5.88)
Model 2		1.00	1.76 (1.05–2.95)	1.05 (0.54–2.05)	2.73 (1.39–5.34)
Model 2 + social engagement <sup>a</sup>		1.00	1.80 (1.07–3.04)	1.02 (0.52–2.00)	2.73 (1.39–5.38)
Telephoning		43 (2)	17 (4)	7 (2)	11 (8)
Model 1		1.00	1.85 (1.03–3.32)	1.10 (0.49–2.47)	4.53 (2.25–9.10)
Model 2		1.00	1.64 (0.88–3.04)	0.75 (0.29–1.93)	3.82 (1.80–8.09)
Model 2 + social engagement <sup>a</sup>		1.00	1.74 (0.93–3.24)	0.78 (0.30–2.03)	4.29 (2.02–9.13)
Managing money		59 (2)	16 (4)	10 (3)	14 (9)
Model 1		1.00	1.27 (0.71–2.25)	1.13 (0.57–2.23)	3.68 (1.99–6.82)
Model 2		1.00	1.29 (0.71–2.35)	0.97 (0.45–2.07)	3.49 (1.84–6.62)
Model 2 + social engagement <sup>a</sup>		1.00	1.32 (0.72–2.41)	0.95 (0.44–2.04)	3.68 (1.94–6.98)
Using public transport		75 (3)	33 (8)	13 (4)	7 (5)
Model 1		1.00	1.98 (1.28–3.06)	1.20 (0.66–2.20)	1.42 (0.64–3.19)
Model 2		1.00	1.97 (1.23–3.16)	1.16 (0.61–2.20)	1.33 (0.58–3.05)
Model 2 + social engagement <sup>a</sup>		1.00	1.93 (1.20–3.11)	1.13 (0.59–2.14)	1.36 (0.60–3.13)

Model 1: adjusted for age; Model 2: adjusted for age, social class, BMI, smoking, physical activity, CVD, hypertension and diabetes.

<sup>a</sup>Social engagement was defined as doing three or fewer of the following activities on a weekly basis: voluntary work, go to the pub or a club, attend religious services, play cards or games, visit the cinema, restaurants or sports events, attend a class or course of study, and, sometimes go on day or overnight trips, and been on a holiday in the last year.

**Table 3.** Hazard ratios (HR) with 95% CIs for associations between all-cause mortality and hearing impairment in British men aged 63–85 in 2003 followed up for 10 years to 2013

		No hearing impairment		Hearing impairment	
		Could hear, no aid	Could hear, used aid	Could not hear, no aid	Could not hear, used aid
.....		.....		.....	
	Rates/1,000 ( <i>n</i> )	39 (974)	54 (216)	48 (169)	58 (76)
Model 1	HR (95% CI)	1.00	1.03 (0.88–1.19)	1.19 (1.01–1.40)	1.18 (0.93–1.49)
Model 2		1.00	1.01 (0.86–1.19)	1.12 (0.93–1.34)	1.14 (0.89–1.45)

Model 1: adjusted for age; Model 2: adjusted for age, social class, BMI, smoking, physical activity, CVD, hypertension and diabetes.

could hear and used a hearing aid and men who could not hear despite aid were both more likely to experience problems, in particular undertaking shopping and light housework, even after further adjustment including social engagement. Men who could hear and used aid were also more likely to have problems using public transport. Only those who could not hear despite aid had increased risks of

difficulty cooking (OR 2.03, 95% CI 1.05–3.94), but the association was attenuated after full adjustment (results not presented). Men who could hear and used aid and men who could not hear despite aid were more likely to have problems telephoning with over fourfold increased risks in men who could not hear despite aid (OR 4.53, 95% CI 2.25–9.10). The association remained in men who could



not hear despite aid after further adjustment including social engagement (OR 4.29, 95% CI 2.02–9.13) and after further adjustment for mobility limitations, depression and poor balance (OR 4.29, 95% CI 2.00–9.18). The associations between hearing impairment and IADL were further analysed without the component of difficulty telephoning. Age-adjusted findings showed that men who could hear and used a hearing aid and men who could not hear despite an aid were more likely to develop difficulties in IADLs (results not presented). The association remained in men who could hear with aid only, after full adjustment including social engagement. None of the hearing impairment groups were associated with difficulties taking medications (results not presented).

Table 3 shows hazard ratios (HR) with 95% CIs for all-cause mortality associated with hearing impairment. Compared with men with no hearing impairment, those who could not hear and did not use a hearing aid had a significantly greater risk of all-cause mortality (HR 1.19, 95% CI 1.01–1.40) but the association was attenuated on further adjustments. No other hearing impairment group was associated with increased risk of all-cause mortality. These findings were confirmed when using age as the time scale.

### Discussion

This study investigated the association of hearing impairment with incident disability (mobility limitations, ADL, IADL) and all-cause mortality in older British men. Our findings show that men with hearing impairment had greater risks in particular of disability affecting IADLs. The associations observed between hearing impairment and incident mobility limitations, incident ADL and all-cause mortality were attenuated on further adjustment for covariates.

The association between hearing impairment and mobility limitations was attenuated particularly on adjustment for social engagement. Communication problems due to hearing impairment may restrict social engagement [7]. Being socially engaged can motivate maintenance of physical functioning [22], reducing the risk of incident disability [23]. Only men who could not hear and did not use a hearing aid had greater risk of subsequent ADL deficits after adjustment including social engagement. However, the association was attenuated after further adjustment for mobility limitations. In contrast, men who could hear with an aid and men who could not hear despite an aid had increased risks of subsequent IADL difficulties and the associations remained after further adjustment. The associations also remained statistically significant after further adjustment for mobility limitations, depression and poor balance. This suggests that hearing impairment has a greater impact on IADLs which involve more complex tasks (such as shopping and light housework) than basic tasks including ADL and mobility limitations [24]. However, this finding should be interpreted with caution as the association between not being able to hear despite an aid and subsequent IADL was driven by difficulty telephoning. Also, the observed associations between hearing impairment and

IADL could be explained by residual confounding due to unmeasured factors such as cognitive functioning, which is important for complex IADL tasks [13, 25]. The degradation hypothesis suggests that a decline in hearing impairment in older age increases the demands on cognitive functioning [8]. Previous research also suggests that family members may steer older relatives with poor physical and cognitive functions away from responsibilities and tasks such as IADLs [8]. Further, lack of consistent findings across the hearing impairment groups and incident IADL with no association observed in those ‘unable to hear, no aid’ suggests that this group may consist of a combination of men with a hearing problem who did not use a hearing aid due to, for instance, lack of access to health services and audiology assessments, reluctance to wear an aid, a perception that aids are unhelpful and men whose hearing problem is not improved by an aid. Finally, the association observed could be due to inflammation, which is related to both hearing impairment and disability [26, 27].

Men who could not hear and did not use hearing aid had greater risks of all-cause mortality compared with men with no hearing impairment. However, the association was attenuated after adjustment for social class, lifestyle factors and co-morbidities. This is consistent with earlier studies demonstrating no association after adjustment for potential confounders including social class and physical functioning [12, 13].

### Strengths and limitations

The major strengths of this study are that it was a large socioeconomically representative cohort with negligible loss to follow-up for disability and mortality [15]. In addition, the cohort was followed up for 2 years for disability and for 10 years for mortality, and the models were adjusted for several confounding variables.

Limitations include that hearing impairment was self-reported rather than objectively measured. However, the questions used have been validated against objective measures [16]. Furthermore, previous research has demonstrated comparable findings when investigating both self-reported (defined as ever had deafness or trouble hearing with one or both ears) and measured hearing impairment and 10-year mortality risk [12]. Further, the question on hearing aid use did not specify whether the participants have been offered a hearing aid and chosen not to use it or whether they do not have a hearing aid at all. Further, despite the large sample, the number of participants in each hearing impairment group with disability was small which might have reduced the statistical power of the study. Furthermore, hearing impairment was measured at baseline only, and no information on the primary cause of and change in hearing impairment were investigated. Finally, the study was in older men, predominantly of white British ethnic origin, and generalisation of findings to women and to other ethnic groups is limited.

## Conclusions and implications

In summary, our study shows that older men who could follow TV and used a hearing aid have greater risks of disability affecting IADLs, which are important for maintaining functional independence in later life. The inconsistent findings across the hearing impairment groups further suggest that it may not be hearing *per se* underlying the association. Future longitudinal studies are required to further assess the association between hearing impairment and incident disability, taking cognitive impairment and inflammation into account.

## Key points

- Hearing problems in later life may increase the risk of having difficulties performing IADLs.
- Inconsistent findings across the hearing impairment groups suggest that something may be underlying the association with IADLs.
- The association of hearing impairment and all-cause mortality was attenuated on adjustment for covariates.

## Conflicts of interest

None declared.

## References

1. Liljas AE, Wannamethee SG, Whincup PH *et al.* Socio-demographic characteristics, lifestyle factors and burden of morbidity associated with self-reported hearing and vision impairments in older British community-dwelling men: a cross-sectional study. *J Public Health (Oxf)* 2015; doi:10.1093/pubmed/fdv095.
2. Fisher D, Li CM, Chiu MS *et al.* Impairments in hearing and vision impact on mortality in older people: the AGES-Reykjavik Study. *Age Ageing* 2014; 43: 69–76.
3. Ebrahim S, Wannamethee SG, Whincup P *et al.* Locomotor disability in a cohort of British men: the impact of lifestyle and disease. *Int J Epidemiol* 2000; 29: 478–86.
4. Harris T, Kovar MG, Suzman R *et al.* Longitudinal study of physical ability in the oldest-old. *Am J Public Health* 1989; 79: 698–702.
5. Chen DS, Genther DJ, Betz J *et al.* Association between hearing impairment and self-reported difficulty in physical functioning. *J Am Geriatr Soc* 2014; 62: 850–6.
6. Wallhagen MI, Strawbridge WJ, Shema SJ *et al.* Comparative impact of hearing and vision impairment on subsequent functioning. *J Am Geriatr Soc* 2001; 49: 1086–92.
7. Chen DS, Betz J, Yaffe K *et al.* Association of hearing impairment with declines in physical functioning and the risk of disability in older adults. *J Gerontol A Biol Sci Med Sci* 2015; 70: 654–61.
8. Yamada M, Nishiwaki Y, Michikawa T *et al.* Self-reported hearing loss in older adults is associated with future decline in instrumental activities of daily living but not in social participation. *J Am Geriatr Soc* 2012; 60: 1304–9.

9. Keller BK, Morton JL, Thomas VS *et al.* The effect of visual and hearing impairments on functional status. *J Am Geriatr Soc* 1999; 47: 1319–25.
10. Rosso AL, Eaton CB, Wallace R *et al.* Geriatric syndromes and incident disability in older women: results from the women's health initiative observational study. *J Am Geriatr Soc* 2013; 61: 371–9.
11. Rudberg MA, Furner SE, Dunn JE *et al.* The relationship of visual and hearing impairments to disability: an analysis using the longitudinal study of aging. *J Gerontol* 1993; 48: M261–5.
12. Reuben DB, Mui S, Damesyn M *et al.* The prognostic value of sensory impairment in older persons. *J Am Geriatr Soc* 1999; 47: 930–5.
13. Laforge R, Spector W, Sternberg J. The relationship of vision and hearing impairment to one-year mortality and functional decline. *J Aging Health* 1992; 4: 126–48.
14. Agrawal S, Mishra A, Jagade M *et al.* Effects of hypertension on hearing. *Indian J Otolaryngol Head Neck Surg* 2013; 65: 614–8.
15. Shaper AG, Pocock SJ, Walker M *et al.* British Regional Heart Study: cardiovascular risk factors in middle-aged men in 24 towns. *Br Med J (Clin Res Ed)* 1981; 283: 179–86.
16. Ranganathan B, Counter P, Johnson I. Validation of self-reported hearing loss using television volume. *J Laryngol Otol* 2011; 125: 18–21.
17. Katz S, Ford AB, Moskowitz RW *et al.* Studies of illness in the aged. The index of ADL: a standardized measure of biological and psychosocial function. *JAMA* 1963; 185: 914–9.
18. Lawton MP, Brody EM. Assessment of older people: self-maintaining and instrumental activities of daily living. *Gerontologist* 1969; 9: 179–86.
19. National Institutes of Health. Clinical guidelines on the identification, evaluation and treatment of overweight and obesity in adults - the evidence report. *Obes Res* 1998; 6: 51S–209S.
20. Wannamethee SG, Lowe GD, Whincup PH *et al.* Physical activity and hemostatic and inflammatory variables in elderly men. *Circulation* 2002; 105: 1785–90.
21. Harwood RH, Pound P, Ebrahim S. Determinants of social engagement in older men. *Psychol Health Med* 2000; 5: 75–85.
22. Avlund K, Lund R, Holstein BE *et al.* Social relations as determinant of onset of disability in aging. *Arch Gerontol Geriatr* 2004; 38: 85–99.
23. James BD, Boyle PA, Buchman AS *et al.* Relation of late-life social activity with incident disability among community-dwelling older adults. *J Gerontol A Biol Sci Med Sci* 2011; 66: 467–73.
24. Brennan M, Horowitz A, Su YP. Dual sensory loss and its impact on everyday competence. *Gerontologist* 2005; 45: 337–46.
25. Tun PA, McCoy S, Wingfield A. Aging, hearing acuity, and the attentional costs of effortful listening. *Psychol Aging* 2009; 24: 761–6.
26. Verschuur CA, Dowell A, Syddall HE *et al.* Markers of inflammatory status are associated with hearing threshold in older people: findings from the Hertfordshire Ageing Study. *Age Ageing* 2012; 41: 92–7.
27. Wannamethee SG, Whincup PH, Lennon L *et al.* Associations between fibrin D-dimer, markers of inflammation, incident self-reported mobility limitation, and all-cause mortality in older men. *J Am Geriatr Soc* 2014; 62: 2357–62.

Received 17 December 2015; accepted in revised form 23 March 2016