


Accepted Manuscript

Migration and allergic diseases in a rural area of a developing country

Alejandro Rodriguez, MSc, Maritza G. Vaca, MD, Martha E. Chico, MD, Laura C. Rodrigues, MD, PhD, Mauricio L. Barreto, MD, PhD, Philip J. Cooper, MD, PhD.



PII: S0091-6749(16)30091-4

DOI: [10.1016/j.jaci.2016.01.052](https://doi.org/10.1016/j.jaci.2016.01.052)

Reference: YMAI 12053

To appear in: *Journal of Allergy and Clinical Immunology*

Received Date: 9 April 2015

Revised Date: 25 December 2015

Accepted Date: 15 January 2016

Please cite this article as: Rodriguez A, Vaca MG, Chico ME, Rodrigues LC, Barreto ML, Cooper PJ, Migration and allergic diseases in a rural area of a developing country, *Journal of Allergy and Clinical Immunology* (2016), doi: 10.1016/j.jaci.2016.01.052.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1 Migration and allergic diseases in a rural area of a developing country

2 Alejandro Rodriguez, MSc,^{a,b,c} Maritza G Vaca, MD,^a Martha E Chico ,MD,^a Laura C
3 Rodrigues, MD ,PhD,^b Mauricio L Barreto,MD, PhD,^{d,e} Philip J Cooper,MD, PhD.^{a,c,f}

4

5 ^aLaboratorio de Investigación FEPIS, Quinindé, Esmeraldas Province, Ecuador

6 ^bFaculty of Epidemiology and Population Health, London School of Hygiene and Tropical
7 Medicine, London, UK

8 ^cCentro de Investigación en Enfermedades Infecciosas y Crónicas, Pontificia Universidad
9 Católica del Ecuador, Quito, Ecuador

10 ^dCentro de Pesquisas Gonçalo Muniz, FIOCRUZ, Salvador, Brazil

11 ^eInstituto de Saude Coletiva, Universidade Federal da Bahia, Salvador, Brazil

12 ^fInstitute of Infection and Immunity, St George's University of London, London, UK.

13

14 §Corresponding author

15 Alejandro Rodríguez, MSc.

16 Faculty of Epidemiology and Population Health, London School of Hygiene and Tropical
17 Medicine, Keppel street, London, UK.

18 Telf: 07517787355 Skype: alejo.rodrigues Email: alejandro.rodriguez@lshtm.ac.uk

19

20 The research was supported by grants from the Wellcome Trust (072405/Z/03/Z and
21 088862/Z/09/Z). The funders had no role in study design, data collection and analysis,
22 decision to publish, or preparation of the manuscript.

23

24

25

26

27

28 **Capsule summary**

29 Migration processes as the absence of the mother at home through temporary or permanent
30 migration could be an important determinant of the increase of allergic diseases in rural
31 areas of developing regions.

32

33 **Key words:** allergic diseases, developing country, migration, rural area

34

35 **Word count: 975**

36

37

38 To the Editor:

39 Studies in Developing Countries (DCs) have frequently reported a lower prevalence of
40 allergic diseases (AllDis) in rural areas compared with urban settings, and this has been
41 attributed to the protective effects of environmental exposures such as rural lifestyle.[1]
42 Recent evidence from studies conducted in Africa and Asia showed that AllDis are
43 increasing in urban and even in rural settings, reducing the urban-rural prevalence gap.[2,3]
44 It has been hypothesized that temporal increases in AllDis prevalence might be associated
45 with urbanization processes, especially with the change from rural to more modern urban
46 lifestyles.[1]

47 Migration is an important component of the urbanization process and involves
48 socioeconomic, environmental and lifestyle changes in rural and urban populations.
49 However, the effects of migration on AllDis in urban and rural settings of DCs have not been
50 explored.[4] The impact of migration on AllDis has been largely investigated by comparing
51 populations that have migrated from DCs (presumed low-risk for AllDis) to developed
52 countries (presumed high-risk).[5] These studies have shown that being born in a country of
53 low risk provides protection against asthma, but this protection may decline with the length of
54 residence in the new environment.[5] Others studies have shown that age of migration and
55 time since migration are associated with the risk of asthma and other AllDis, often leading to
56 a higher risk of atopy and allergy among migrants than the local population.[6]

57 The SCAALA (Social Changes, Asthma and Allergy in Latin America) study has been
58 investigating the effects of migration on the prevalence of AllDis in schoolchildren living in
59 rural and urban areas.[4] We studied 4295 rural and 2510 urban children aged 5-16 years
60 attending a convenience sample of schools in Esmeraldas province, Ecuador. Data on
61 potential risk factors, migration (direction and distance of migration, age at migration, and
62 time since migration), and wheeze, rhinitis, eczema symptoms within the previous 12 months
63 were collected using an investigator-administered questionnaire that included the core

64 allergy questions of ISAAC phase II.[4] Atopy was measured by skin prick testing to 7
65 aeroallergens.

66 Results from the rural area showed that children who migrated during the first year of life had
67 a greater risk of wheeze and rhinitis compared to non-migrant children, and children with
68 history of international migration (children from rural areas of Colombia) had a higher
69 prevalence of rhinitis than non-migrant children (Table 1). The study also evaluated the
70 effects of maternal migration on allergic outcomes in children using the variables, maternal
71 history of migration and children living with one or no parent. These analyses suggested that
72 children whose mothers had a history of migration had a greater risk of eczema than children
73 whose mother did not and children who did not live with any parent had more wheeze than
74 children living with both parents (Table 1). The magnitude of the latter association was
75 greater for all allergic symptoms among children of migrant mothers (Table 2). No
76 associations were observed for atopy (at least one positive allergen skin test).

77 The present study is unique in investigating migrants within a rural area of a DC, where
78 migrants come from urban and rural settings. In this setting, age at migration and
79 international migration were important factors associated with a higher risk of AllDis in rural
80 populations. A novel observation was the effect on the prevalence of AllDis of migrant status
81 of the mother: children of migrant mothers not living with either parent had a two-fold greater
82 risk of all 3 AllDis compared to children living with both parents. These data raise a question:
83 *Could it be that social effects of migration, such as absence of parents at home, are*
84 *important determinants of the increase in AllDis in rural populations of DCs?* In order to
85 answer this question, we need to consider some demographic patterns in these regions. It is
86 well known that people in rural villages move to urban areas, temporally or permanently, in
87 search of work to improve their quality of life. A high proportion of these rural migrants are
88 single women who provide economic support for their families. Most of these women leave
89 their children in the community of origin to be cared for by relatives. Some of these
90 immigrants are able to settle in the city while others return to their rural communities.[7] In

91 the SCAALA rural population 31% of the children and 23% of the mothers had history of
92 migration, and 15% of the children lived with no parent.

93 If the absence of parents at home (especially the mother) is an important determinant of the
94 increase of AllDis in DCs, then two migration trends that have occurred over recent decades
95 might help us understand temporal trends in AllDis. In the past, most economic migrants
96 were young men, but now "*feminization of migration*" is a growing trend worldwide because
97 of a greater demand for female labour.[8] Second, "*circular migration*" is a common
98 phenomenon in regions that are undergoing high levels of urbanization, and it refers to
99 repeated migrations between rural and urban areas due to improvements in transport and
100 modern forms of communication.[9]

101 Migration affects not only the individual who migrates but also their family. Migration impacts
102 on roles, support structures, and responsibilities of family members resulting in changes in
103 social and psychological factors. In the case of maternal migration, children who remain in
104 their community may experience heightened levels of stress and depression due to
105 separation from their primary carer. Psychological mechanisms have been proposed to
106 explain how emotional factors, in the context of family, might affect the development of
107 allergic diseases.[10] For this reason, we propose that the absence of the parents at home,
108 through temporary or permanent migration, may contribute to the increase of AllDis in rural
109 and urban populations of DCs.

110 Finally, further analyses in different populations living in rural and urban areas evaluating the
111 effects on migration on AllDis are required. A better understanding of the social,
112 psychological and environmental effects of migration on AllDis in DCs is required.

113

114

115

116

117

118 **References:**

- 119 1. von Hertzen L, Haahtela T. Disconnection of man and the soil: reason for the asthma and
120 atopy epidemic? *J Allergy Clin Immunol.* 2006;117(2):334-344.
121 doi:10.1016/j.jaci.2005.11.013.
- 122 2. Addo-Yobo EOD, Woodcock A, Allotey A, Baffoe-Bonnie B, Strachan D, Custovic A.
123 Exercise-induced bronchospasm and atopy in Ghana: Two surveys ten years apart. *PLoS*
124 *Med.* 2007;4(2):0355-0360. doi:10.1371/journal.pmed.0040070.
- 125 3. Selcuk ZT, Demir AU, Tabakoglu E, Caglar T. Prevalence of asthma and allergic
126 diseases in primary school children in Edirne, Turkey, two surveys 10 years apart. *Pediatr*
127 *Allergy Immunol.* 2010;21(4 Pt 2):e711-e717. doi:10.1111/j.1399-3038.2010.01008.x.
- 128 4. Cooper PJ, Chico ME, Vaca MG, et al. Risk factors for asthma and allergy associated with
129 urban migration: background and methodology of a cross-sectional study in Afro-Ecuadorian
130 school children in Northeastern Ecuador (Esmeraldas-SCAALA Study). *BMC Pulm Med.*
131 2006;6:24. doi:10.1186/1471-2466-6-24.
- 132 5. Cabieses B, Uphoff E, Pinart M, Antó JM, Wright J. A Systematic Review on the
133 Development of Asthma and Allergic Diseases in Relation to International Immigration: The
134 Leading Role of the Environment Confirmed. *PLoS One.* 2014;9(8):e105347.
135 doi:10.1371/journal.pone.0105347.
- 136 6. Rottem M, Szyper-Kravitz M, Shoenfeld Y. Atopy and asthma in migrants. *Int Arch Allergy*
137 *Immunol.* 2005;136:198-204. doi:10.1159/000083894.
- 138
- 139 7. Mujeres migrantes de América Latina y el Caribe: derechos humanos, mitos y duras
140 realidades. Santiago de Chile: Comisión Económica para América Latina y el Caribe; 2005.
141 Available at: [http://www.cepal.org/es/publicaciones/7200-mujeres-migrantes-de-america-](http://www.cepal.org/es/publicaciones/7200-mujeres-migrantes-de-america-latina-y-el-caribe-derechos-humanos-mitos-y-duras)
142 [latina-y-el-caribe-derechos-humanos-mitos-y-duras.](http://www.cepal.org/es/publicaciones/7200-mujeres-migrantes-de-america-latina-y-el-caribe-derechos-humanos-mitos-y-duras) Accessed February 22, 2015
- 143
- 144 8. Chammartin G. The feminization of international migration. *Int Migr Program Int Labour*
145 *Organ.* 2002:37-40. <http://library.fes.de/pdf-files/gurn/00072.pdf>.
- 146
- 147 9. Beguy D, Bocquier P, Zulu EM. Circular migration patterns and determinants in Nairobi
148 slum settlements. *Demogr Res.* 2010;23:549-586. doi:10.4054/DemRes.2010.23.20.
- 149 10. Kaugars AS, Klinnert MD, Bender BG. Family influences on pediatric asthma. *J Pediatr*
150 *Psychol.* 2004;29:475-491. doi:10.1093/jpepsy/jsh051

151

152

153

154

155

156

157

158

159 **Table 1. Odds ratios (OR) and 95% confidence intervals (95% CI) for associations**
 160 **between migration variables and allergic symptoms adjusted for sex, age and**
 161 **socioeconomic status.**

Variables	Categories ^A	n	Wheeze	Rhinitis	Eczema
			OR (95% CI)	OR (95% CI)	OR (95% CI)
Direction of migration	No Migrant	2964	1	1	1
	Rural to Rural	555	1.13 (0.84-1.52)	1.02 (0.7-1.49)	1.23(0.82-1.83)
	Urban to Rural	776	0.97 (0.74-1.27)	1.18 (0.86-1.61)	1.16 (0.81-1.66)
Distance of migration	No Migrant	2964	1	1	1
	National	1263	0.99 (0.79-1.25)	1.04 (0.79-1.38)	1.21(0.90-1.64)
	International	68	1.71 (0.88-3.32)	2.39(1.16-4.92)*	0.64(0.16-2.66)
Age at migration (years)	No Migrant	2964	1	1	1
	<1	269	1.47(1.02-2.12)*	1.59(1.03-2.46)*	1.25 (0.73-2.14)
	1-5	560	0.96 (0.71-1.31)	1.18 (0.83-1.69)	1.17 (0.78-1.75)
	>5	502	0.88 (0.62-1.24)	0.76 (0.48-1.19)	1.16 (0.75-1.79)
Time since migration (years)	No Migrant	2964	1	1	1
	<3 vs NM	383	0.98 (0.68-1.4)	0.94 (0.6-1.49)	0.96 (0.57-1.61)
	3-5 vs NM	197	0.56 (0.31-1.02)	0.9 (0.48-1.69)	1.53 (0.86-2.7)
	>5 vs NM	751	1.21 (0.94-1.58)	1.26 (0.92-1.73)	1.21 (0.85-1.73)
Maternal history of Migration	No	3314	1	1	1
	Yes	981	1.22 (0.96-1.53)	1.24 (0.93-1.65)	1.88(1.39-2.53)*
Parents living in the child's house	Both	2490	1	1	1
	One	1146	1.07 (0.84-1.36)	1.16 (0.87-1.54)	1.21 (0.88-1.67)
	None	659	1.57 (1.2-2.05)*	1.29 (0.92-1.81)	1.27 (0.86-1.86)

162 Outcomes were defined as: recent wheeze—reported wheezing during the previous 12 months; recent eczema—having a
 163 reported itchy rash with a flexural distribution in the previous 12 months; and recent rhinitis—nasal stuffiness or sneezing
 164 without a cold accompanied by itchy eyes in the previous 12 months. * p value < 0.05

165

166

167

168

169

170

171 **Table 2. Odds ratios (OR) and 95% confidence intervals (95% CI) for associations**
 172 **between allergic symptoms and parents living in the child's home (live with parents)**
 173 **stratified by maternal history of migration. ORs adjusted for sex, age and**
 174 **socioeconomic status.**

175

		Maternal history of migration					
		NO			YES		
	Live with parents	OR	95% CI	p value	OR	95% CI	p value
Wheeze	One vs. both	1	0.76-1.34	0.976	1.2	0.77-1.87	0.429
	None vs. Both	1.44	1.06-1.95	0.02	2.17	1.25-3.77	0.006
Rhinitis	One vs. both	1.03	0.73-1.46	0.858	1.46	0.85-2.52	0.171
	None vs. Both	1.1	0.74-1.64	0.627	2.07	1.05-4.08	0.036
Eczema	One vs. both	0.96	0.63-1.46	0.857	1.63	0.95-2.77	0.074
	None vs. Both	1.03	0.64-1.65	0.916	2.12	1.07-4.17	0.031

176

177

178

179

180

181

182

183

184

185

186