

Corresponding numerical data for Figure 1.

A)

Birthweight interval (grams)	Number of infants	
	Infants who had enterococcal infections	Infants who had non-enterococcal infections only
300-400	1	3
400-500	2	59
500-600	36	181
600-700	58	329
700-800	66	283
800-900	34	240
900-1000	23	172
1000-1100	23	138
1100-1200	5	101
1200-1300	10	100
1300-1400	6	72
1400-1500	9	52
1500-1600	4	43
1600-1700	11	46
1700-1800	2	37
1800-1900	6	33
1900-2000	4	25
2000-2100	7	26
2100-2200	2	38
2200-2300	3	31
2300-2400	3	22
2400-2500	4	30
2500-2600	6	28
2600-2700	4	33
2700-2800	0	30
2800-2900	0	24
2900-3000	2	22
3000-3100	2	24
3100-3200	3	32
3200-3300	1	28
3300-3400	2	36
3400-3500	2	21

3500-3600	3	31
3600-3700	0	20
3700-3800	4	13
3800-3900	2	12
3900-4000	2	11
4000-4100	0	7
4100-4200	0	3
4200-4300	0	6
4300-4400	2	2
4400-4500	0	4
4500-4600	0	1
4600-4700	0	2
4700-4800	0	0
4800-4900	0	3
4900-5000	0	1
5000-5100	0	3
5100-5200	0	0
5200-5300	0	1
5300-5400	0	1

B)

Gestation interval (weeks)	Number of infants	
	Infants who had enterococcal infections	Infants who had non-enterococcal infections only
20-21	0	0
21-22	0	0
22-23	1	2
23-24	34	148
24-25	69	326
25-26	45	278
26-27	36	265
27-28	30	203
28-29	17	204
29-30	19	131
30-31	12	106
31-32	9	107
32-33	8	80
33-34	12	63

34-35	13	59
35-36	6	55
36-37	9	63
37-38	12	86
38-39	8	92
39-40	4	69
40-41	12	84
41-42	0	47