

Letter

Advice on Avoiding the Valley of Death: Insights from a 3Rs Model of Aversive and Emetic Compound Identification

Robin S.B. Williams^{1*} and Paul L.R. Andrews²

¹Centre for Biomedical Sciences, School of Biological Sciences, Royal Hollow University of London, Egham, TW20 0EX, United Kingdom; ²Division of Biomedical Sciences, St George's University of London, Tooting, London, SW17 0RE, United Kingdom

The gap between the development of new 3Rs (replace, reduce, refine) technologies and their implementation into routine use has been called the “Valley of Death”¹. Reasons for reluctance in adoption of these approaches, particularly by industry, are often complex, and may be related to a lack of awareness of, or confidence in, 3Rs approaches. We report from the Valley of Death, on a long-term project developing the social amoeba *Dictyostelium discoideum* as an innovative 3Rs alternative to the use of animals for early identification of novel chemical entities (NCEs) with aversive and emetic properties, to provide guidance so others may learn from our experience.

The side-effects of nausea and vomiting are factors limiting drug development or may reduce patient compliance for established treatments (Holmes et al., 2009); identification of emetic effects of candidate drugs utilises *in vivo* animal studies. A bitter or pungent taste of a medication can also reduce patient compliance particularly in paediatrics (Mennella et al., 2013) and bitter tastants can be nauseogenic (Peyrot des Gachons et al., 2011). The taste profile of candidate drugs is assessed by the rodent Brief Aversion Taste Assay (BATA) (e.g. Soto et al., 2015). In these experiments, around 12 animals are housed individually with some confinement, and with water deprivation, where a specially designed apparatus allows the number of licks taken from bottles containing test substance to be measured (Clapham et al., 2012). Lick number provides an estimation of aversive or bitter taste (Soto et al., 2015). The BATA test falls within the European Union legislation 2010/63/EU (EU, 2010) regulating the use of animals in research (e.g. Soto et al., 2015) as it exceeds the threshold for a regulated procedure. Although this method provides the industry standard, it is time consuming and requires animal experimentation as does testing for emetic liability.

Microorganisms were proposed as an early non-animal methodology (NAM) in a tiered approach to identification of emetic liability and aversive properties of NCEs (Holmes et al., 2009). The social amoeba *Dictyostelium discoideum* (Fig. 1) was considered particularly attractive due to its use at the time in pharmacogenetic studies (Waheed et al., 2014; Chang et al., 2012; Kelly et al., 2018) and its recognition by the US National Institute of Health as a biomedical model system, and its use as an innovative model in a range of biomedical-related studies (Muller-Taubenberger et al., 2013).




Fig. 1: The social amoeba *Dictyostelium discoideum*

(A) *Dictyostelium* is found in leaf litter of temperate forests, with a single cells life involving the consumption of microorganisms and division by binary fission. Cells are around 10 μm long, and have a structure typical of eukaryotes (with a nucleus, endoplasmic reticulum, and mitochondria) and are haploid. Cells can be grown in the laboratory in nutrient-rich media or in the presence of bacteria as a food source. (B) Following the initiation of starvation, *Dictyostelium* cells progress through a series of stages leading to differentiation and multicellularity leading to the formation of a mature fruiting body, around 1mm tall, consisting of a spore head held aloft by a stalk. Spores within this structure are dormant, and resistant to dehydration, and upon release germinate to form single-celled amoeba. (C) This development process can be easily reproduced in laboratory conditions, and can be employed as a useful model for development and pharmacogenetic studies.

Received October 19, 2018; Accepted March 18, 2019;
Epub April 2, 2019; © The Authors, 2019.

ALTEX 36(X), ###-###. doi:10.14573/altex.1810182




Correspondence: Robin S. B. Williams, PhD, Centre for Biomedical Sciences,
School of Biological Sciences, Royal Hollow University of London,
Egham, TW20 0EX, United Kingdom
(robin.williams@rhul.ac.uk)

This is an Open Access article distributed under the terms of the Creative Commons Attribution 4.0 International license (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution and reproduction in any medium, provided the original work is appropriately cited.

¹ <https://www.nc3rs.org.uk/news/bridging-3rs-valley-death-nc3rs-strategy-2017-2019>

In a series of papers, we reported the utility of *Dictyostelium* as a model for bitter and emetic substance screening. These studies established the utility of *Dictyostelium* to detect emetic agents and bitter tastants (Robery et al., 2011), including proposing novel molecular targets (Robery et al., 2013). They demonstrated that the model responds (cell movement and shape change) in a concentration-dependent manner to bitter tastants occupying a diverse chemical space, and showed a correlation of response data from eleven bitter tastants between *Dictyostelium* and the rat BATA assay which itself correlated with human taste panel data (Cocorocchio et al., 2015; Otto et al., 2016). Overall, these results support the use of *Dictyostelium* to estimate the probability that an NCE will have a bitter, aversive taste and hence its continued consideration as a potential 3Rs model in the drug discovery and development pipeline. Comparisons between the human taste panel, rodent BATA and *Dictyostelium* for the detection of potentially aversive (particularly bitter tasting) substances shows some of the key features and advantages of each system (Tab. 1). It is unlikely that one system will provide comprehensive data regarding a NAM, but selective combinations of multiple systems, including *Dictyostelium*, may provide a fast, cheap, and animal reduction approach.

Tab. 1: Comparison between bitter tastant model systems

Parameter	 Human taste panel	 Rat BATA	 <i>Dictyostelium</i> cell behaviour
Requires ethical approval/regulated in EU	Yes	Yes	No
Prior information on toxicity <i>in vivo</i> needed	Yes	Yes	No
Throughput capacity	Relatively low	Relatively low	Relatively high
Training required for subjects	Yes	Yes	No
Read out	Perceived sensation & intensity	Licks/unit time	Cell behaviour (chemotaxis/cell shape)
Automated data collection and analysis possible	Yes	Yes	Yes
Detects concentration related effects	Yes	Yes	Yes
ED ₅₀ measurement	Yes	Yes	Yes
Time taken per compound to identify ED ₅₀	Hours	Days	Hours
Face validity	Excellent	Yes, but readout is the response to the taste rather than the sensation	Limited
Construct validity	Yes	Yes	Yes
Predictive validity	Total	Yes, based on relatively limited data	Yes, dependent upon additional validation
Experimental genetic manipulation possible	No	Yes	Yes
Mechanistic studies feasible	Potentially, but complex	Yes, but complex	Yes, relatively straightforward

Despite the progress made over the last 10 years, much of it in collaboration with the potential end-user (i.e., industry), we have been unable to obtain further support to finally validate (or not) the *Dictyostelium* assay as a reduction/replacement technology for the BATA assay and exit the Valley of Death. The basis for this lack of support is likely to be multifactorial. Firstly, a single celled organism, without a mouth, gut or nervous system and lacking a range of proteins that may provide a mechanistic target for aversive effects clearly has limited face validity. Secondly, in this area industry seems fundamentally conservative, with a strong focus on maintaining consistent testing using established models. Finally, funding environments remain highly selective, with grant applications often relying upon guidance from experts working in currently established technologies without a drive to supplant these technologies with new approaches, and the smallest criticism can lead to the rejection of a funding application. In an industry setting, significant financial investment in novel technologies without a strategic decision to move into that technology is likely to block the development of new initiatives. Industry commitment to novel 3Rs technologies is sometimes difficult to identify. It is particularly interesting to note that one shortcoming of approaches for the development of alternative methods is “*little input from end users*”, and that

methods were produced that “*did not adequately meet the testing requirements of end users*”². In our study, although we had input from a major industry partner at an early stage in the project to address these points, this did not ensure continued research support to fully assess the potential for industry adoption.

We believe that the barriers we have encountered are not unique and hence the lessons we have learned may have wider applicability; we make some recommendations for researchers looking to develop other NAMs.

1. Develop a relationship with industry partners. Key to implementing a new technology is to address all concerns of the end user, in this case our industrial partner. Thus, listening to their concerns, and addressing these issues is essential for them to ultimately engage with the technology. Relationships with industry can also give access to unpublished animal or human data enabling comparison with NAM data² to facilitate validation. One difficulty with engaging large multinational partners is that it may be hard to find the right person to contact regarding 3Rs-orientated research, and ensuring that applications for funding reach both the 3Rs coordinator and the decision maker in the section where the technology will be relevant. In our case, trying to develop a novel technology to improve identification of a potential side effect was not a standard type of novel drug-target or mechanism-related approach regarding a particular therapeutic area. Furthermore, we had developed good working relationships with various industry colleagues, who understood and supported the new technology, but perhaps we should also have focused on developing relationships with those who decide on future research initiatives and subsequent investment.

2. Publish, publish, and publish. Validation of new technologies is considerably strengthened through the peer review process in publishing papers. Publications provide clear evidence of innovation or discovery that has been reviewed independently, increasing trust in the technology. This approach also enhances outreach and supports further funding applications. The journals to publish in may also be worth considering – where impact factor may not be as important as access to the target audience, and should this audience be the relevant industry-focused group, or those with specific interest in 3Rs-technology or a broad readership from all areas of science and society?

3. Network. Develop impact through targeted industry and academic networks. Presenting talks at industry, academic or government meetings is likely to both improve the potential for engaging industry partners and will additionally provide feedback on concerns that still need to be addressed. We gave at least 12 presentations on our *Dictyostelium* research model including several to predominantly industrial audiences, with subsequent discussions related to project and funding.

4. Don't give up but recognise the limitations. With the highly competitive state of funding, all potential avenues of support must be investigated to maximise chances of continued investment, so keep looking for alternative mechanisms of support. However, projects such as this, where funding is required at the final step to validate a method for reduction/ replacement, are particularly problematic as the final data set (a graph of ID₅₀ values for a range of substances in *Dictyostelium* vs. rodent, and human data) may look like the pilot data just with more data points and a better correlation statistic. In addition, the number of animals that would be replaced if *Dictyostelium* exactly matched the predictability of the BATA assay would be relatively small, partially because of repeated use of individuals. We suspect that a few thousand animals are used globally in the BATA assay but the number used by industry is impossible to know. In competition with projects developing methods aimed at reducing/replacing a large number of animals in procedures with severity classified as moderate or even severe (particularly involving pain) it seems inevitable that a project aiming to reduce/replace fewer animals in a mild procedure will have a lower priority. Finally, our project focuses on detecting a potential drug side effect rather than investigating disease or drug mechanisms and this may also reduce the priority in competition with other projects.

Thus, following 10 years of research into developing a 3Rs model for screening NCE for emetic and aversive effects, we have arrived in the Valley of Death from which few 3Rs projects seem to emerge. Through highlighting key points in advancing the development of new 3Rs technologies that we have recognised through this time, we hope to help others champion their new technologies into industrial settings.

References

- Chang, P., Orabi, B., Deranieh, R. M. et al. (2012). The antiepileptic drug valproic acid and other medium-chain fatty acids acutely reduce phosphoinositide levels independently of inositol in dictyostelium. *Dis. Model. Mech* 5, 115-124. doi:10.1242/dmm.008029
- Clapham, D., Kirsanov, D., Legin, A. et al. (2012). Assessing taste without using humans: Rat brief access aversion model and electronic tongue. *Int J Pharm* 435, 137-139. doi:10.1016/j.ijpharm.2012.05.056
- Cocorocchio, M., Ives, R., Clapham, D. et al. (2016). Bitter tastant responses in the amoeba dictyostelium correlate with rat and human taste assays. *ALTEX* 33, 225-236. doi:10.14573/altex.1509011
- EU - European Union (2010). Directive 2010/63/EU of the European Parliament and of the Council of 22 September 2010 on the protection of animals used for scientific purposes. <https://bit.ly/2TdfGiT>
- Holmes, A. M., Rudd, J. A., Tattersall, F. D. et al. (2009). Opportunities for the replacement of animals in the study of nausea and vomiting. *Br J Pharmacol* 157, 865-880. doi:10.1111/j.1476-5381.2009.00176.x
- Kelly, E., Sharma, D., Wilkinson, C. J. et al. (2018). Diacylglycerol kinase (dgka) regulates the effect of the epilepsy and bipolar disorder treatment valproic acid in dictyostelium discoideum. *Dis Model Mech* 11, doi:10.1242/dmm.035600
- Mennella, J. A., Spector, A. C., Reed, D. R. et al. (2013). The bad taste of medicines: Overview of basic research on bitter taste. *Clin Ther* 35, 1225-1246. doi:10.1016/j.clinthera.2013.06.007

² ICCVAM (2018). A strategic roadmap for establishing new approaches to evaluate the safety of chemicals and medical products in the United States. DOI:10.22427/NTP-ICCVAM-ROADMAP2018

- Muller-Taubenberger, A., Kortholt, A. and Eichinger, L. (2013). Simple system--substantial share: The use of dictyostelium in cell biology and molecular medicine. *Eur J Cell Biol* 92, 45-53. doi:10.1016/j.ejcb.2012.10.003
- Otto, G. P., Cocorocchio, M., Munoz, L. et al. (2016). Employing dictyostelium as an advantageous 3rs model for pharmacogenetic research. *Methods Mol Biol* 1407, 123-130. doi:10.1007/978-1-4939-3480-5_9
- Peyrot des Gachons, C., Beauchamp, G. K., Stern, R. M. et al. (2011). Bitter taste induces nausea. *Curr Biol* 21, R247-248. doi:10.1016/j.cub.2011.02.028
- Robery, S., Mukanowa, J., Percie du, S. N. et al. (2011). Investigating the effect of emetic compounds on chemotaxis in dictyostelium identifies a non-sentient model for bitter and hot tastant research. *PLoS. One* 6, e24439. doi:10.1371/journal.pone.0024439
- Robery, S., Tyson, R., Dinh, C. et al. (2013). A novel human receptor involved in bitter tastant detection identified using dictyostelium discoideum. *J. Cell Sci* 126, 5465-5476. doi:10.1242/jcs.136440
- Soto, J., Sheng, Y., Standing, J. F. et al. (2015). Development of a model for robust and exploratory analysis of the rodent brief-access taste aversion data. *Eur. J. Pharm. Biopharm* 91, 47-51. doi:10.1016/j.ejpb.2015.01.016
- Waheed, A., Ludtmann, M. H., Pakes, N. et al. (2014). Naringenin inhibits the growth of dictyostelium and mdck-derived cysts in a trpp2 (polycystin-2)-dependent manner. *Br J Pharmacol* 171, 2659-2670. doi:10.1111/bph.12443

Conflict of Interest

The authors have no conflicts of interest.

Acknowledgements

We wish to acknowledge Janina Mukanowa, Steve Robery and Marco Cocorocchio. We also thank Robert Ives and Dave Clapham at Glaxo SmithKline for stimulating discussions about adapting a new model. Finally, we thank the Universities Federation for Animal Welfare, SouthWest London Academic Network, GlaxoSmithKline and the United Kingdom National Centre for Replacement, Refinement & Reduction of Animals in Research (Grant NC/S01201/1, NC/S01301/1) for funding support.