

PREDICTIVE ACCURACY OF THE CEREBROPLACENTAL RATIO FOR ADVERSE PERINATAL AND NEURODEVELOPMENTAL OUTCOMES IN SUSPECTED FETAL GROWTH RESTRICTION: A SYSTEMATIC REVIEW AND META-ANALYSIS

Agustin Conde-Agudelo¹

José Villar^{2,3}

Stephen H. Kennedy^{2,3}

Aris T. Papageorghiou^{2,3}

Authors' affiliations

¹ Perinatology Research Branch, *Eunice Kennedy Shriver* National Institute of Child Health and Human Development/National Institutes of Health/Department of Health and Human Services, Bethesda, MD and Detroit, MI, USA

² Nuffield Department of Obstetrics & Gynaecology, University of Oxford, Women's Centre, John Radcliffe Hospital, Oxford OX3 9DU, UK

³ Oxford Maternal & Perinatal Health Institute, Green Templeton College, Oxford OX2 6HG, UK

Short title: The cerebroplacental ratio predicts perinatal death in pregnancies with suspected fetal growth restriction

Corresponding author

Prof. José Villar, Nuffield Department of Obstetrics & Gynaecology, University of Oxford, Women's Centre, John Radcliffe Hospital, Oxford OX3 9DU, UK

e-mail: jose.villar@obs-gyn.ox.ac.uk

Tel. + 44 1865 740434

Financial support: Departmental funds

Disclosure: All authors report no conflicts of interest

Reprints will not be available

Word count: 3960

Key words: fetal growth restriction; cerebroplacental ratio; Doppler; predictive accuracy; perinatal outcomes; neurodevelopmental outcomes

ABSTRACT

Objective The cerebroplacental ratio (CPR) has been proposed for the routine surveillance of pregnancies with suspected fetal growth restriction (FGR), but the predictive performance of this test is unclear. The aim of this study was to determine the accuracy of the CPR for predicting adverse perinatal and neurodevelopmental outcomes in suspected FGR.

Methods PubMed, EMBASE, CINAHL, and Lilacs (all from inception to July 31, 2017) were searched for cohort or cross-sectional studies that reported on the accuracy of the CPR for predicting adverse perinatal and/or neurodevelopmental outcomes in singleton pregnancies with antenatally suspected FGR based on sonographic parameters. Summary receiver operating characteristic (ROC) curves, pooled sensitivities and specificities, and summary likelihood ratios (LRs) were generated.

Results Twenty-two studies (4301 women) met the inclusion criteria. Summary ROC curves showed that the best predictive accuracy of the CPR was for perinatal death and the worst was for neonatal acidosis, with areas under the summary ROC curves of 0.83 and 0.57, respectively. The predictive accuracy of the CPR was moderate-to-high for perinatal death (pooled sensitivity and specificity of 93% and 76%, respectively, and summary positive and negative LRs of 3.9 and 0.09, respectively), and low for composite of adverse perinatal outcomes, cesarean section for non-reassuring fetal status, Apgar <7 at 5 minutes, admission to the neonatal intensive care unit, neonatal acidosis, and neonatal morbidities with summary positive and negative LRs ranging from 1.1-2.5, and 0.3-0.9, respectively. An abnormal CPR result had moderate accuracy for predicting small for gestational age at birth (summary positive LR of 7.4). The

CPR had a higher predictive accuracy in pregnancies with suspected early-onset FGR. No study provided data for assessing the predictive accuracy of the CPR for adverse neurodevelopmental outcomes.

Conclusion The CPR appears to be useful in predicting perinatal death in pregnancies with suspected FGR. Nevertheless, before incorporating the CPR into the routine clinical management of suspected FGR, randomized controlled trials should assess whether the use of the CPR reduces perinatal death or other adverse perinatal outcomes.

INTRODUCTION

Fetal growth restriction (FGR) is a major clinical and public health challenge around the world^{1,2}. Small for gestational age (SGA) at birth, based on different cut-off values, is a commonly used proxy measure of FGR³. FGR is associated with an increased risk of short- and long-term morbidity and mortality, as well as impaired neurological and cognitive development⁴⁻¹¹.

Suspected FGR is defined in the antenatal period by sonographic estimation of fetal anthropometric measures using a wide range of seldom validated definitions and cut-off values¹²⁻¹⁶. The clinical management of suspected FGR is challenging and no consensus exists for the best way to monitor fetal well-being in these pregnancies; consequently, clinical practice varies considerably around the world¹⁷⁻¹⁹. The use of umbilical artery (UA) Doppler velocimetry in high-risk pregnancies, including those with suspected FGR, has been shown to be associated with a significant reduction in perinatal mortality and fewer cesarean deliveries and inductions of labour²⁰.

In 1987, Arbeille *et al.*²¹ reported that the cerebroplacental ratio (CPR), a measure of cerebral centralization of fetal blood flow, appeared to be superior to either the middle cerebral artery (MCA) or UA Doppler indices alone in predicting SGA among women with gestational hypertension. The CPR is calculated by dividing the Doppler indices (pulsatility index [PI], resistance index [RI], or systolic/diastolic ratio [S/D]) of the MCA by the UA. Physiologically, the CPR represents the interaction of alterations in blood flow to the brain as manifest by increased diastolic flow as a result of cerebrovascular dilatation due to hypoxia and increased placental resistance, leading to decreased diastolic flow in the UA²². Integrating the CPR into the clinical management of suspected FGR has

recently been proposed²²⁻²⁷, but the test's ability to predict adverse perinatal outcomes in this entity has been questioned^{28,29}. Hence, we carried out a systematic review and meta-analysis to assess the accuracy of the CPR to predict adverse perinatal and neurodevelopmental outcomes in antenatally suspected FGR.

METHODS

The systematic review was conducted following a prospectively prepared protocol and reported in accordance with recommended methods for systematic reviews of diagnostic test accuracy^{30,31}. The protocol was registered with PROSPERO on March 2016 (CRD42016036488; available from: http://www.crd.york.ac.uk/PROSPERO/display_record.php?ID=CRD42016036488).

Literature search

We searched PubMed, EMBASE, CINAHL, and Lilacs (all from inception to July 31, 2017) using a combination of keywords and text words related to *cerebroplacental ratio* and *fetal growth restriction* without language restrictions (Appendix S1).

Eligibility criteria

We included cohort or cross-sectional studies that reported on the accuracy of the CPR for predicting adverse perinatal and/or neurodevelopmental outcomes in singleton pregnancies with antenatally suspected FGR based on sonographic parameters, and provided the necessary information to generate 2x2 tables. Studies were excluded if they: (1) assessed retrospectively the predictive accuracy of the CPR in infants categorized as SGA or FGR based on postnatal parameters such as birthweight or other anthropometric measures, and/or

placental histopathology; (2) assessed the CPR in a mix of high-risk pregnancies but did not report results separately for pregnancies with suspected FGR; (3) assessed the CPR in the general population as a screening tool; (4) were case-control studies without complete information for cases with suspected FGR, case series or reports, editorials, comments, reviews, or letters without original data; (5) reported data for the CPR only as mean or median values; (6) did not publish accuracy test estimates or sufficient information to calculate them could not be retrieved.

One reviewer (A.C.-A.) screened titles and abstracts of all identified citations and selected potentially eligible studies. Then, these studies were retrieved and assessed by the same reviewer for inclusion and data extraction, and a 10% sample of the papers was examined by a second independent reviewer (J.V.). Disagreements were resolved through consensus. In cases of duplicate publication, we included only the most recent or complete version.

Reference standard outcomes

The reference standard outcomes included the following: perinatal death; any composite of adverse perinatal outcomes (as defined in the original study and regardless of its individual components); cesarean delivery for fetal distress/non-reassuring fetal status; Apgar <7 at 5 minutes; admission to the neonatal intensive care unit (NICU); neonatal acidosis; neonatal brain lesions; neonatal morbidities other than brain lesions; use of mechanical ventilation; SGA at birth (birthweight <10th, <5th or <3rd perpercentile or <2 standard deviations of mean adjusted for gestational age and based on local population values), and adverse neurodevelopmental outcomes (suspected or diagnosed developmental delay, cerebral palsy, intellectual disabilities, vision impairment, hearing loss, cognitive

and behavioral impairments, and motor, communication and learning disorders at any age in childhood).

Assessment of risk of bias

The risk of bias in each included study was evaluated by at least one investigator using a modified version of the QUADAS (Quality Assessment of Diagnostic Accuracy Studies)-2 tool³². The following domains were assessed: study design, description of the test, selection of test cut-off value, blinding of clinicians to the CPR results, inclusion of participants recruited into the study in the analysis, and use of interventions aimed to prevent adverse perinatal outcomes based on the CPR results. Each domain was scored as “low risk”, “high risk”, or “unclear risk” of bias (Appendix S2). We did not calculate a summary score estimating the overall quality of each study because of the well-known problems associated with such scores³³.

Data extraction

Data were extracted from each article using a specially designed form for capturing information on study characteristics (authors, setting, year of publication, method of recruiting women, design, prospective or retrospective data collection, blinding of test results, flow diagram, completeness of follow-up and reporting of withdrawals, and use of interventions after performing the test), patient characteristics (inclusion and exclusion criteria, sample size, and demographic characteristics), how the test was carried out (gestational age at testing, frequency of test, method of performance of test, type of Doppler and route, site of measurement, plane in which images were obtained, Doppler index used [PI, RI, or S/D], cut-off value used, and interval from Doppler examination to delivery), and reference standard outcomes assessed and their prevalences.

For each study and for all cut-off values defining an abnormal CPR result, we extracted the number of true positive, false positive, true negative, and false negative test results. When predictive accuracy data were not available, we recalculated them from the reported results including scatterplot and bar graphs. The corresponding authors of primary studies were contacted to obtain additional information on methods used and/or unpublished relevant data. Only three authors supplied additional data.

Data synthesis

Data extracted from each study were used to construct 2x2 contingency tables. When any single cell in these tables contained a zero, we added 0.5 to each cell to enable calculation of predictive values³⁴. Sensitivity and specificity with 95% confidence intervals (CIs) were calculated separately for all Doppler indices and cut-off values used, and reference standard outcomes reported. Then, we constructed summary receiver-operating characteristic (ROC) curves for each predefined reference standard outcome using the hierarchical summary ROC model, regardless of Doppler indices and cut-off values used to define abnormality³⁵. Variation in cut-off values across studies is taken into account by using this model. Pooled estimates and 95% CIs of sensitivity and specificity were generated using random-effects bivariate meta-regression models³⁶. For studies that reported results for more than one Doppler index and/or cut-off value, we selected the most commonly used. We also calculated area under the summary ROC curves with their corresponding 95% CIs, which allowed for comparison of the predictive accuracy of the CPR for different outcomes³⁷.

Thereafter, summary likelihood ratios (LRs) with 95% CIs were calculated from the pooled sensitivities and specificities³⁸. A guide for the interpretation of

LRs suggests that LRs >10 for a positive test result and LRs <0.1 for a negative test result generate large changes from pretest to post-test probabilities of disease; LRs of 5 to 10 and 0.1 to 0.2 generate moderate changes in probability; LRs of 2 to 5 and 0.2 to 0.5 generate small (but sometimes important) changes in probability; and LRs of 1 to 2 and 0.5 to 1 generate minimal (and rarely important) changes in probability³⁹. Finally, we planned to calculate the post-test probabilities of the most important reference standard outcomes by combining summary LRs obtained from meta-analyses for positive and negative test results and a global prevalence (pretest probability) of these reference standard outcomes across the studies³⁹.

Prespecified subgroup analyses were carried out to assess the predictive accuracy of the CPR for any composite of adverse perinatal outcomes according to gestational age (GA) at diagnosis or delivery (early-onset [<32 or <34 weeks] and late-onset [≥ 32 or ≥ 34 weeks], as defined by the authors), definition of abnormal CPR result (MCA-PI/UA-PI ≤ 1.08 , MCA-PI/UA-PI <5 th percentile, and MCA-RI/UA-RI <1 or <1.05), interval from CPR to delivery (≤ 7 and >7 days), and definition of suspected FGR used (estimated fetal weight [EFW] <10 th percentile for GA and EFW <10 th percentile for GA and abnormal UA Doppler). In addition, a post-hoc subgroup analysis according to the use or non-use of the CPR results for managing pregnancies with suspected FGR was performed. We also assessed the effect of risk of bias of the included studies on the predictive accuracy of the CPR by performing a sensitivity analysis, including only studies with a low risk of bias in at least five of the six domains evaluated.

As it is common in diagnostic accuracy studies, we anticipated that there would be substantial between-study variation in reported pairs of sensitivity and

specificity values. As forest plots, which display both sensitivity and specificity, depict estimates with associated CIs it is possible to discern the presence of high levels of heterogeneity where there is little overlap in the CIs from different studies. In order to formally investigate potential sources of heterogeneity, we used subgroup analysis and meta-regression by including covariates defined *a priori* (Doppler indices and cut-off values used, definition of suspected FGR used, GA at diagnosis or delivery, interval from CPR to delivery, and study's risk of bias) in the bivariate model, which enabled us to assess the effect of various factors on the predictive accuracy of the CPR^{40,41}. If there were at least 10 studies included in a meta-analysis, we assessed publication and related biases by examining the symmetry of the funnel plots with the Deeks' test⁴². A value of $P < 0.1$ for the slope coefficient indicated significant asymmetry of the funnel plot.

We used SAS version 9.2 (SAS Institute Inc, Cary, NC) for the analyses and Review Manager 5.3.5 (The Nordic Cochrane Centre, Copenhagen, Denmark) to generate forest plots and summary ROC curves.

RESULTS

Selection, characteristics and quality of studies

Of 1191 citations initially identified, 22 studies⁴³⁻⁶⁴ including a total of 4301 women/fetuses met the inclusion criteria (Figure 1). Two studies were performed using the same cohort^{60,61}, one reporting results for all cases of suspected FGR⁶⁰ and the other for cases of suspected early-onset FGR⁶¹. We included results of this last study only in the subgroup analysis according to GA at diagnosis or delivery.

The main characteristics of the included studies are displayed in Table 1. All studies but two^{44,52} were performed in European or North American countries.

The sample size ranged from 29⁴⁸ to 881⁵⁶ (median, 159). The definitions of suspected FGR used in the studies were as follows: EFW <10th percentile for GA (11 studies)^{44,45,47,50,54,56-59,62,64}, EFW <10th percentile for GA and/or abdominal circumference (AC) <5th percentile (two studies)^{60,61}, EFW <10th percentile for GA and abnormal UA Doppler indices (two studies)^{53,63}, AC <5th percentile for GA and abnormal UA Doppler indices (two studies)^{46,49}, AC <10th percentile for GA (one study)⁴³, AC <10th percentile for GA on at least two consecutive measurements, two weeks apart (one study)⁵⁵, EFW <10th percentile for GA with growth rate slower than normal and abnormal UA Doppler indices (one study)⁴⁸, EFW or AC <10th percentile for GA and abnormal UA Doppler indices (one study)⁵², and EFW below the GA-adjusted mean value minus 2 SD (2.3rd percentile), or a fall of ≥10% weight deviation from the mean weight between two ultrasound examinations (one study)⁵¹. Ten studies reported results for suspected late-onset FGR^{47,50,51,54,56-59,62,63}, four for suspected early-onset FGR^{47,56,58,61}, and 13 for suspected FGR at all GAs^{43-49,52,53,55,56,58,60}.

The most common definitions of an abnormal CPR result were MCA-PI/UA-PI ≤1.08 (eight studies^{43,47,51-53,56,60,61}) and MCA-PI/UA-PI <5th percentile for GA (six studies^{47,50,54,56,58,62}). The mean or median interval between the CPR and delivery was <48 hours in five studies^{46,48,50,51,62}, <7 days in six studies^{49,53,56,57,59,64}, 7 to 14 days in three studies^{44,47,58}, >14 days in five studies^{45,54,55,60,61}, and unreported in three studies^{43,52,63}. Most studies (N=16) used the last CPR result before delivery in analyses^{44,46-55,57-59,62,64}. Sixteen studies reported that the CPR results were not used to manage the pregnancies^{43-45,47,48,50-53,56,58-63}, one reported that the CPR results were used to manage the pregnancies⁵⁴, and the remaining five studies did not report on this

issue^{46,49,55,57,64}. Eleven studies provided data on the predictive accuracy of the CPR for a composite of adverse perinatal outcomes^{43,44,47,53,54,56-60,62}, nine on admission to the NICU^{43-45,51,56,58-60,62}, seven on cesarean delivery for non-reassuring fetal status^{43,45,50,58-60,62}, six each on perinatal death^{45,52,55,56,58,60} and Apgar <7 at 5 minutes^{43,44,51,58,59,64}, five each on neonatal acidosis^{50,51,59,62,64} and neonatal brain lesions^{45,48,52,55,63}, four on neonatal morbidities other than brain lesions^{43,45,46,49}, two on SGA at birth^{43,59}, and one on use of mechanical ventilation⁵⁵. No study provided data on adverse neurodevelopmental outcomes.

The risk of bias in each included study is shown in Figure 2. Eight studies (36%) fulfilled ≥ 5 criteria, whereas the remaining 14 studies (64%) had ≥ 2 methodological flaws. The most common deficiencies were related to blinding of clinicians to the CPR results and inclusion of participants recruited into the study in the analyses.

Predictive accuracy for adverse perinatal outcomes

Summary ROC curves of the CPR for predicting adverse perinatal outcomes in pregnancies with suspected FGR are shown in Figure 3. The best predictive accuracy was for perinatal death and the worst was for neonatal acidosis, with areas under the summary ROC curves of 0.83 (95% CI, 0.74-0.92) and 0.57 (95% CI, 0.51-0.63), respectively. Similar summary ROC curves were obtained for the prediction of any composite of adverse perinatal outcomes, cesarean delivery for non-reassuring fetal status, and admission to the NICU (areas under the summary ROC curves between 0.71 and 0.74). The sensitivity and specificity of the CPR ratio to predict adverse perinatal outcomes in suspected fetal growth restriction in the individual studies are shown in Figure S1.

Table 2 presents the pooled estimates of the predictive accuracy of the CPR for adverse perinatal outcomes. Overall, the CPR showed a moderate-to-high predictive ability for perinatal death with pooled sensitivity and specificity of 93% and 76%, respectively, and summary positive and negative LR of 3.9 and 0.09, respectively (six studies, 1495 fetuses, 29 perinatal deaths). The CPR had a low predictive performance for any composite of adverse perinatal outcomes, cesarean delivery for non-reassuring fetal status, Apgar <7 at 5 minutes, admission to the NICU, neonatal acidosis, neonatal brain lesions, neonatal morbidities other than brain lesions, and use of mechanical ventilation with summary positive and negative LR that varied between 1.1 and 2.5, and 0.3 and 0.9, respectively. An abnormal CPR result had moderate accuracy for predicting SGA at birth (summary positive LR of 7.4; two studies, 554 fetuses). Based on all included studies, we estimated that fetuses with suspected growth restriction had a prevalence rate (pretest probability) of 25% for the composite of adverse perinatal outcomes, 2.0% for perinatal death, and 90% for SGA at birth (birthweight <10th percentile for GA). Then, based on estimated pretest probabilities and summary positive and negative LR, we calculated that an abnormal CPR result would increase the pretest probability of the composite of adverse perinatal outcomes, perinatal death, and SGA at birth from 25% to 45%, 2% to 7.4%, and 90% to 98.5%, respectively, whereas a normal CPR result would decrease the pretest probability to 17%, 0.2%, and 84%, respectively.

Visual assessment of both forest plots (Figure S1) and summary ROC curves (Figure 3) suggested substantial between-study heterogeneity, mainly for perinatal death, any composite of adverse perinatal outcomes, cesarean delivery for non-reassuring fetal status, and admission to the NICU. Meta-regression

analyses showed that none of the prespecified covariates explained the heterogeneity (Table S1). Sensitivity analyses revealed that pooled predictive accuracy estimates obtained from studies with low risk of bias in ≥ 5 domains did not differ significantly from those obtained in the overall analysis (data not shown). The funnel plot of the meta-analysis that included at least ten studies showed no significant asymmetry ($P = 0.19$, by Deeks' test).

Subgroup analyses

Subgroup analyses of the accuracy of the CPR to predict any composite of adverse perinatal outcomes are depicted in Table 3. The CPR had a significantly higher predictive accuracy for any composite of adverse perinatal outcomes among pregnancies with suspected early-onset FGR than among those with suspected late-onset FGR. Moreover, the accuracy of the CPR for predicting any composite of adverse perinatal outcomes was lower when using a MCA-PI/UA-PI < 5 th percentile as the abnormal result and when the CPR results were used to manage pregnancies. There were no differences in the predictive ability of the CPR between studies in which the interval from CPR to delivery was ≤ 7 days or > 7 days, and between studies using an EFW < 10 th percentile for GA as definition of suspected FGR and that using an EFW < 10 th percentile for GA and abnormal UA Doppler as definition of suspected FGR.

DISCUSSION

Main findings

The results of this systematic review indicate that the CPR had a moderate-to-high predictive accuracy for perinatal death, the most important outcome measure in relation with utero-placental insufficiency in suspected FGR. In particular, a normal CPR result had high accuracy to identify which fetuses with

suspected growth restriction are at low risk of dying in the perinatal period, decreasing the pretest probability of perinatal death from 2% to 0.2%. Overall, the CPR had a low predictive accuracy for the other adverse perinatal outcomes considered, several of which are less correlated with utero-placental insufficiency in suspected FGR. Notwithstanding, the presence of an abnormal CPR result increased the pretest probability of having an adverse perinatal outcome from 25% to 45%. In addition, subgroup analyses suggest that the predictive accuracy of the CPR is higher in pregnancies with suspected early-onset FGR and when a $MCA-PI/UA-PI \leq 1.08$ or a $MCA-RI/UA-RI < 1$ or < 1.05 is used as the definition of an abnormal CPR.

Previously, it has been suggested that the CPR is a stronger predictor of adverse perinatal outcomes in suspected late-onset FGR than suspected early-onset FGR^{23,25-27}. Unexpectedly, our subgroup analysis showed the opposite: a higher predictive accuracy for adverse perinatal outcomes in pregnancies with suspected early-onset FGR. Usually, suspected late-onset FGR is characterized by abnormal Doppler indices involving the MCA, with a normal or minimally elevated resistance of the UA²². In contrast, suspected early-onset FGR is characterized by abnormal Doppler indices of both the UA and MCA²². Abnormality in both vessels included in the calculation of the CPR, in particular high values of UA indices, could explain the better predictive accuracy of the CPR in suspected early-onset FGR in comparison with suspected late-onset FGR in which there are abnormal indices in only one vessel.

It was noteworthy that no included study provided data to assess the predictive ability of the CPR for adverse neurodevelopmental outcomes in pregnancies with suspected FGR. A secondary analysis of the TRUFFLE study⁶⁵

reported that the CPR was not associated with neurodevelopmental impairment at 2 years' corrected age in fetuses with suspected early-onset growth restriction⁶⁶. Two studies reported similar results for a decreased MCA-PI in suspected FGR^{67,68}. A systematic review reported that SGA or growth-restricted fetuses with cerebral redistribution may be at higher risk of adverse neurodevelopmental outcomes⁶⁹. However, none of the studies included in this review used the CPR for defining cerebral redistribution.

Strengths and limitations

The reliability and robustness of our systematic review are supported by the: (1) adherence to guidelines for the conduct and reporting of systematic reviews of predictive test accuracy; (2) use of a prospective protocol designed to address a highly specific research question; (3) comprehensive literature search without language restrictions; (4) inclusion of a relatively large number of studies, mostly published in recent years; (5) strict study quality assessment; (6) quantitative synthesis of the evidence; (7) use of contemporary statistical methods to obtain summary measures of predictive accuracy including subgroup and sensitivity analyses; and (8) exploration of potential sources of heterogeneity.

Limitations include the lack of blinding of the CPR results or omission of information on this subject in approximately two-thirds of the included studies. Although most studies reported that the CPR results were not used to manage pregnancies with suspected FGR, it is possible that women with abnormal CPR results were followed-up more closely or received interventions, which could have introduced bias in the assessment of the test's predictive accuracy. However, sensitivity analyses that were restricted to studies at low risk of

blinding bias showed no significant differences in the results obtained with overall meta-analyses.

There were considerable differences among studies in the definition of suspected FGR and Doppler indices/cut-off values used for defining an abnormal CPR, which limit the generalisability of our findings. Moreover, prespecified variables did not explain substantial heterogeneity and therefore, pooled estimates of predictive accuracy should be interpreted cautiously. Finally, the statistical power of some of our meta-analyses was limited by the small number of studies within each subgroup and the relatively small number of outcome events in some included studies.

Interpretation in the light of previous systematic reviews

We identified three systematic reviews on the predictive accuracy of the CPR for adverse perinatal outcomes⁷⁰⁻⁷². Nassr et al⁷⁰ included seven studies, and reported that an abnormal CPR in pregnancies at high risk for FGR or with diagnosis of FGR increased the risk for adverse perinatal outcomes. Summary ROC curves showed that the CPR had a better predictive accuracy for neonatal complications and NICU admission. Dunn et al⁷¹ reported that the CPR was predictive of cesarean section for intrapartum fetal compromise, SGA, and NICU admission in pregnancies at term. These reviews did not report pooled estimates of predictive accuracy. Finally, Vollgraaf Heidweiller-Schreurs *et al.*⁷² assessed the accuracy of the CPR to predict adverse perinatal outcomes in singleton pregnancies of all risk profiles. The CPR was significantly superior to UA and MCA Doppler in predicting a composite of adverse perinatal outcomes and emergency delivery for fetal distress. No differences were found between the CPR and either UA Doppler or MCA Doppler in the prediction of perinatal death,

low Apgar score, or NICU admission. Overall, our estimates of the predictive accuracy of the CPR for adverse perinatal outcomes among pregnancies with suspected FGR were lower than those reported in this review among pregnancies of all risk profiles.

The CPR has been hypothesized to be a more accurate test for predicting adverse perinatal outcomes than its individual components UA and MCA Doppler. When comparing the estimates obtained in our study with those reported in two meta-analyses that assessed the accuracy of UA Doppler⁷³ and MCA Doppler⁷⁴ to predict adverse perinatal outcomes in high-risk pregnancies, the CPR had better predictive accuracy for perinatal death (summary positive and negative LR of 3.9 and 0.09, respectively) than both UA Doppler (summary positive and negative LR of 2.5 and 0.3, respectively) and MCA Doppler (summary positive and negative LR of 1.4 and 0.5, respectively). In general, the predictive accuracy of the CPR for other adverse perinatal outcomes appeared to be comparable to those of UA Doppler and MCA Doppler.

Conclusions

The CPR appears to be useful in predicting perinatal death in pregnancies with suspected FGR. Nevertheless, before incorporating the CPR into the routine clinical management of suspected FGR, randomized controlled trials –ideally blinded- should assess whether the use of the CPR reduces perinatal death or other adverse perinatal outcomes. Studies are required to assess the predictive accuracy of the CPR for adverse neurodevelopmental outcomes in fetuses with antenatally suspected growth restriction.

ACKNOWLEDGEMENTS

Aris T. Papageorgiou is supported by the Oxford Partnership Comprehensive Biomedical Research Centre with funding from the Department of Health NIHR Biomedical Research Centres funding scheme. We are very grateful to Drs. Francesc Figueras, Inas Babic, and Carri R. Warshak for providing unpublished data from their studies and for clarifying other queries.

REFERENCES

1. de Onis M, Blössner M, Villar J. Levels and patterns of intrauterine growth retardation in developing countries. *Eur J Clin Nutr* 1998; **52**(Suppl 1): S5–15.
2. Lee AC, Katz J, Blencowe H, Cousens S, Kozuki N, Vogel JP, Adair L, Baqui AH, Bhutta ZA, Caulfield LE, Christian P, Clarke SE, Ezzati M, Fawzi W, Gonzalez R, Huybregts L, Kariuki S, Kolsteren P, Lusingu J, Marchant T, Meriardi M, Mongkolchat A, Mullany LC, Ndirangu J, Newell ML, Nien JK, Osrin D, Roberfroid D, Rosen HE, Sania A, Silveira MF, Tielsch J, Vaidya A, Willey BA, Lawn JE, Black RE; CHERG SGA-Preterm Birth Working Group.. National and regional estimates of term and preterm babies born small for gestational age in 138 low-income and middle-income countries in 2010. *Lancet Glob Health* 2013; **1**: e26-36.
3. WHO Expert Committee on Physical Status. Physical status: the use and interpretation of anthropometry - Report of a WHO expert committee. WHO Technical Report Series 854. Geneva: World Health Organization, 1995.
4. Katz J, Lee AC, Kozuki N, Lawn JE, Cousens S, Blencowe H, Ezzati M, Bhutta ZA, Marchant T, Willey BA, Adair L, Barros F, Baqui AH, Christian P, Fawzi W, Gonzalez R, Humphrey J, Huybregts L, Kolsteren P, Mongkolchat A, Mullany LC, Ndyomugenyi R, Nien JK, Osrin D, Roberfroid D, Sania A, Schmiegelow C, Silveira MF, Tielsch J, Vaidya A, Velaphi SC, Victora CG, Watson-Jones D, Black RE; CHERG Small-for-Gestational-Age-Preterm Birth Working Group. Mortality risk in preterm and small-for-gestational-age infants in low-income and middle-income countries: a pooled country analysis. *Lancet* 2013; **382**: 417-425.
5. Hirst JE, Villar J, Victora CG, Papageorghiou AT, Finkton D, Barros FC, Gravett MG, Giuliani F, Purwar M, Frederick IO, Pang R, Cheikh Ismail L, Lambert A, Stones W, Jaffer YA, Altman DG, Noble JA, Ohuma EO, Kennedy SH, Bhutta ZA; International Fetal and Newborn Growth Consortium for the 21st Century (INTERGROWTH-21st). The antepartum stillbirth syndrome: risk factors and pregnancy conditions identified from the INTERGROWTH-21st Project. *BJOG* 2016 Dec 28. doi: 10.1111/1471-0528.14463. [Epub ahead of print].
6. Longo S, Borghesi A, Tzialla C, Stronati M. IUGR and infections. *Early Hum Dev* 2014; **90**(Suppl 1): S42-4.
7. Chernauskas SD. Update: consequences of abnormal fetal growth. *J Clin Endocrinol Metab* 2012; **97**: 689-695.
8. Baschat AA. Neurodevelopment after fetal growth restriction. *Fetal Diagn Ther* 2014; **36**: 136-142.
9. Murray E, Fernandes M, Fazel M, Kennedy SH, Villar J, Stein A. Differential effect of intrauterine growth restriction on childhood neurodevelopment: a systematic review. *BJOG* 2015; **122**: 1062-1072.

10. Barker DJ. Adult consequences of fetal growth restriction. *Clin Obstet Gynecol* 2006; **49**: 270-283.
11. Salam RA, Das JK, Bhutta ZA. Impact of intrauterine growth restriction on long-term health. *Curr Opin Clin Nutr Metab Care* 2014; **17**: 249-254.
12. American College of Obstetricians and Gynecologists. ACOG Practice bulletin no. 134: fetal growth restriction. *Obstet Gynecol* 2013; **121**: 1122-1133.
13. Green-top Guideline No. 31. *The Investigation and Management of the Small-for-Gestational-Age Fetus*. London: Royal College of Obstetricians and Gynaecologists, 2013.
14. Society for Maternal-Fetal Medicine Publications Committee, Berkley E, Chauhan SP, Abuhamad A. Doppler assessment of the fetus with intrauterine growth restriction. *Am J Obstet Gynecol* 2012; **206**: 300-308.
15. Lausman A, Kingdom J; MATERNAL FETAL MEDICINE COMMITTEE. Intrauterine growth restriction: screening, diagnosis, and management. *J Obstet Gynaecol Can* 2013; **35**: 741-748.
16. Gordijn SJ, Beune IM, Thilaganathan B, Papageorghiou A, Baschat AA, Baker PN, Silver RM, Wynia K, Ganzevoort W. Consensus definition of fetal growth restriction: a Delphi procedure. *Ultrasound Obstet Gynecol* 2016; **48**: 333-339.
17. Chauhan SP, Dahlke JD, Magann EF, Chang E, Gupta L, Mol BW, Lewis DF. Isolated intrauterine growth restriction: a survey of Central Association of Obstetricians Gynecologists (CAOG) members. *J Matern Fetal Neonatal Med* 2013; **26**: 497-502.
18. Unterscheider J, Daly S, Geary MP, Kennelly MM, McAuliffe FM, O'Donoghue K, Hunter A, Morrison JJ, Burke G, Dicker P, Tully EC, Malone FD. Definition and management of fetal growth restriction: a survey of contemporary attitudes. *Eur J Obstet Gynecol Reprod Biol* 2014; **174**: 41-45.
19. Savchev S, Figueras F, Gratacos E. Survey on the current trends in managing intrauterine growth restriction. *Fetal Diagn Ther* 2014; **36**: 129-135.
20. Alfirevic Z, Stampalija T, Dowswell T. Fetal and umbilical Doppler ultrasound in high-risk pregnancies. *Cochrane Database Syst Rev* 2017; **6**: CD007529.
21. Arbeille P, Roncin A, Berson M, Patat F, Pourcelot L. Exploration of the fetal cerebral blood flow by duplex Doppler--linear array system in normal and pathological pregnancies. *Ultrasound Med Biol* 1987; **13**: 329-337.

22. DeVore GR. The importance of the cerebroplacental ratio in the evaluation of fetal well-being in SGA and AGA fetuses. *Am J Obstet Gynecol* 2015; **213**: 5-15.
23. Hernandez-Andrade E, Benavides Serralde JA, Cruz-Martinez R. Can anomalies of fetal brain circulation be useful in the management of growth restricted fetuses? *Prenat Diagn* 2012; **32**: 103-112.
24. Seravalli V, Baschat AA. A uniform management approach to optimize outcome in fetal growth restriction. *Obstet Gynecol Clin North Am* 2015; **42**: 275-288.
25. Morales-Roselló J, Khalil A. Fetal cerebral redistribution: a marker of compromise regardless of fetal size. *Ultrasound Obstet Gynecol* 2015; **46**: 385-388.
26. Khalil A, Thilaganathan B. Role of uteroplacental and fetal Doppler in identifying fetal growth restriction at term. *Best Pract Res Clin Obstet Gynaecol* 2017; **38**: 38-47.
27. Figueras F, Gratacos E. An integrated approach to fetal growth restriction. *Best Pract Res Clin Obstet Gynaecol* 2017; **38**: 48-58.
28. Nawathe A, Lees C. Early onset fetal growth restriction. *Best Pract Res Clin Obstet Gynaecol* 2017; **38**: 24-37.
29. Dall'Asta A, Brunelli V, Prefumo F, Frusca T, Lees CC. Early onset fetal growth restriction. *Matern Health Neonatol Perinatol* 2017; **3**: 2.
30. Bossuyt PM, Reitsma JB, Bruns DE, Gatsonis CA, Glasziou PP, Irwig LM, Lijmer JG, Moher D, Rennie D, de Vet HC; Standards for Reporting of Diagnostic Accuracy.. Towards complete and accurate reporting of studies of diagnostic accuracy: the STARD initiative. *BMJ* 2003; **326**: 41-44.
31. Leeflang MM, Deeks JJ, Gatsonis C, Bossuyt PM; Cochrane Diagnostic Test Accuracy Working Group. Systematic reviews of diagnostic test accuracy. *Ann Intern Med* 2008; **149**: 889-897.
32. Whiting PF, Rutjes AW, Westwood ME, Mallett S, Deeks JJ, Reitsma JB, Leeflang MM, Sterne JA, Bossuyt PM; QUADAS-2 Group QUADAS-2: a revised tool for the quality assessment of diagnostic accuracy studies. *Ann Intern Med* 2011; **155**: 529-536.
33. Whiting P, Harbord R, Kleijnen J. No role for quality scores in systematic reviews of diagnostic accuracy studies. *BMC Med Res Methodol* 2005; **5**: 19.
34. Sankey S, Weisfiels L, Fine M, Kapoor W. An assessment of the use of the continuity correction for sparse data in meta-analysis. *Commun Stat Simulation Computation* 1996; **25**: 1031-1056.

35. Rutter CM, Gatsonis CA. A hierarchical regression approach to meta-analysis of diagnostic test accuracy evaluations. *Stat Med* 2001; **20**: 2865-2884.
36. Reitsma JB, Glas AS, Rutjes AW, Scholten RJ, Bossuyt PM, Zwinderman AH. Bivariate analysis of sensitivity and specificity produces informative summary measures in diagnostic reviews. *J Clin Epidemiol* 2005; **58**: 982-990.
37. Walter SD. Properties of the summary receiver operating characteristic (SROC) curve for diagnostic test data. *Stat Med* 2002; **21**: 1237-1256.
38. Zwinderman AH, Bossuyt PM. We should not pool diagnostic likelihood ratios in systematic reviews. *Stat Med* 2008; **27**: 687-697.
39. Jaeschke R, Guyatt GH, Sackett DL. Users' guides to the medical literature. III. How to use an article about a diagnostic test. B. What are the results and will they help me in caring for my patients? The Evidence-Based Medicine Working Group. *JAMA* 1994; **271**: 703-707.
40. Macaskill P, Gatsonis C, Deeks JJ, Harbord RM, Takwoingi Y. Analysing and Presenting Results. In *Cochrane Handbook for Systematic Reviews of Diagnostic Test Accuracy* Version 1.0, Deeks JJ, Bossuyt PM, Gatsonis C (eds), Chapter 10. The Cochrane Collaboration, 2010.
41. Lijmer JG, Bossuyt PM, Heisterkamp SH. Exploring sources of heterogeneity in systematic reviews of diagnostic tests. *Stat Med* 2002; **21**: 1525-1537.
42. Deeks JJ, Macaskill P, Irwig L. The performance of tests of publication bias and other sample size effects in systematic reviews of diagnostic test accuracy was assessed. *J Clin Epidemiol* 2005; **58**: 882-893.
43. Gramellini D, Folli MC, Raboni S, Vadora E, Merialdi A. Cerebral-umbilical Doppler ratio as a predictor of adverse perinatal outcome. *Obstet Gynecol* 1992; **79**: 416-420.
44. Makhseed M, Jirous J, Ahmed MA, Viswanathan DL. Middle cerebral artery to umbilical artery resistance index ratio in the prediction of neonatal outcome. *Int J Gynaecol Obstet* 2000; **71**: 119-125.
45. Sterne G, Shields LE, Dubinsky TJ. Abnormal fetal cerebral and umbilical Doppler measurements in fetuses with intrauterine growth restriction predicts the severity of perinatal morbidity. *J Clin Ultrasound* 2001; **29**: 146-151.
46. Makh DS, Harman CR, Baschat AA. Is Doppler prediction of anemia effective in the growth-restricted fetus? *Ultrasound Obstet Gynecol* 2003; **22**: 489-492.
47. Odibo AO, Riddick C, Pare E, Stamilio DM, Macones GA. Cerebroplacental Doppler ratio and adverse perinatal outcomes in intrauterine growth restriction: evaluating the impact of using gestational age-specific reference values. *J Ultrasound Med* 2005; **24**: 1223-1228.

48. Jugović D, Tumbri J, Medić M, Jukić MK, Kurjak A, Arbeille P, Salihagić-Kadić A.. New Doppler index for prediction of perinatal brain damage in growth-restricted and hypoxic fetuses. *Ultrasound Obstet Gynecol* 2007; **30**: 303-311.
49. Manogura AC, Turan O, Kush ML, Berg C, Bhide A, Turan S, Moyano D, Bower S, Nicolaides KH, Galan HL, Müller T, Thilaganathan B, Gembruch U, Harman CR, Baschat AA. Predictors of necrotizing enterocolitis in preterm growth-restricted neonates. *Am J Obstet Gynecol* 2008; **198**: 638.e1-5.
50. Cruz-Martínez R, Figueras F, Hernandez-Andrade E, Oros D, Gratacos E. Fetal brain Doppler to predict cesarean delivery for nonreassuring fetal status in term small-for-gestational-age fetuses. *Obstet Gynecol* 2011; **117**: 618-626.
51. Fu J, Olofsson P. Relations between fetal brain-sparing circulation, oxytocin challenge test, mode of delivery and fetal outcome in growth-restricted term fetuses. *Acta Obstet Gynecol Scand* 2011; **90**: 227-230.
52. Marsoosi V, Bahadori F, Esfahani F, Ghasemi-Rad M. The role of Doppler indices in predicting intra ventricular hemorrhage and perinatal mortality in fetal growth restriction. *Med Ultrason* 2012; **14**: 125-132.
53. Odibo AO, Goetzinger KR, Cahill AG, Odibo L, Macones GA. Combined sonographic testing index and prediction of adverse outcome in preterm fetal growth restriction. *Am J Perinatol* 2014; **31**: 139-144.
54. Lobmaier SM, Figueras F, Mercade I, Perello M, Peguero A, Crovetto F, Ortiz JU, Crispi F, Gratacós E. Angiogenic factors vs Doppler surveillance in the prediction of adverse outcome among late-pregnancy small-for-gestational-age fetuses. *Ultrasound Obstet Gynecol* 2014; **43**: 533-540.
55. Spinillo A, Gardella B, Bariselli S, Alfei A, Silini EM, Bello BD. Cerebroplacental Doppler ratio and placental histopathological features in pregnancies complicated by fetal growth restriction. *J Perinat Med* 2014; **42**: 321-328.
56. Flood K, Unterscheider J, Daly S, Geary MP, Kennelly MM, McAuliffe FM, O'Donoghue K, Hunter A, Morrison JJ, Burke G, Dicker P, Tully EC, Malone FD. The role of brain sparing in the prediction of adverse outcomes in intrauterine growth restriction: results of the multicenter PORTO Study. *Am J Obstet Gynecol* 2014; **211**: 288.e1-5.
57. Crimmins S, Fitzgerald G, Block-Abraham D, Atkins K, Harman C, Turan O. Use of cerebroplacental ratio to predict outcome in late onset fetal growth restriction (FGR). *Am J Obstet Gynecol* 2015; **212**(Suppl 1): S269-S270.
58. Babic I, Alameri S, Tulbah M, Moretti F, Kurdi W. Cerebroplacental ratio and its association with adverse perinatal outcomes in early and late onset

- intrauterine growth restriction (IUGR). *Ultrasound Obstet Gynecol* 2015; **46**(Suppl 1): 149.
59. Figueras F, Savchev S, Triunfo S, Crovetto F, Gratacos E. An integrated model with classification criteria to predict small-for-gestational-age fetuses at risk of adverse perinatal outcome. *Ultrasound Obstet Gynecol* 2015; **45**: 279-285.
60. Regan J, Masters H, Warshak C. Association between an abnormal cerebroplacental ratio and the development of severe pre-eclampsia. *J Perinatol* 2015; **35**: 322-327.
61. Warshak CR, Masters H, Regan J, DeFranco E. Doppler for growth restriction: the association between the cerebroplacental ratio and a reduced interval to delivery. *J Perinatol* 2015; **35**: 332-337.
62. Garcia-Simon R, Figueras F, Savchev S, Fabre E, Gratacos E, Oros D. Cervical condition and fetal cerebral Doppler as determinants of adverse perinatal outcome after labor induction for late-onset small-for-gestational-age fetuses. *Ultrasound Obstet Gynecol* 2015; **46**: 713-717.
63. Starčević M, Predojević M, Butorac D, Tumbri J, Konjevoda P, Kadić AS. Early functional and morphological brain disturbances in late-onset intrauterine growth restriction. *Early Hum Dev* 2016; **93**: 33-38.
64. Sirico A, Diemert A, Glosemeyer P, Hecher K. Prediction of adverse perinatal outcome by cerebroplacental ratio adjusted for estimated fetal weight. *Ultrasound Obstet Gynecol* 2018; **51**: 381-386.
65. Lees CC, Marlow N, van Wassenaer-Leemhuis A, Arabin B, Bilardo CM, Brezinka C, Calvert S, Derks JB, Diemert A, Duvkot JJ, Ferrazzi E, Frusca T, Ganzevoort W, Hecher K, Martinelli P, Ostermayer E, Papageorghiou AT, Schlembach D, Schneider KT, Thilaganathan B, Todros T, Valcamonico A, Visser GH, Wolf H; TRUFFLE study group. 2 year neurodevelopmental and intermediate perinatal outcomes in infants with very preterm fetal growth restriction (TRUFFLE): a randomised trial. *Lancet* 2015; **385**: 2162-2172.
66. Stampalija T, Arabin B, Wolf H, Bilardo CM, Lees C; TRUFFLE investigators. Is middle cerebral artery Doppler related to neonatal and 2-year infant outcome in early fetal growth restriction? *Am J Obstet Gynecol* 2017; **216**: 521.e1-521.e13.
67. Baschat AA, Viscardi RM, Hussey-Gardner B, Hashmi N, Harman C. Infant neurodevelopment following fetal growth restriction: relationship with antepartum surveillance parameters. *Ultrasound Obstet Gynecol* 2009; **33**: 44-50.
68. Constantinescu S, Denes M, Vladareanu R. Intrauterine growth restriction: Perinatal assessment in predicting the offspring neurologic impairment. A 2 years prospective study. *Ginecoeu* 2013; **9**: 35-40.

69. Meher S, Hernandez-Andrade E, Basheer SN, Lees C. Impact of cerebral redistribution on neurodevelopmental outcome in small-for-gestational-age or growth-restricted babies: a systematic review. *Ultrasound Obstet Gynecol* 2015; **46**: 398-404.
70. Nassr AA, Abdelmagied AM, Shazly SA. Fetal cerebro-placental ratio and adverse perinatal outcome: systematic review and meta-analysis of the association and diagnostic performance. *J Perinat Med* 2016; **44**: 249-256.
71. Dunn L, Sherrell H, Kumar S. Review: Systematic review of the utility of the fetal cerebroplacental ratio measured at term for the prediction of adverse perinatal outcome. *Placenta* 2017; **54**: 68-75.
72. Vollgraff Heidweiller-Schreurs CA, de Boer MA, Heymans MW, Schoonmade LJ, Bossuyt PMM, Mol BWJ, De Groot CJM, Bax CJ. Prognostic accuracy of cerebroplacental ratio and middle cerebral artery Doppler for adverse perinatal outcomes: a systematic review and meta-analysis. *Ultrasound Obstet Gynecol* 2018; **51**: 313-322.
73. Morris RK, Malin G, Robson SC, Kleijnen J, Zamora J, Khan KS. Fetal umbilical artery Doppler to predict compromise of fetal/neonatal wellbeing in a high-risk population: systematic review and bivariate meta-analysis. *Ultrasound Obstet Gynecol* 2011; **37**: 135-142.
74. Morris RK, Say R, Robson SC, Kleijnen J, Khan KS. Systematic review and meta-analysis of middle cerebral artery Doppler to predict perinatal wellbeing. *Eur J Obstet Gynecol Reprod Biol* 2012; **165**: 141-155.

FIGURE LEGENDS

Legend for Figure 1: Study selection process

Legend for Figure 2: Risk of bias of included studies

Legend for Figure 3: Summary ROC curves of the cerebroplacental ratio to predict adverse perinatal outcomes in suspected fetal growth restriction

Legend for Figure S1: Forest plots of the cerebroplacental ratio to predict adverse perinatal outcomes in suspected fetal growth restriction