





International standards for fetal brain structures based on serial ultrasound measurements from Fetal Growth Longitudinal Study of INTERGROWTH-21st Project

R. NAPOLITANO¹ , M. MOLLOHOLLI¹, V. DONADONO¹, E. O. OHUMA^{1,2,3}, S. Z. WANYONYI¹, B. KEMP¹, M. K. YAQUB⁴, S. ASH¹, F. C. BARROS⁵, M. CARVALHO⁶, Y. A. JAFFER⁷, J. A. NOBLE⁴, M. OBERTO⁸, M. PURWAR⁹, R. PANG¹⁰, L. CHEIKH ISMAIL¹¹, A. LAMBERT¹, M. G. GRAVETT¹², L. J. SALOMON¹³, Z. A. BHUTTA¹⁴, S. H. KENNEDY^{1,2}, J. VILLAR^{1,2} and A. T. PAPAGEORGHIOU^{1,2} ; International Fetal and Newborn Growth Consortium for the 21st Century (INTERGROWTH-21st)

¹Nuffield Department of Women's & Reproductive Health, University of Oxford, Oxford, UK; ²Oxford Maternal & Perinatal Health Institute, Green Templeton College, University of Oxford, Oxford, UK; ³Centre for Statistics in Medicine, Botnar Research Centre, University of Oxford, Oxford, UK; ⁴Department of Engineering Science, Institute of Biomedical Engineering, University of Oxford, Oxford, UK; ⁵Programa de Pós-Graduação em Saúde e Comportamento, Universidade Católica de Pelotas, Pelotas, Brazil; ⁶Faculty of Health Sciences, Aga Khan University, Nairobi, Kenya; ⁷Department of Family & Community Health, Ministry of Health, Muscat, Sultanate of Oman; ⁸S.C. Ostetricia 2U, Città della Salute e della Scienza di Torino, Italy; ⁹Nagpur INTERGROWTH-21st Research Centre, Ketkar Hospital, Nagpur, India; ¹⁰School of Public Health, Peking University, Beijing, China; ¹¹Clinical Nutrition and Dietetics Department, University of Sharjah, Sharjah, United Arab Emirates; ¹²Departments of Obstetrics and Gynecology, and of Public Health, University of Washington, Seattle, WA, USA; ¹³Department of Obstetrics and Fetal Medicine, Hôpital Necker Enfants Malades, Université Paris Descartes, Paris, France; ¹⁴Center for Global Child Health, Hospital for Sick Children, Toronto, Canada

KEYWORDS: anterior horn of lateral ventricle; atrium; cisterna magna; fetal brain structures; growth; parieto-occipital fissure; screening; Sylvian fissure

CONTRIBUTION

What are the novel findings of this work?

Charts of fetal brain measurements in current use are based mostly on small studies with suboptimal methodology and no follow-up. In this international study, we created standards for the size of five fetal brain structures based on a prospective cohort of fetuses followed up into childhood, demonstrating normal neurodevelopment.

What are the clinical implications of this work?

Clinical use of such objective fetal brain structure measurements may help to improve the screening and diagnostic performance of prenatal ultrasonography. It should also allow a unified approach to fetal assessment by integration with other standards from the same population and result in a common language when describing aberrations from expected norms.

ABSTRACT

Objective To create prescriptive growth standards for five fetal brain structures, measured using ultrasound, in healthy, well-nourished women at low risk of impaired fetal growth and poor perinatal outcome, taking part in the Fetal Growth Longitudinal Study (FGLS) of the INTERGROWTH-21st Project.

Methods This was a complementary analysis of a large, population-based, multicenter, longitudinal study. The sample analyzed was selected randomly from the overall FGLS population, ensuring an equal distribution among the eight diverse participating sites and of three-dimensional (3D) ultrasound volumes across pregnancy (range: 15–36 weeks' gestation). We measured, in planes reconstructed from 3D ultrasound volumes of the fetal head at different timepoints in pregnancy, the size of the parieto-occipital fissure (POF), Sylvian fissure

Correspondence to: Dr A. T. Papageorghiou, Nuffield Department of Women's & Reproductive Health, University of Oxford, The Women's Centre, John Radcliffe Hospital, Oxford, OX3 9DU, UK (e-mail: aris.papageorghiou@wrh.ox.ac.uk)

Accepted: 27 January 2020

(SF), anterior horn of the lateral ventricle, atrium of the posterior horn of the lateral ventricle (PV) and cisterna magna (CM). Fractional polynomials were used to construct the standards. Growth and development of the infants were assessed at 1 and 2 years of age to confirm their adequacy for constructing international standards.

Results From the entire FGLS cohort of 4321 women, 451 (10.4%) were selected at random. After exclusions, 3D ultrasound volumes from 442 fetuses born without a congenital malformation were used to create the charts. The fetal brain structures of interest were identified in 90% of cases. All structures, except the PV, showed increasing size with gestational age, and the size of the POF, SF, PV and CM showed increasing variability. The 3rd, 5th, 50th, 95th and 97th smoothed centiles are presented. The 5th centiles for the POF and SF were 3.1 mm and 4.7 mm at 22 weeks' gestation and 4.6 mm and 9.9 mm at 32 weeks, respectively. The 95th centiles for the PV and CM were 8.5 mm and 7.5 mm at 22 weeks and 8.6 mm and 9.5 mm at 32 weeks, respectively.

Conclusions We have produced prescriptive size standards for fetal brain structures based on prospectively enrolled pregnancies at low risk of abnormal outcome. We recommend these as international standards for the assessment of measurements obtained using ultrasound from fetal brain structures. © 2020 The Authors. *Ultrasound in Obstetrics & Gynecology* published by John Wiley & Sons Ltd on behalf of the International Society of Ultrasound in Obstetrics and Gynecology.

INTRODUCTION

In most settings, the anatomy of the fetal brain is assessed routinely as part of the mid-trimester anomaly scan at around 20 weeks' gestation, the main aims being to demonstrate anatomical integrity and diagnose abnormalities of the central nervous system (CNS). Measurement of intracranial structures forms part of the assessment, and includes the width of the atrium of the lateral ventricle measured posteriorly (PV) and cisterna magna (CM)^{1,2}. On more advanced neurosonography, undertaken for indications such as a previous or suspected abnormality, other structures, e.g. the Sylvian fissure (SF), are examined either earlier in cases of a previous abnormality or late in pregnancy to assess gyration and sulcation patterns, which change with advancing gestational age^{3–8}.

Fetal brain structures can be evaluated by assessing their appearance subjectively or measured quantitatively, which is recommended whenever possible, as subjective assessment is associated with higher variability². Currently, the normality of any measurements obtained is evaluated in relation to one of several reported reference charts for fetal brain structures². However, many studies reporting reference charts have important methodological limitations⁹. There can also be a lack of consistency in the interpretation of ultrasound images of the fetal

CNS, leading to inconsistent clinical management, if the same measurement from a fetus is plotted on two different charts. These issues are generic to the measurement of all fetal anatomical structures, as reported in systematic reviews of studies aimed at creating charts for fetal biometry and pregnancy dating^{10,11}.

To overcome these issues with regard to ultrasound assessment of the fetal brain, we have followed, as before, World Health Organization (WHO) recommendations and adopted a prescriptive approach to the construction of international size standards for five fetal brain structures, as a secondary analysis of data collected in the Fetal Growth Longitudinal Study (FGLS), one of the key components of the INTERGROWTH-21st Project (www.intergrowth21.org.uk)¹². Three of the brain structures relate to clinical evaluation of cerebrospinal fluid, namely the PV, CM and anterior horn of the lateral ventricle (AV)¹; the two other structures are clinically relevant to the assessment of gyration and sulcation, namely the parieto-occipital fissure (POF) and the SF. The international standards produced complement those published previously for early and late pregnancy dating^{13,14}, fetal growth and estimated fetal weight^{15,16}, symphysis–fundal height¹⁷, gestational weight gain¹⁸, neonatal size and body composition^{19,20} and postnatal growth of preterm infants²¹.

PATIENTS AND METHODS

Study population

INTERGROWTH-21st is an international, multicenter, population-based project, conducted between 2009 and 2016 in eight delimited geographical areas: Pelotas (Brazil), Turin (Italy), Muscat (Oman), Oxford (UK), Seattle (USA), Shunyi County in Beijing (China), the central area of Nagpur (India) and the Parklands suburb of Nairobi (Kenya). In the FGLS, serial two-dimensional (2D) and three-dimensional (3D) fetal scans were performed every 5 ± 1 weeks from 14 + 0 weeks' gestation to delivery¹⁵. Women participating in the study, who initiated antenatal care before 14 weeks' gestation, were selected based upon the WHO recommended criteria for optimal health, nutrition, education and socioeconomic status needed to construct international standards^{22,23}. Hence, they had a low-risk pregnancy that fulfilled well defined and strict inclusion criteria at both population and individual levels²³. Briefly, the individual inclusion criteria were maternal age between 18 and 35 years, body mass index ≥ 18.5 kg/m² and < 30 kg/m², naturally conceived singleton pregnancy, normal pregnancy history without relevant past medical history, no evidence of socioeconomic constraints likely to impede fetal growth, no use of tobacco or recreational drugs and no heavy alcohol consumption. Women also had to have a known date of their last menstrual period (LMP), with regular cycles without the use of hormonal contraceptives or breastfeeding in the 2 months before pregnancy. Gestational age was LMP-based provided that

standardized ultrasound measurement of crown–rump length between 9 + 0 and 14 + 0 weeks was in agreement within 7 days²⁴.

In the FGLS all ultrasound scans were performed by sonographers who were trained, standardized and audited regularly^{25,26}. The same type of commercially available ultrasound equipment (Philips HD-9; Philips Ultrasound, Bothell, WA, USA), with curvilinear abdominal 2D transducers (C5-2, C6-3) and a curvilinear abdominal 3D transducer (V7-3), was used for all growth scans. For the purposes of the INTERGROWTH-21st Project, the manufacturer reprogrammed the machines' software to ensure that the measurement values did not appear on the screen during the scan in order to reduce operator 'expected value' bias. A detailed description of the ultrasound methodology has been reported previously²⁵.

Infants from sites that participated in the follow-up study (Brazil, India, Italy, Kenya and the UK) were assessed at the ages of 1 and 2 years to obtain a detailed evaluation of growth, nutrition, morbidity and motor development. These data were collected by interviewing parents and assessment by a certified examiner. Achievement of milestones ('sitting without support', 'standing with assistance', 'hand-and-knees-crawling', 'walking with assistance', 'standing alone' and 'walking alone') were considered normal if the age at achievement was within the expected WHO windows (less than the 99th centile for each of the expected windows)²⁷.

The INTERGROWTH-21st Project was approved by the Oxfordshire Research Ethics Committee "C" (ref: 08/H0606/139), the research ethics committees of the individual institutions and the regional health authorities in which the project was implemented; all the women involved gave written informed consent.

Structures measured and sample-size considerations

The fetal brain structures were measured on ultrasound images extracted from 3D volumes of the fetal head, acquired at all eight participating sites. The decision regarding which structures to evaluate was based on a combination of factors: an extensive scoping exercise and review of the literature demonstrating their clinical utility⁹; structures that can be assessed in axial planes that are acquired routinely; and a pilot study involving 90 ultrasound volumes assessing feasibility and reproducibility.

The sample size was based on pragmatic and statistical considerations. The main pragmatic consideration was the

considerable length of time required for volume upload, manipulation, plane extraction and measurement (20 min per volume on average). As a result, we decided to take a random sample from the entire FGLS cohort, bearing in mind the need for precision at the 5th and 95th centiles. A sample of 300 scans would obtain a precision of 0.1 SD at the 5th or the 95th centile²⁸. Using conservative estimates, we assumed a possible 5% exclusion rate due to loss to follow-up in pregnancy or at birth, withdrawal of consent, miscarriage, stillbirth, maternal death, fetal or neonatal structural abnormality or severely abnormal outcome at 2-year follow-up, which was defined as any of the following: meningitis, hearing loss, blindness or major visual problems, seizures, cerebral palsy, neurological disorders, malignancy, malaria, tuberculosis, hepatitis, HIV/AIDS, cystic fibrosis or hemolytic conditions. We also assumed that, in up to 40% of cases, all five structures might not be measurable (based on a conservative estimate, as the actual upper limit of the confidence interval from the pilot study was 20%, primarily due to movement artifact). Based on these assumptions, we estimated that 451 3D volumes would lead to a minimum of 300 measurements for each structure. Therefore, we selected 451 3D volumes from the overall FGLS population using computer randomization, ensuring an equal distribution among the eight participating sites and of ultrasound volumes across pregnancy (range: 15–36 weeks' gestation). The random selection was performed using SAS software© (SAS Institute Inc., Cary, NC, USA).

The study was cross-sectional, as only one volume per pregnancy was included.

Volume acquisition, offline analysis and quality control

Detailed descriptions of the volume acquisition methods are provided elsewhere^{25,29}. Briefly, head volumes were acquired at the level of the axial transthalamic plane. Five predefined quality-control criteria for the transthalamic plane had to be satisfied to acquire the volume (Table 1; Figure 1)²⁶. Acquisition was undertaken with the volume data box and angle of sweep (usually 70°) adjusted to include the entire head during fetal quiescence, and with the mother asked to hold her breath and the transducer held steady. The real-time image was observed during acquisition to confirm that the sweep included the entire head with no maternal or fetal movement during the sweep, otherwise the process was repeated. All data were

Table 1 Quality criteria for acquisition of the three planes of the fetal brain on ultrasound

<i>Transthalamic plane</i>	<i>Transventricular plane</i>	<i>Transcerebellar plane</i>
Symmetrical hemispheres	Symmetrical hemispheres	Symmetrical hemispheres
Cavum of the septum pellucidum present	Cavum of the septum pellucidum present	Cavum of the septum pellucidum present
Thalami visible	Lateral ventricles visible	Thalami visible
No cerebellum visible	No cerebellum visible	Cerebellum present at maximum diameter
Magnification of 30% image	Magnification of 30% image	Magnification of 30% image




Figure 1 Reconstructed ultrasound planes showing caliper placement for fetal brain structure acquisition at different gestational ages for transventricular plane (a), transthalamic plane (b) and transcerebellar plane (c). AV, anterior horn of lateral ventricle; CM, cisterna magna; POF, parieto-occipital fissure; PV, atrium of posterior horn of lateral ventricle; SF, Sylvian fissure; TCD, transcerebellar diameter; w, completed weeks' gestation.

then sent to the Ultrasound Quality Coordinating Unit in Oxford.

Offline analysis was undertaken by four experienced sonographers at the coordinating unit. All were trained in neurosonography and their training specifically standardized for the purposes of this study in volume manipulation for plane reconstruction and measurement (Videoclip S1, Figure 1). The volume manipulations and measurements were performed using the software of the ultrasound machines' manufacturer or an open-source image analysis software program (Medical Imaging Interaction Toolkit MITK, version 0.12.2; German Cancer Research Center, Division of Medical and Biological Informatics, www.mitk.org)³⁰. This was done because the open-source software was more 'user friendly'. Comparability of measurements between the manufacturer's software and the open-source image-analysis software program was confirmed (mean reproducibility was within 0.7 mm).

All sonographers were blinded to the measurements during the study. In addition, strict quality control was undertaken in the whole sample: image quality criteria were used to ensure the maximum possible score for each extracted plane (Table 1) before measurement of the following five structures: the POF and SF in the transthalamic plane; the AV and PV in the transventricular plane; and the CM in the transcerebellar plane. The POF, SF, AV and PV were measured in the distal hemisphere of the respective plane (because of poorer visualization in the proximal hemisphere). Further details

of volume manipulation and caliper placement are given in Appendix S1.

Reproducibility

Reproducibility was assessed in a subset of 90 volumes. The first sonographer uploaded the volume, manually extracted the three planes and measured the five structures twice (intraobserver reproducibility for plane reconstruction and measurement acquisition). A second sonographer re-uploaded the same volume and repeated this process (this second set of data was used to assess interobserver reproducibility for plane reconstruction and measurement acquisition). To assess the contribution of caliper replacement, the second sonographer replaced the calipers on still images and repositioned them to measure all structures in each plane stored by the first sonographer (interobserver reproducibility for caliper replacement on stored images). As in the main study, all sonographers were blinded to their own and the other sonographer's measurements during the reproducibility study.

Statistical analysis

We followed the modeling approach used previously by our group to construct fetal growth charts¹⁵. In summary, fractional polynomials that model the means and SD were used to model biometric measurements of brain structures as a function of gestational age. Our overall aim was to

produce centiles that change smoothly with age and maximize simplicity without compromising model fit. Goodness of fit was assessed by Q–Q plots and a scatterplot of Z-scores by gestational age. Mean differences between the observed and fitted centiles were also calculated.

For the reproducibility study, Bland–Altman plots were used to quantify the level of agreement and variability in the measurements. Differences between and within observers were expressed in absolute values (mm). Analysis was performed using Stata 11 (StataCorp., College Station, TX, USA).

RESULTS

After exclusions, 442/451 (98.0%) volumes were used to reconstruct planes and create the fetal brain charts (Figure 2). No congenital malformations were detected antenatally or postnatally in the selected fetuses, and no infants met the exclusion criteria set for the 2-year follow-up. As expected, given the random selection, maternal demographics and pregnancy outcomes were similar to those in the overall FGLS population, confirming a low risk of perinatal complications (Table S1).

Of the 442 infants, 297 (67.2%) were assessed by their parent(s) at 1 year of age; of these, 289 (97.3%) were also assessed by a certified examiner at a mean age of 12.3 months (range, 10.9–19.4 months). As reported by the parent(s), 99% of the infants had entirely normal motor development. Only three (1%) infants did not achieve the milestones ‘sitting without support’ and ‘standing with assistance’; brain structure measurements for these children were within the 5th and 95th centile range. There was overall good agreement between the achievement of milestones as reported by the parent(s) and that found on assessment by a certified examiner (average agreement, 96% (range, 92–100%)). Reassuringly, in almost all cases in which disagreement between the two assessments was present, the examiner reported more precocious milestone achievement than did the parent(s), confirming the low risk for abnormal long-term outcome in our cohort. Follow-up at 2 years of age was available in 304 children; the findings of this detailed assessment demonstrate comparability with the morbidity reported in children from the overall FGLS cohort who underwent motor and neurodevelopment assessment (Table S2; Figure 3)³¹. The mean and SD of the children’s weight, length and head circumference at 2 years of age were 12.3 ± 1.7 kg, 87.4 ± 3.7 cm and 47.7 ± 1.6 cm, respectively, and Z-scores were within the expected values of the WHO Child Growth Standards. Motor development for the two milestones not reached by the age of 1 year (‘standing alone’ and ‘walking alone’) was confirmed as normal at 2 years in 99% and 98%, respectively.

In total, 2439 measurements of fetal brain structures were acquired. On average, structures were measurable in a high-quality extracted plane in 90% of cases, the CM being the structure measurable the least frequently. After removal of outliers, measurements were available to create centiles for the POF, SF, AV, PV and CM in




Figure 2 Flowchart summarizing inclusion of pregnancies for development of fetal brain charts. FGLS, Fetal Growth Longitudinal Study.




Figure 3 Forest plot showing median (3rd and 97th centiles) age at achievement of four gross motor developmental milestones in children included in INTERGROWTH-21st Fetal Growth Standards study (●) and those included in current study (■). For comparison, median, 3rd and 97th centiles of WHO windows for achievement of the same milestones are presented as gray bars.

420 (95%), 404 (91%), 378 (86%), 422 (95%) and 352 (80%) cases, respectively. The overall mean (SD) measurements were POF, 5.47 ± 1.91 mm; SF, 9.45 ± 4.22 mm; AV, 7.61 ± 1.54 mm; PV, 6.00 ± 1.59 mm; and CM, 5.27 ± 1.66 mm. All fetal brain measurements were normally distributed conditional on gestational age.




Figure 4 Fitted 5th, 50th and 95th smoothed centile curves of fetal brain structure measurements: (a) parieto-occipital fissure; (b) Sylvian fissure; (c) anterior horn of lateral ventricle (AV); (d) atrium of posterior horn of lateral ventricle (PV); (e) cisterna magna.

The best fitting powers were provided by second-degree fractional polynomials and further modeled in a multilevel framework to account for the cross-sectional design of the study. The gestational-age-specific smoothed centiles for the POF, SF, AV, PV and CM are presented in Figure 4 and Tables 2–6. One fetus had a PV > 10 mm and four had a CM > 10 mm; all had normal perinatal outcome.

Both visual assessment of scatterplots of Z-scores by gestational age and goodness-of-fit tests, assessed by gestational-age-specific comparisons of empirical centiles with smoothed centile curves, showed good agreement.

The equations for the mean and SD from the fractional polynomial regression models for each structure measured are presented in Table 7, allowing for the calculation of any desired centile according to gestational age in exact weeks.

Results of the reproducibility study are shown in Table 8. All measurements were reproducible within less than 3 mm or 12% (all mean differences were less than 0.1 mm or 0.5%). The greatest proportion of variability was due to caliper replacement, accounting for more than 50% of the intra- and interobserver variability for measurements of all structures, as observed previously for fetal biometry measurements (Figure S1)³².

DISCUSSION

We have produced international size standards for ultrasound measurements of clinically relevant fetal brain structures. The study population consisted of women at low risk of adverse pregnancy and perinatal outcomes¹⁵. Unlike previous studies reporting fetal brain standards, we followed up the infants and demonstrated satisfactory growth and development at 1 and 2 years of age, confirming that our initial selection criteria met the WHO requirements for constructing international growth standards^{12,31}. The sequence and timing of the attainment of neurodevelopmental milestones and associated behaviors in early childhood were very similar to those reported previously by our group, i.e. we have demonstrated that there are similarities across diverse geographical regions, as long as nutritional and health needs are met³¹.

We performed a systematic review of the literature that assessed the methodology used to create fetal brain structure charts⁹. This showed that some studies did not strictly adhere to plane standardization. Using different planes for fetal head biometry can lead to significant measurement differences³³. In some studies, landmarks for plane acquisition were not specified^{34–44}, while in others, various oblique planes with numerous landmarks were proposed^{45,46}. One of the strengths of our study is the use of standardized axial planes recommended in routine

Table 2 Smoothed centiles for parieto-occipital fissure (mm), according to exact gestational age (GA)

GA (weeks)	Sample size (n)	Centile				
		3 rd	5 th	50 th	95 th	97 th
15+0	18	1.29	1.39	2.14	2.88	2.99
16+0	18	1.55	1.69	2.66	3.63	3.77
17+0	18	1.79	1.96	3.12	4.28	4.45
18+0	19	2.02	2.21	3.53	4.85	5.04
19+0	19	2.24	2.45	3.90	5.35	5.56
20+0	21	2.44	2.67	4.24	5.80	6.03
21+0	16	2.64	2.87	4.54	6.21	6.44
22+0	18	2.82	3.07	4.82	6.57	6.82
23+0	21	2.99	3.25	5.08	6.90	7.17
24+0	18	3.16	3.43	5.32	7.21	7.48
25+0	20	3.32	3.59	5.54	7.49	7.77
26+0	19	3.47	3.75	5.75	7.75	8.04
27+0	19	3.61	3.90	5.95	7.99	8.29
28+0	19	3.75	4.05	6.13	8.22	8.52
29+0	22	3.89	4.19	6.31	8.43	8.74
30+0	21	4.02	4.32	6.48	8.63	8.94
31+0	20	4.14	4.46	6.64	8.83	9.14
32+0	17	4.27	4.58	6.80	9.01	9.32
33+0	22	4.39	4.71	6.94	9.18	9.50
34+0	21	4.51	4.83	7.09	9.35	9.67
35+0	19	4.62	4.95	7.23	9.51	9.84
36+0	15	4.74	5.07	7.37	9.67	9.99
Total measurements	420					

Table 3 Smoothed centiles for Sylvian fissure (mm), according to exact gestational age (GA)

GA (weeks)	Sample size (n)	Centile				
		3 rd	5 th	50 th	95 th	97 th
15+0	18	0.40	0.57	1.77	2.98	3.15
16+0	15	0.91	1.13	2.65	4.17	4.38
17+0	18	1.46	1.72	3.49	5.27	5.52
18+0	18	2.03	2.31	4.31	6.30	6.59
19+0	17	2.60	2.91	5.09	7.27	7.58
20+0	20	3.18	3.51	5.85	8.18	8.51
21+0	15	3.75	4.10	6.57	9.04	9.40
22+0	18	4.32	4.69	7.27	9.86	10.23
23+0	20	4.87	5.26	7.95	10.64	11.02
24+0	17	5.42	5.82	8.60	11.38	11.78
25+0	20	5.96	6.37	9.23	12.09	12.50
26+0	18	6.49	6.91	9.84	12.77	13.19
27+0	16	7.01	7.44	10.43	13.42	13.85
28+0	19	7.52	7.95	11.00	14.05	14.48
29+0	22	8.01	8.45	11.55	14.65	15.09
30+0	20	8.49	8.94	12.09	15.23	15.68
31+0	20	8.97	9.42	12.61	15.79	16.25
32+0	17	9.43	9.89	13.11	16.33	16.79
33+0	22	9.88	10.34	13.60	16.86	17.32
34+0	22	10.32	10.79	14.07	17.36	17.83
35+0	18	10.75	11.22	14.54	17.85	18.33
36+0	14	11.17	11.64	14.99	18.33	18.80
Total measurements	404					

clinical practice for biometry assessment (Table 1). We believe that this approach of using standardized planes improves reproducibility, a view that is supported by the findings of previous studies^{46,47}. In our case, this led to a high proportion of structures that could be measured on stored volumes (90% on average) and

resulted in reproducible measurements, with 95% limits of agreement within < 3 mm (or < 12%) (Table 8). Studies involving experts in neurosonography report similar results in visualizing structures from volume analysis⁴⁸. This is in contrast to previous studies on subjective assessment of brain fissures, which report variable results

Table 4 Smoothed centiles for anterior horn of lateral ventricle (mm), according to exact gestational age (GA)

GA (weeks)	Sample size (n)	Centile				
		3 rd	5 th	50 th	95 th	97 th
15+0	17	4.34	4.62	6.61	8.59	8.87
16+0	15	4.39	4.67	6.65	8.63	8.91
17+0	17	4.44	4.72	6.70	8.68	8.97
18+0	18	4.49	4.78	6.76	8.74	9.02
19+0	19	4.56	4.84	6.82	8.80	9.09
20+0	20	4.63	4.91	6.89	8.87	9.16
21+0	15	4.71	4.99	6.97	8.95	9.24
22+0	18	4.79	5.08	7.06	9.04	9.32
23+0	21	4.89	5.17	7.15	9.13	9.42
24+0	15	4.99	5.27	7.25	9.24	9.52
25+0	19	5.10	5.38	7.37	9.35	9.63
26+0	18	5.22	5.51	7.49	9.47	9.75
27+0	17	5.35	5.64	7.62	9.60	9.88
28+0	19	5.49	5.78	7.76	9.74	10.02
29+0	22	5.65	5.93	7.91	9.89	10.17
30+0	19	5.81	6.09	8.07	10.05	10.34
31+0	17	5.98	6.26	8.24	10.23	10.51
32+0	13	6.17	6.45	8.43	10.41	10.69
33+0	18	6.36	6.65	8.63	10.61	10.89
34+0	17	6.57	6.85	8.84	10.82	11.10
35+0	15	6.79	7.08	9.06	11.04	11.32
36+0	9	7.03	7.31	9.29	11.27	11.56
Total measurements	378					

Table 5 Smoothed centiles for atrium of posterior horn of lateral ventricle (mm), according to exact gestational age (GA)

GA (weeks)	Sample size (n)	Centile				
		3 rd	5 th	50 th	95 th	97 th
15+0	18	4.71	4.99	6.94	8.88	9.16
16+0	19	4.49	4.77	6.78	8.78	9.07
17+0	18	4.28	4.58	6.64	8.70	9.00
18+0	19	4.10	4.40	6.52	8.64	8.94
19+0	19	3.92	4.23	6.41	8.58	8.89
20+0	22	3.76	4.08	6.31	8.54	8.86
21+0	16	3.61	3.94	6.22	8.51	8.84
22+0	18	3.46	3.80	6.14	8.49	8.82
23+0	21	3.33	3.67	6.07	8.47	8.81
24+0	18	3.20	3.55	6.00	8.46	8.81
25+0	20	3.07	3.43	5.94	8.46	8.82
26+0	19	2.95	3.32	5.89	8.46	8.83
27+0	19	2.84	3.22	5.84	8.46	8.84
28+0	19	2.73	3.11	5.79	8.48	8.86
29+0	22	2.62	3.01	5.75	8.49	8.88
30+0	21	2.52	2.92	5.71	8.51	8.91
31+0	20	2.42	2.83	5.68	8.53	8.94
32+0	17	2.32	2.74	5.65	8.55	8.97
33+0	22	2.23	2.65	5.62	8.58	9.00
34+0	22	2.14	2.57	5.59	8.61	9.04
35+0	19	2.05	2.49	5.56	8.64	9.08
36+0	14	1.96	2.41	5.54	8.67	9.12
Total measurements	422					

in terms of reproducibility (kappa coefficients varying from 0.56 to 0.95)^{45,49}. Improving reproducibility was also one of the aims of our study, in order to move to quantitative assessment of fetal brain development^{45,46,50}.

To achieve our objectives, we used international guidelines to obtain measurements of the PV and CM^{1,2}, and we provide detailed methods for AV, POF

and SF measurements, based on existing publications (Appendix S1), as we were unable to find generally accepted guidelines.

Our study overcomes many of the methodological limitations of previous studies⁹. These include a high risk of bias in the selection of the population, ultrasound protocol and data analysis. For example, fewer than

Table 6 Smoothed centiles for cisterna magna (mm), according to exact gestational age (GA)

GA (weeks)	Sample size (n)	Centile				
		3 rd	5 th	50 th	95 th	97 th
15+0	19	1.71	1.82	2.82	4.36	4.64
16+0	17	1.96	2.08	3.20	4.92	5.24
17+0	17	2.19	2.33	3.56	5.44	5.79
18+0	18	2.41	2.56	3.89	5.92	6.29
19+0	19	2.61	2.77	4.20	6.36	6.75
20+0	21	2.80	2.97	4.48	6.76	7.17
21+0	15	2.97	3.15	4.73	7.12	7.55
22+0	18	3.12	3.31	4.97	7.45	7.90
23+0	21	3.26	3.46	5.18	7.75	8.21
24+0	16	3.39	3.60	5.37	8.02	8.50
25+0	17	3.51	3.72	5.55	8.27	8.76
26+0	19	3.62	3.83	5.71	8.50	8.99
27+0	15	3.72	3.94	5.85	8.70	9.21
28+0	16	3.81	4.03	5.99	8.89	9.41
29+0	20	3.90	4.12	6.11	9.06	9.59
30+0	16	3.97	4.20	6.22	9.22	9.75
31+0	13	4.04	4.27	6.33	9.36	9.90
32+0	14	4.11	4.34	6.42	9.49	10.04
33+0	12	4.17	4.40	6.51	9.62	10.17
34+0	13	4.22	4.46	6.59	9.73	10.28
35+0	11	4.27	4.51	6.66	9.83	10.39
36+0	5	4.32	4.56	6.73	9.92	10.49
Total measurements	352					

Table 7 Equations for estimation of mean and SD (mm) of each fetal brain structure measurement, according to exact gestational age (GA) (weeks)

Structure	Equation
Parieto-occipital fissure	
Mean	$10.29428 - (122.8447 \times GA^{-1}) + (0.00001038 \times GA^3)$
SD	$1.596042 - (257.2297 \times GA^{-2})$
Sylvian fissure	
Mean	$80.27012 - (32.7877 \times GA^{-0.5}) - (100.1593 \times GA^{-0.5} \times \ln(GA))$
SD	$2.304501 - (353.814 \times GA^{-2})$
Anterior horn of lateral ventricle	
Mean	$6.396214 + (0.00006205 \times GA^3)$
SD	1.204454
Atrium of posterior horn of lateral ventricle	
Mean	$4.389214 + (38.10015 \times GA^{-1}) + (0.0000020063 \times GA^3)$
SD	$0.6707227 + (0.034258 \times GA)$
Cisterna magna	
Mean	$\text{Exp}(2.098095 - (239.0659 \times GA^{-2}) - 0.0000001547 \times GA^3)$
SD	$0.2297936 + (8.1872 \times GA^{-2})$

Ln, natural logarithm.

Table 8 Intra- and interobserver reproducibility for measurement of fetal brain structures

Structure	Mean difference (95% limits of agreement)		
	Intraobserver reproducibility	Interobserver reproducibility	Caliper replacement reproducibility
Parieto-occipital fissure (mm)	-0.02 (1.6)	0 (0.19)	-0.01 (0.19)
Sylvian fissure (mm)	-0.01 (0.21)	0 (0.22)	0 (0.28)
Anterior horn of lateral ventricle (mm)	-0.01 (0.18)	-0.02 (0.2)	0 (0.18)
Atrium of posterior horn of lateral ventricle (mm)	0 (0.11)	0 (0.18)	0.01 (0.17)
Cisterna magna (mm)	0 (0.16)	-0.02 (0.19)	0.01 (0.18)

10% of previous studies reported on maternal and fetal inclusion/exclusion criteria, pregnancy outcome or ultrasound quality control. Goodness of fit of the model to create the charts was reported in only 35% of the studies. Most importantly, no studies reported long-term infant outcomes, most probably owing to their retrospective descriptive design (30%); thus, data were often not collected specifically for the purpose of the study. Not surprisingly, these are some of the same challenges seen in previous studies to construct fetal biometry charts^{10,11}. Nevertheless, some previous studies did have a relatively low risk of methodological bias, and the ranges of our observed measurements did not differ substantially from their findings^{34,51–54}.

Strengths and limitations

A large number of sonographers were involved in this study; however, this more accurately reflects clinical practice⁵⁵. In addition, the quality of the images obtained in the study was of a high standard and in accordance with a predefined protocol²⁵. We set near-optimal conditions for scanning to minimize the potential contribution of confounding factors, which could also be seen as a strength. It is possible that measurements acquired on planes extracted from 3D volumes are not equivalent to measurements made from 2D image acquisition. Although volumetry is associated with a high degree of variability if not standardized⁵⁰, once rigorous methodology is adopted, 2D measurements from reconstructed planes can be as reproducible as measurements obtained in real time^{29,37}.

A key strength of our study is that we adopted a prescriptive design, as recommended by the WHO. We identified urban regions in which women were at low risk of perinatal complications; participants were then enrolled within these regions based on their individual characteristics. All ultrasound measurements were taken specifically for the purpose of constructing international standards with standardization of all study sites, using centrally trained staff and specially adapted ultrasound equipment to allow masking of measurements. For the offline analysis, we developed a novel quality-control strategy. The most appropriate statistical methods were used to analyze the dataset.

It could be argued that only longitudinal data should be used to assess fetal growth. However, given the design of FGLS, in which women mostly had an equal number of visits during pregnancy and these visits were according to what was prespecified in the protocol, cross-sectional data were acquired in order to ensure a representative number of brain-structure measurements per gestational week. The fitted model took this into account.

The INTERGROWTH-21st Project and WHO Multicentre Growth Reference Study have demonstrated previously the generalizability across geographically diverse international populations of anthropometric standards produced using the prescriptive approach^{12,31,56}. Follow-up of infants in the FGLS cohort has also been reported, and demonstrates strong similarities across sites

when assessed by variance components analysis and standardized site differences, showing that the sequence and timing of attainment of neurodevelopmental milestones and associated behaviors in early childhood are probably innate and universal³¹.

Conclusions

We report international standards for the size of five fetal brain structures throughout gestation. These standards use reproducible and highly controlled ultrasound measurements, and were created using a prospective cohort of fetuses that was followed up into childhood. Clinical use of such objective measurements may help to improve the screening and diagnostic performance of prenatal ultrasound. It should also allow a unified approach to fetal assessment by integrating with other standards from the same population and result in a common language when describing aberrations from expected norms^{57,58}. The proposed standards should not replace currently accepted cut-off values for triggering referral or further investigation; for example, we do not propose that we should redefine the diagnosis of antenatally diagnosed ventriculomegaly. This is because previous studies on the association between infant outcome and antenatally detected congenital brain abnormalities cannot simply be replicated^{57–59}.

ACKNOWLEDGMENTS

This project was supported by a generous grant from the Bill & Melinda Gates Foundation to the University of Oxford (Oxford, UK), for which we are very grateful. Aris T. Papageorgiou is supported by the National Institute for Health Research (NIHR) Oxford Biomedical Research Centre (BRC).

We thank the Health Authorities in Pelotas, Brazil; Beijing, China; Nagpur, India; Turin, Italy; Nairobi, Kenya; Muscat, Oman; Oxford, UK; and Seattle, USA, who facilitated the project by allowing participation of these study sites as collaborating centers. We are grateful to Philips Medical Systems, who provided the ultrasound equipment and technical assistance throughout the project. We thank MedSciNet U.K. Ltd for setting up the INTERGROWTH-21st website and for the development, maintenance and support of the online data management system.

Finally, we thank the parents and infants who participated in the studies and the more than 200 members of the research teams who made implementation of this project possible. The participating hospitals included: Brazil, Pelotas (Hospital Miguel Piltcher, Hospital São Francisco de Paula, Santa Casa de Misericórdia de Pelotas, and Hospital Escola da Universidade Federal de Pelotas); China, Beijing (Beijing Obstetrics & Gynecology Hospital, Shunyi Maternal & Child Health Centre, and Shunyi General Hospital); India, Nagpur (Ketkar Hospital, Avanti Institute of Cardiology Private Limited, Avantika Hospital, Gurukrupa Maternity Hospital,

Mulik Hospital & Research Centre, Nandlok Hospital, Om Women's Hospital, Renuka Hospital & Maternity Home, Saboo Hospital, Brajmonhan Taori Memorial Hospital, and Somani Nursing Home); Kenya, Nairobi (Aga Khan University Hospital, MP Shah Hospital and Avenue Hospital); Italy, Turin (Ospedale Infantile Regina Margherita Sant' Anna and Azienda Ospedaliera Ordine Mauriziano); Oman, Muscat (Khoula Hospital, Royal Hospital, Wattayah Obstetrics & Gynaecology Poly Clinic, Wattayah Health Centre, Ruwi Health Centre, Al-Ghoubra Health Centre and Al-Khuwair Health Centre); UK, Oxford (John Radcliffe Hospital) and USA, Seattle (University of Washington Hospital, Swedish Hospital, and Providence Everett Hospital).

Disclosure

A.T.P. and J.A.N. are Senior Advisors of Intelligent Ultrasound. All other authors declare no competing interests.


REFERENCES


- Salomon LJ, Alfirevic Z, Berghella V, Bilardo C, Hernandez-Andrade E, Johnsen SL, Kalache K, Leung KY, Malinger G, Munoz H, Prefumo F, Toi A, Lee W. Practice guidelines for performance of the routine mid-trimester fetal ultrasound scan. *Ultrasound Obstet Gynecol* 2011; 37: 116–126.
- International Society of Ultrasound in Obstetrics & Gynecology Education Committee. Sonographic examination of the fetal central nervous system: guidelines for performing the 'basic examination' and the 'fetal neurosonogram'. *Ultrasound Obstet Gynecol* 2007; 29: 109–116.
- Malinger G, Zakut H. The corpus callosum: normal fetal development as shown by transvaginal sonography. *AJR Am J Roentgenol* 1993; 161: 1041–1043.
- Paladini D, Volpe P. Posterior fossa and vermian morphometry in the characterization of fetal cerebellar abnormalities: a prospective three-dimensional ultrasound study. *Ultrasound Obstet Gynecol* 2006; 27: 482–489.
- Napolitano R, Thilaganathan B. Late termination of pregnancy and foetal reduction for foetal anomaly. *Best Pract Res Clin Obstet Gynaecol* 2010; 24: 529–537.
- Kuklisova-Murgasova M, Cifor A, Napolitano R, Papageorgiou A, Quaghebeur G, Rutherford MA, Hajnal JV, Noble JA, Schnabel JA. Registration of 3D fetal neurosonography and MRI. *Med Image Anal* 2013; 17: 1137–1150.
- Malinger G, Lerman-Sagie T, Waterberg N, Rotmensh S, Lev D, Glezerman M. A normal second-trimester ultrasound does not exclude intracranial structural pathology. *Ultrasound Obstet Gynecol* 2002; 20: 51–56.
- Namburete ALL, Stebbing RV, Kemp B, Yaqub M, Papageorgiou AT, Alison Noble J. Learning-based prediction of gestational age from ultrasound images of the fetal brain. *Med Image Anal* 2015; 21: 72–86.
- Donadono V, Napolitano R, Cavallaro A, Roberts NM, Papageorgiou AT. Charts of fetal brain structures: a systematic review of methodological quality. *Ultrasound Obstet Gynecol* 2017; 50: 59.
- Napolitano R, Dhami J, Ohuma EO, Ioannou C, Conde-Agudelo A, Kennedy SH, Villar J, Papageorgiou AT. Pregnancy dating by fetal crown–rump length: a systematic review of charts. *BJOG* 2014; 121: 556–565.
- Ioannou C, Talbot K, Ohuma E, Sarris I, Villar J, Conde-Agudelo A, Papageorgiou A. Systematic review of methodology used in ultrasound studies aimed at creating charts of fetal size. *BJOG* 2012; 119: 1425–1439.
- Villar J, Cheikh Ismail L, Staines-Urias E, Giuliani F, Ohuma EO, Victora CG, Papageorgiou AT, Altman DG, Garza C, Barros FC, Puglia F, Ochieng R *et al*; International Fetal and Newborn Growth Consortium for the 21st Century (INTERGROWTH-21st). The satisfactory growth and development at 2 years of age of the INTERGROWTH-21st Fetal Growth Standards cohort support its appropriateness for constructing international standards. *Am J Obstet Gynecol* 2018; 218: S841–S854.e2.
- Papageorgiou AT, Kennedy SH, Salomon LJ, Ohuma EO, Cheikh Ismail L, Barros FC, Lambert A, Carvalho M, Jaffer YA, Bertino E, Gravett MG, Altman DG *et al*; International Fetal and Newborn Growth Consortium for the 21st Century (INTERGROWTH-21st). International standards for early fetal size and pregnancy dating based on ultrasound measurement of crown–rump length in the first trimester of pregnancy. *Ultrasound Obstet Gynecol* 2014; 44: 641–648.
- Papageorgiou AT, Kemp B, Stones W, Ohuma EO, Kennedy SH, Purwar M, Salomon LJ, Altman DG, Noble JA, Bertino E, Gravett MG, Pang R *et al*; International Fetal and Newborn Growth Consortium for the 21st Century (INTERGROWTH-21st). Ultrasound-based gestational-age estimation in late pregnancy. *Ultrasound Obstet Gynecol* 2016; 48: 719–726.
- Papageorgiou AT, Ohuma EO, Altman DG, Todros T, Cheikh Ismail L, Lambert A, Jaffer YA, Bertino E, Gravett MG, Purwar M, Noble JA, Pang R *et al*; International Fetal and Newborn Growth Consortium for the 21st Century (INTERGROWTH-21st). International standards for fetal growth based on serial ultrasound measurements: the Fetal Growth Longitudinal Study of the INTERGROWTH-21st Project. *Lancet* 2014; 384: 869–879.
- Stirnemann J, Villar J, Salomon LJ, Ohuma E, Ruyan P, Altman DG, Nosten F, Craik R, Munim S, Cheikh Ismail L, Barros FC, Lambert A *et al*; International Fetal and Newborn Growth Consortium for the 21st Century (INTERGROWTH-21st). International estimated fetal weight standards of the INTERGROWTH-21st Project. *Ultrasound Obstet Gynecol* 2017; 49: 478–486.
- Papageorgiou AT, Ohuma EO, Gravett MG, Hirst J, da Silveira MF, Lambert A, Carvalho M, Jaffer YA, Altman DG, Noble JA, Bertino E, Purwar M *et al*; International Fetal and Newborn Growth Consortium for the 21st Century (INTERGROWTH-21st). International standards for symphysis–fundal height based on serial measurements from the Fetal Growth Longitudinal Study of the INTERGROWTH-21st Project: prospective cohort study in eight countries. *BMJ* 2016; 355: i5662.
- Cheikh Ismail L, Bishop DC, Pang R, Ohuma EO, Kac G, Abrams B, Rasmussen K, Barros FC, Hirst JE, Lambert A, Papageorgiou AT, Stones W *et al*. Gestational weight gain standards based on women enrolled in the Fetal Growth Longitudinal Study of the INTERGROWTH-21st Project: a prospective longitudinal cohort study. *BMJ* 2016; 352: i555.
- Villar J, Cheikh Ismail L, Victora CG, Ohuma EO, Bertino E, Altman DG, Lambert A, Papageorgiou AT, Carvalho M, Jaffer YA, Gravett MG, Purwar M *et al*; International Fetal and Newborn Growth Consortium for the 21st Century (INTERGROWTH-21st). International standards for newborn weight, length, and head circumference by gestational age and sex: the Newborn Cross-Sectional Study of the INTERGROWTH-21st Project. *Lancet* 2014; 384: 857–868.
- Villar J, Puglia FA, Fenton TR, Cheikh Ismail L, Staines-Urias E, Giuliani F, Ohuma EO, Victora CG, Sullivan P, Barros FC, Lambert A, Papageorgiou AT *et al*. Body composition at birth and its relationship with neonatal anthropometric ratios: the newborn body composition study of the INTERGROWTH-21st project. *Pediatr Res* 2017; 82: 305–316.
- Villar J, Giuliani F, Bhutta ZA, Bertino E, Ohuma EO, Ismail LC, Barros FC, Altman DG, Victora C, Noble JA, Gravett MG, Purwar M *et al*; International Fetal and Newborn Growth Consortium for the 21st Century (INTERGROWTH-21st). Postnatal growth standards for preterm infants: the Preterm Postnatal Follow-up Study of the INTERGROWTH-21(st) Project. *Lancet Glob Health* 2015; 3: e681–691.
- Villar J, Papageorgiou AT, Pang R, Ohuma EO, Cheikh Ismail L, Barros FC, Lambert A, Carvalho M, Jaffer YA, Bertino E, Gravett MG, Altman DG *et al*; International Fetal and Newborn Growth Consortium for the 21st Century (INTERGROWTH-21st). The likeness of fetal growth and newborn size across non-isolated populations in the INTERGROWTH-21st Project: the Fetal Growth Longitudinal Study and Newborn Cross-Sectional Study. *Lancet Diabetes Endocrinol* 2014; 2: 781–792.
- Villar J, Altman DG, Purwar M, Noble JA, Knight HE, Ruyan P, Cheikh Ismail L, Barros FC, Lambert A, Papageorgiou AT, Carvalho M, Jaffer YA *et al*. The objectives, design and implementation of the INTERGROWTH-21st Project. *BJOG* 2013; 120 (Suppl 2): 9–26.
- Wanyonyi SZ, Napolitano R, Ohuma EO, Salomon LJ, Papageorgiou AT. Image-scoring system for crown–rump length measurement. *Ultrasound Obstet Gynecol* 2014; 44: 649–654.
- Papageorgiou AT, Sarris I, Ioannou C, Todros T, Carvalho M, Pilu G, Salomon LJ. Ultrasound methodology used to construct the fetal growth standards in the INTERGROWTH-21st Project. *BJOG* 2013; 120 (Suppl 2): 27–32.
- Sarris I, Ioannou C, Ohuma E, Altman D, Hoch L, Cosgrove C, Fathima S, Salomon L, Papageorgiou A. Standardisation and quality control of ultrasound measurements taken in the INTERGROWTH-21st Project. *BJOG* 2013; 120 (Suppl 2): 33–37.
- WHO Multicentre Growth Reference Study Group. WHO Motor Development Study: windows of achievement for six gross motor development milestones. *Acta Paediatr Suppl* 2006; 450: 86–95.
- Altman DG, Ohuma EO. Statistical considerations for the development of prescriptive fetal and newborn growth standards in the INTERGROWTH-21st Project. *BJOG* 2013; 120 (Suppl 2): 71–76.
- Sarris I, Ioannou C, Ohuma E, Altman D, Hoch L, Cosgrove C, Fathima S, Salomon L, Papageorgiou A. Fetal biometry: how well can offline measurements from three-dimensional volumes substitute real-time two-dimensional measurements? *Ultrasound Obstet Gynecol* 2013; 42: 560–570.
- Ioannou C, Sarris I, Napolitano R, Ohuma E, Javadi MK, Papageorgiou AT. A longitudinal study of normal fetal femur volume. *Prenat Diagn* 2013; 33: 1088–1094.
- Villar J, Fernandes M, Purwar M, Staines-Urias E, Di Nicola P, Cheikh Ismail L, Ochieng R, Barros F, Albernaz E, Victora C, Kunnawar N, Temple S *et al*. Neurodevelopmental milestones and associated behaviours are similar among healthy children across diverse geographical locations. *Nat Commun* 2019; 10: 511.
- Cavallaro A, Ash ST, Napolitano R, Wanyonyi S, Ohuma EO, Molloholli M, Sande J, Sarris I, Ioannou C, Norris T, Donadono V, Carvalho M *et al*. Quality control of ultrasound for fetal biometry: results from the INTERGROWTH-21st Project. *Ultrasound Obstet Gynecol* 2018; 52: 332–339.
- Napolitano R, Donadono V, Ohuma EO, Knight CL, Wanyonyi SZ, Kemp B, Norris T, Papageorgiou AT. Scientific basis for standardization of fetal head measurements by ultrasound: a reproducibility study. *Ultrasound Obstet Gynecol* 2016; 48: 80–85.
- Alonso I, Borenstein M, Grant G, Narbona I, Azumendi G. Depth of brain fissures in normal fetuses by prenatal ultrasound between 19 and 30 weeks of gestation. *Ultrasound Obstet Gynecol* 2010; 36: 693–699.
- Alagappan R, Browning PD, Laorr A, McGahan JP. Distal lateral ventricular atrium: reevaluation of normal range. *Radiology* 1994; 193: 405–408.

36. Araujo Junior E, Martins WP, Nardoza LM, Pires CR, Filho SM. Reference range of fetal transverse cerebellar diameter between 18 and 24 weeks of pregnancy in a Brazilian population. *J Child Neurol* 2015; 30: 250–253.
37. Chang CH, Chang FM, Yu CH, Ko HC, Chen HY. Three-dimensional ultrasound in the assessment of fetal cerebellar transverse and antero-posterior diameters. *Ultrasound Med Biol* 2000; 26: 175–182.
38. Chavez MR, Ananth CV, Smulian JC, Lashley S, Kontopoulos EV, Vintzileos AM. Fetal transverse cerebellar diameter nomogram in singleton gestations with special emphasis in the third trimester: a comparison with previously published nomograms. *Am J Obstet Gynecol* 2003; 189: 1021–1025.
39. Hata K, Hata T, Senoh D, Makihara K, Aoki S, Takamiya O, Kitao M. Ultrasonographic measurement of the fetal transverse cerebellum in utero. *Gynecol Obstet Invest* 1989; 28: 111–112.
40. Joshi BR. Fetal transverse cerebellar diameter nomogram in Nepalese population. *J Inst Med* 2010; 32: 19–23.
41. Koktencer A, Dilmen G, Kurt A. The cisterna magna size in normal second-trimester fetuses. *J Perinat Med* 2007; 35: 217–219.
42. Lei H, Wen SW. Ultrasonographic examination of intrauterine growth for multiple fetal dimensions in a Chinese population. Central–South China Fetal Growth Study Group. *Am J Obstet Gynecol* 1998; 178: 916–921.
43. Mahony BS, Callen PW, Filly RA, Hoddick WK. The fetal cisterna magna. *Radiology* 1984; 153: 773–776.
44. Uerpaiojkit B, Charoenvidhya D, Manotaya S, Tanawattanachareon S, Wacharaprechanont T, Tannirandom Y. Fetal transverse cerebellar diameter in Thai population. *J Med Assoc Thai* 2001; 84 (Suppl 1): S346–351.
45. Pistorius LR, Stoutenbeek P, Groenendaal F, de Vries L, Manten G, Mulder E, Visser G. Grade and symmetry of normal fetal cortical development: a longitudinal two- and three-dimensional ultrasound study. *Ultrasound Obstet Gynecol* 2010; 36: 700–708.
46. Timor-Tritsch IE, Monteagudo A. Transvaginal fetal neurosonography: standardization of the planes and sections by anatomic landmarks. *Ultrasound Obstet Gynecol* 1996; 8: 42–47.
47. Sarris I, Ioannou C, Dighe M, Mitidieri A, Oberio M, Qingqing W, Shah J, Sohoni S, Al Zidjaji W, Hoch L, Altman DG, Papageorgiou AT; International Fetal and Newborn Growth Consortium for the 21st Century (INTERGROWTH-21st). Standardization of fetal ultrasound biometry measurements: improving the quality and consistency of measurements. *Ultrasound Obstet Gynecol* 2011; 38: 681–687.
48. Bornstein E, Monteagudo A, Santos R, Strock I, Tsybal T, Lenchner E, Timor-Tritsch IE. Basic as well as detailed neurosonograms can be performed by offline analysis of three-dimensional fetal brain volumes. *Ultrasound Obstet Gynecol* 2010; 36: 20–25.
49. Quarello E, Stirnemann J, Ville Y, Guibaud L. Assessment of fetal Sylvian fissure operculization between 22 and 32 weeks: a subjective approach. *Ultrasound Obstet Gynecol* 2008; 32: 44–49.
50. Ioannou C, Sarris I, Salomon LJ, Papageorgiou AT. A review of fetal volumetry: the need for standardization and definitions in measurement methodology. *Ultrasound Obstet Gynecol* 2011; 38: 613–619.
51. Almog B, Gamzu R, Achiron R, Fainaru O, Zalel Y. Fetal lateral ventricular width: what should be its upper limit? A prospective cohort study and reanalysis of the current and previous data. *J Ultrasound Med* 2003; 22: 39–43.
52. Alves CM, Araujo Junior E, Nardoza LM, Goldman SM, Martinez LH, Martins WP, Oliveira PS, Moron AF. Reference ranges for fetal brain fissure development on 3-dimensional sonography in the multiplanar mode. *J Ultrasound Med* 2013; 32: 269–277.
53. Snijders RJ, Nicolaides KH. Fetal biometry at 14–40 weeks' gestation. *Ultrasound Obstet Gynecol* 1994; 4: 34–48.
54. Verburg BO, Steegers EA, De Ridder M, Snijders RJ, Smith E, Hofman A, Moll HA, Jaddoe VW, Witteman JC. New charts for ultrasound dating of pregnancy and assessment of fetal growth: longitudinal data from a population-based cohort study. *Ultrasound Obstet Gynecol* 2008; 31: 388–396.
55. Sarris I, Ioannou C, Chamberlain P, Ohuma E, Roseman F, Hoch L, Altman DG, Papageorgiou AT; International Fetal and Newborn Growth Consortium for the 21st Century (INTERGROWTH-21st). Intra- and interobserver variability in fetal ultrasound measurements. *Ultrasound Obstet Gynecol* 2012; 39: 266–273.
56. de Onis M, Garza C, Onyango AW, Martorell R. WHO Child Growth Standards. *Acta Paediatr Suppl* 2006; 450: 1–101.
57. D'Antonio F, Khalil A, Garel C, Pilu G, Rizzo G, Lerman-Sagie T, Bhide A, Thilaganathan B, Manzoli L, Papageorgiou AT. Systematic review and meta-analysis of isolated posterior fossa malformations on prenatal ultrasound imaging (part 1): nomenclature, diagnostic accuracy and associated anomalies. *Ultrasound Obstet Gynecol* 2016; 47: 690–697.
58. Melchiorre K, Bhide A, Gika AD, Pilu G, Papageorgiou AT. Counseling in isolated mild fetal ventriculomegaly. *Ultrasound Obstet Gynecol* 2009; 34: 212–224.
59. D'Antonio F, Khalil A, Garel C, Pilu G, Rizzo G, Lerman-Sagie T, Bhide A, Thilaganathan B, Manzoli L, Papageorgiou AT. Systematic review and meta-analysis of isolated posterior fossa malformations on prenatal imaging (part 2): neurodevelopmental outcome. *Ultrasound Obstet Gynecol* 2016; 48: 28–37.

SUPPORTING INFORMATION ON THE INTERNET

The following supporting information may be found in the online version of this article:

 **Videoclip S1** Demonstration of methodology for volume manipulation and caliper placement for measurement acquisition of fetal brain structures using MITK software.

 **Figure S1** Bland–Altman plots showing intra- (a) and inter- (b) observer reproducibility for volume manipulation and caliper placement for measurement acquisition of fetal brain structures, and interobserver reproducibility for caliper replacement on stored images (c).

Appendix S1 Detailed methodology for 3D volume manipulation and caliper placement

Table S1 Characteristics of all pregnancies in Fetal Growth Longitudinal Study (FGLS) and those included in current study

Table S2 Morbidity at 2 years of age in all pregnancies in Fetal Growth Longitudinal Study (FGLS) and those included in current study