Table SDC 7: Concordance and discordance of empiric regimen by pathogen with no fatalities from overall cohort (n=452 overall).

Pathogen			Survivors
Salmonella spp. (NTS & Typhi)	Empiric treatment	Concordant	38 (95.0%)
		Discordant	2 (5.0%)
	Total		40 (100%)
Streptococcus pneumoniae	Empiric treatment	Concordant	33 (91.7%)
		Discordant	3 (8.3%)
	Total		36 (100%)
Streptococcus agalactiae	Empiric treatment	Concordant	20 (87.0%)
		Discordant	3 (13.0%)
	Tota	ıl	23 (100%)
Enterococcus faecalis	Empiric treatment	Concordant	15 (75.0%)
		Discordant	5 (25.0%)
	Tota	ıl	20 (100%)
Enterococcus faecium	Empiric treatment	Concordant	3 (30.0%)
		Discordant	7 (70.0%)
	Tota	ıl .	10 (100%)
Enterococcus spp.	Empiric treatment	Concordant	2 (40.0%)
		Discordant	3 (60.0%)
	Total		5 (100%)
Haemophilus influenzae	Empiric treatment	Concordant	2 (66.7%)
		Discordant	1 (33.3%)
	Tota	ıl	3 (100%)
Burkholderia cepacia	Empiric treatment	Concordant	0 (0%)
		Discordant	2 (100%)
	Total		2 (100%)
Proteus spp.	Empiric treatment	Concordant	2 (100%)
		Discordant	0 (0%)
	Tota	ıl	2 (100%)
Total			141