



MRSA surveillance programmes worldwide: moving towards a harmonised international approach



Valérie O. Baede¹, Michael Z. David², Arjana Tambic Andrasevic³, Dominique S. Blanc^{4,5}, Michael Borg⁶, Grainne Brennan⁷, Boudewijn Catry^{8,9}, Aurélie Chabaud¹⁰, Joanna Empel¹¹, Hege Enger¹², Marie Hallin¹³, Marina Ivanova¹⁴, Andreas Kronenberg¹⁵, Kuntaman Kuntaman^{16,17}, Anders Rhod Larsen¹⁸, Katrien Latour⁸, Jodi A. Lindsay¹⁹, Bruno Pichon²⁰, Dewi Santosaningsih²¹, Leo M. Schouls²², François Vandenesch^{23,24}, Guido Werner²⁵, Dorota Żabicka¹¹, Helena Žemličková^{26,27}, Harald Seifert²⁸, Margreet C. Vos^{1,*}, for the MRSA Surveillance Worldwide Study Group (ISAC), the ESCMID Study Group for Nosocomial Infections (ESGNI), the ESCMID Study Group for Staphylococci and Staphylococcal Diseases (ESGS)

¹ Department of Medical Microbiology and Infectious Diseases, Erasmus MC University Medical Center Rotterdam, Rotterdam, the Netherlands

² Department of Medicine, University of Pennsylvania, Philadelphia, Pennsylvania, USA

³ Department of Clinical Microbiology, University Hospital for Infectious Diseases, University of Zagreb School of Dental Medicine, Zagreb, Croatia

⁴ Hospital Preventive Medicine, Lausanne University Hospital and University of Lausanne, Lausanne, Switzerland

⁵ Swiss National Reference Center for Emerging Antibiotic Resistance, Lausanne site, Lausanne, Switzerland

⁶ Department of Infection Control, Mater Dei Hospital, Msida, Malta

⁷ National MRSA Reference Laboratory, St James's Hospital, Dublin 8, Ireland

⁸ Healthcare-associated infections & antimicrobial resistance, Sciensano, Brussels, Belgium

⁹ Faculty of Medicine, Université libre de Bruxelles, Brussels, Belgium

¹⁰ Mission nationale SPARES (Surveillance et Prévention de l'AntibioRésistance en Etablissement de santé), Limoges University Hospital, Limoges, France

¹¹ Department of Epidemiology and Clinical Microbiology, National Medicines Institute, Warsaw, Poland

¹² The Norwegian MRSA Reference Laboratory, St Olavs Hospital, Trondheim, Norway

¹³ Department of Microbiology, LHUB-ULB, Université Libre de Bruxelles, Brussels, Belgium

¹⁴ East-Tallinn Central Hospital Laboratory, Tallinn, Estonia

¹⁵ Swiss Centre for Antibiotic Resistance, Institute for Infectious Diseases, University of Bern, Bern, Switzerland

¹⁶ Department of Medical Microbiology, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia

¹⁷ Dr Soetomo General Academic Hospital, Surabaya, Indonesia

¹⁸ National Reference Laboratory for Antimicrobial Resistance, Department of Bacteria, Fungi and Parasites, Statens Serum Institut, Copenhagen, Denmark

¹⁹ Institute for Infection and Immunity, St George's, University of London, London, UK

²⁰ National Infection Service, Public Health England, London, UK

²¹ Department of Clinical Microbiology, Faculty of Medicine, Brawijaya University/Dr Saiful Anwar Hospital, Malang, Indonesia

²² Centre for Infectious Disease Control, National Institute for Public Health and the Environment (RIVM), Bilthoven, the Netherlands

²³ Centre International de Recherche en Infectiologie (CIRI), Université de Lyon, INSERM, U1111, Université Claude Bernard Lyon 1, CNRS, UMR5308, ENS de Lyon, Lyon, France

²⁴ Centre National de Référence des Staphylocoques, Institut des Agents Infectieux, Hospices Civils de Lyon, Lyon, France

²⁵ Robert Koch Institute, Department of Infectious Diseases, National Reference Centre for Staphylococci and Enterococci, Wernigerode Branch, Germany

²⁶ National Reference Laboratory for Antibiotics, Centre for Epidemiology and Microbiology, National Institute of Public Health, 10000 Prague, Czech Republic

²⁷ Department of Microbiology, 3rd Faculty of Medicine Charles University, University Hospital Kralovske Vinohrady and National Institute of Public Health, 10000 Prague, Czech Republic

²⁸ Institute for Medical Microbiology, Immunology and Hygiene, University of Cologne, Köln, Germany

ARTICLE INFO

Article history:

Received 23 July 2021

Accepted 19 January 2022

Editor: Stefania Stefani

ABSTRACT

Multinational surveillance programmes for methicillin-resistant *Staphylococcus aureus* (MRSA) are dependent on national structures for data collection. This study aimed to capture the diversity of national MRSA surveillance programmes and to propose a framework for harmonisation of MRSA surveillance. The International Society of Antimicrobial Chemotherapy (ISAC) MRSA Working Group conducted a structured

* Corresponding author. Mailing address: Department of Medical Microbiology and Infectious Diseases, Erasmus MC University Medical Centre Rotterdam, P.O. Box 2040, 3000 CA Rotterdam, the Netherlands.

E-mail address: m.vos@erasmusmc.nl (M.C. Vos).

Keywords:

Antimicrobial resistance
Staphylococcus aureus
Monitoring
Epidemiology

survey on MRSA surveillance programmes and organised a webinar to discuss the programmes' strengths and challenges as well as guidelines for harmonisation. Completed surveys represented 24 MRSA surveillance programmes in 16 countries. Several countries reported separate epidemiological and microbiological surveillance. Informing clinicians and national policy-makers were the most common purposes of surveillance. Surveillance of bloodstream infections (BSIs) was present in all programmes. Other invasive infections were often included. Three countries reported active surveillance of MRSA carriage. Methodology and reporting of antimicrobial susceptibility, virulence factors, molecular genotyping and epidemiological metadata varied greatly. Current MRSA surveillance programmes rely upon heterogeneous data collection systems, which hampers international epidemiological monitoring and research. To harmonise MRSA surveillance, we suggest improving the integration of microbiological and epidemiological data, implementation of central biobanks for MRSA isolate collection, and inclusion of a representative sample of skin and soft-tissue infection cases in addition to all BSI cases.

Crown Copyright © 2022 Published by Elsevier Ltd.

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

1. Introduction

Antimicrobial resistance is currently one of the greatest threats to public health. Methicillin-resistant *Staphylococcus aureus* (MRSA) is the second most common cause of antibiotic-resistant bacterial infections in the European Union (EU) and European Economic Area (EEA) [1]. Many MRSA originate from a limited number of historically dominant clonal lineages [2]. While some MRSA clones are found worldwide, others are restricted to certain geographic areas, implying differences in transmission [3]. To analyse MRSA transmission and to decrease the incidence of new infections, international epidemiological research is crucial, and this research depends on MRSA surveillance programmes.

Many MRSA surveillance programmes exist worldwide, but only a few are multinational [4]. One European multinational programme is the European Antimicrobial Resistance Surveillance Network (EARS-Net) [5]. EARS-Net is co-ordinated by the European Centre for Disease Prevention and Control (ECDC) and depends on national surveillance systems. While susceptibility testing and interpretation recommendations have been harmonised [European Committee on Antimicrobial Susceptibility Testing (EUCAST)] [6], national surveillance programmes use different sampling strategies and laboratory techniques that can bias analyses [5]. Also, non-European multinational MRSA surveillance programmes mostly depend on national networks using different methodologies. Examples are the Asian Network for Surveillance of Resistant Pathogens (ANSORP), the Latin American Network for Antimicrobial Resistance Surveillance (ReLAVRA), the SENTRY Antimicrobial Surveillance Program and the Tigecycline Evaluation and Surveillance Trial (T.E.S.T.), now embedded in the Antimicrobial Testing Leadership and Surveillance (ATLAS) database [7–11].

Heterogeneity in testing and sampling practices hampers international epidemiological surveillance and the establishment of an early warning system for emerging MRSA clones [4,12,13]. Additionally, it lowers the quality of available data. This can be illustrated by the experiences of the MACOTRA Study Group, which aimed to establish an MRSA strain collection to analyse transmission success of MRSA. However, drafted definitions of successful versus unsuccessful MRSA strains were not applicable due to the heterogeneity described above. As a result, multiple strategies for strain selection were adopted, leading to selection bias and decreased data comparability. This demonstrates that the current organisation of MRSA surveillance systems and reference laboratories is not sufficient to support a greater understanding of MRSA transmission, nor to detect emerging virulent strains.

The aim of this project was to capture the diversity of existing national and institutional MRSA surveillance programmes and to propose a framework for a standardised (inter)national surveil-

lance network. A structured survey on current MRSA surveillance practices was conducted, followed by a webinar organised by the International Society of Antimicrobial Chemotherapy (ISAC) MRSA Working Group.

2. Methods

ISAC MRSA Working Group members were contacted to identify directors or head microbiologists of national or regional MRSA surveillance programmes or staphylococcal reference laboratories in their respective countries. Other representatives of national organisations participating in EARS-Net were contacted directly [5]. All representatives were invited to participate in a structured survey drafted by the executive committee of the ISAC MRSA Working Group [MCV (chair), MZD, HS, VB and SS]. The survey contained sections regarding organisational structure, surveillance goals, strain and sample characteristics, epidemiological metadata and laboratory reports. An overview of the survey is given in the Supplementary data.

Additionally, surveillance programme representatives were invited to participate in a webinar, held on 10 March 2021, organised by the ISAC MRSA Working Group and the MACOTRA Study Group, which was entitled 'Regional and National MRSA Surveillance Programs Worldwide: Results of a Survey and Discussion of Current Practices'. Its purpose was to present an overview of surveillance programmes to an international audience, to discuss these programmes' strengths and challenges, and to discuss the requirements for harmonisation of MRSA surveillance.

3. Results

Representatives of 12 MRSA surveillance programmes in nine countries were invited through the ISAC MRSA Working Group (Fig. 1). Another 21 national organisations participating in EARS-Net were also invited. In total, 18 surveys were completed between January and April 2021, representing 24 MRSA surveillance programmes in 14 European and 2 non-European countries. Multiple surveillance programmes were described for Belgium (3), Germany (3), France (2), Indonesia (2), Switzerland (2) and the USA (2). Fourteen surveillance programmes in eight countries were presented at the webinar.

3.1. Survey

A summary of survey results is given in Table 1.

3.1.1. Surveillance structure and purpose

All countries conducted surveillance at the national level, except Malta. In Malta, surveillance was performed at the sole ter-

Table 1
Survey results

	BE	CH-1	CH-2	CZ	DE	DK	EE	FR	GB	HR	ID-1	ID-2	IE	MT	NL	NO	PL	US	
Surveillance structure																			
MRSA surveillance standardised	x		x		x	x	x	x	x	x				x	x	x		x	
MRSA surveillance on national level	x		x	x	x	x	x	x	x	x	x	x	x	999	x	x	x	x	
MRSA surveillance on regional level		x	x		x	x	U					U		999	x			x	
MRSA surveillance on local/hospital level	x	x	x	x	x	x	x		x	x	x	x	x	x	x		U	x	
General community included			x			x		x	x	x		x	x		x	x		x	
Outpatient clinics included			x			x	x	x		x			x	x	x				
Mandatory for specific communities	x					x													
Mandatory for specific anatomic sites or infections (e.g. BSI)	x			x	x	x	x		x	x	x		x			x		x	
Results reported in an annual report, scientific publications, website etc.	x		x	x	x	x	x	x	x	x	x		x	x	x	x	x	x	
Organisational structure																			
SRL is a governmental organisation				x	x	x	x	x	x	x	x				x	x	x	x	
No extra costs for genotyping	x			x	x	x		x	x	x			x	999	x	x	x	x	
No extra costs for other tests	x			x	x	x		x	x	x			x	999	NA	x	x	x	
Mandatory submission of strains						x				x						x			
Purpose of surveillance																			
ECDC data collection	x			x	x	x	x	x	U	x			x	x	x			x	
National epidemiology	x		x		x	x	x	x	x	x	x			x	x			x	
Clinical question	x	x	x	x	x	x	x	x	x	x	x		x	x	x	x	x	x	
Research question, e.g. virulence factors	x	x		x	x	x		x	x	x		x	x		x	x	x		
Research question, e.g. molecular typing	x	x		x	x	x		x	x			x	x			x	x		
Other research questions					x	x	x		x			x				x	x		
Sample data																			
MRSA collection in biobank for research	x	x		x	x	x		x	x	x		x	x	x	x	x	x	x	
Infection isolates included	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	
Carriage isolates included	x	x	x		x	x	x		x	x		x		x	x	x	x		
Case/sample inclusion restricted per period	x	x		x	x				Other	x				x	x	x		x	
Case/sample inclusion restricted per individual							U		Other	x	x			x	x	x		x	
Amount of MRSA isolates collected each year	100–1000	100–1000	NA	100–1000	1000–10000	1000–10000	NA	1000–10000	1000–10000	0–100	NA	NA	100–1000	100–1000	1000–10000	1000–10000	100–1000	100–1000	
Sample types																			
Blood	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	
Wound	x	x	x	x	x	x		x	x		x	x	x	x	x	x	x		
Skin	x	x	x	x	x	x		x	x				x	x	x	x	x		
Nose/throat/perineum	x	x	x	x	x	x			x			x	x		x	x	x		
Other	x	x	x		x	x		x			x		x		x	x		x	
Strain data																			
Default genotyping done	x	x		x	x	x		x	x			x	999		x		x	x	
WGS	x	x			x	x		x	x						x	x	x	x	
MLST	x			x											x	x	x	x	
spa	x			x	x	x							x			x	x	x	
PFGE	x	x																x	

(continued on next page)

Table 1 (continued)

	BE	CH-1	CH-2	CZ	DE	DK	EE	FR	GB	HR	ID-1	ID-2	IE	MT	NL	NO	PL	US
Other genotyping technique	x	x				x							x		x		x	x
Antimicrobial susceptibility tested	x		x	x	x	x	x	x	Other	x	x	x	x	999		x	x	x
Virulence factors tested	x	x			x	x		x	x				x	999	x	x	x	x
Recorded epidemiological metadata																		
Age			x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
Sex			x	x	x	x	x		x	x	x	x	x	x	x	x	x	x
Date of sampling	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
Place of residence			x	x		x	x		x	x	x	x	x	x	x	x	x	x
MRSA acquisition risk group					x	x	999					x			x	x		x
Medical specialty			x	x	x	x	999	x	x	x	x	x	x	x		x		
Patient admission to ICU or general ward			x	x	x	x	999	x		x	x	x	x	x	x	x	x	x
Admission history in same HCC			x			x	999		x			x		NA				x
Admission history in other HCCs						x	999					x		NA				x
Outbreak metadata					x	x	999	x								x		
Recorded metadata enables outbreak traceability				U		x	999							x				
Laboratory reports																		
Laboratory reports returned to submitter	x	x	NA	x	x	x	x	x	x	x	NA	U	x	NA	x	x	x	
Turnaround time for genotyping results	7d	1m	NA	3w	3w	2–7d		2w	2w		NA	U	10d	NA	2d	5d	1m	
Turnaround time for virulence factor results	7d	1m	NA	1w	1w	2–7d		2w	2w		NA	U	5d	NA	2d	5d	1w	
<i>Data on laboratory reports include:</i>																		
Genotyping result in relation to strains in the HCC	x	x			x		999		x		NA		x	NA	x	x		
Genotyping result in relation to strains in the region		x					999				NA			NA	x	x		
Genotyping result in relation to strains in the country	x	x					999	x			NA			NA	x	x		
Admission history of patients with the same genotype in the HCC							999		x		NA			NA				
Admission history of patients with the same genotype in the region							999		x		NA			NA				
Admission history of patients with the same genotype in the country							999				NA			NA				
Place of residence of patients with the same genotype							999		x		NA			NA	x			
A combination of all mentioned above, but time period restricted							999	x			NA			NA	x			

Abbreviations: MRSA, methicillin-resistant *Staphylococcus aureus*; BSI, bloodstream infection; ECDC, European Centre for Disease Prevention and Control; WGS, whole-genome sequencing; MLST, multilocus sequence typing; PFGE, pulsed-field gel electrophoresis; ICU, intensive care unit; HCC, healthcare centre; d, days; m, months; w, weeks.

Country abbreviations: BE, Belgium; CH, Switzerland; CZ, Czech Republic; DE, Germany; DK, Denmark; EE, Estonia; FR, France; GB, United Kingdom; HR, Croatia; ID, Indonesia; IE, Ireland; MT, Malta; NL, Netherlands; NO, Norway; PL, Poland; US, United States of America.

NOTE: For Belgium, France, Germany and the USA, results mentioned in this table described multiple surveillance programmes. Details are available in the Supplementary data. ID-1: this survey described MRSA surveillance at Dr Saiful Anwar Hospital in Malang, Java, Indonesia. ID-2: this survey described MRSA surveillance at Dr Soetomo Hospital, Surabaya, Java, Indonesia. CH-1: this survey described a regional MRSA surveillance programme in the French part of Switzerland, co-ordinated by Lausanne University Hospital. CH-2: this survey described the national antimicrobial resistance surveillance programme ANRESIS, Switzerland.

NOTE: All positive answers are depicted by an x, and blanks are negative answers. U, unknown; Other, other answer was applicable; NA, not applicable; 999, missing data.




Fig. 1. Overview of participating methicillin-resistant *Staphylococcus aureus* (MRSA) surveillance programmes. Representatives of MRSA surveillance programmes were identified through the network of the International Society of Antimicrobial Chemotherapy MRSA Working Group (ISAC MRSA-WG) or through participation in the European Antimicrobial Resistance Surveillance Network (EARS-Net). Listed are the number of contacted organisations and respective number of countries. Also listed are the number of returned surveys and presentations given at the webinar, for the respective number of included countries and surveillance programmes.

tiary hospital but covered >90% of all national testing. In four countries, surveillance was primarily conducted at the hospital level and was organised around the surveillance of bloodstream infections (BSIs). In the Czech Republic, all hospitals performed some MRSA surveillance, and MRSA BSI surveillance captured ~80% of the population. In Ireland and Poland, passive surveillance was performed through EARS-Net participation, and several national structured surveys were conducted in the past 20 years. For Indonesia, active MRSA surveillance was performed in several hospitals, but most surveillance was conducted for research purposes.

In Belgium, France and Germany, multiple separate programmes for epidemiological and microbiological surveillance were reported. In Switzerland, a local initiative focused on molecular surveillance of MRSA exists in addition to the national surveillance system (ANRESIS), which gathers epidemiological data for all antimicrobial-resistant micro-organisms. In the USA, at least two large MRSA surveillance programmes exist: a national programme on MRSA BSI in which most hospitals participate; and a population-based programme of invasive MRSA infections covering ~5% of the population [14].

Most surveillance programmes served multiple goals. The most common purpose of surveillance was to inform clinicians, public-health workers and laboratories about current resistance trends (17/18). Other epidemiological goals were informing national policy-makers (14/18) or EARS-Net participation (for all current EU/EEA countries except Norway). Research goals included studies on staphylococcal virulence factors (12/18), resistance profiles, specific clones such as livestock-associated MRSA (LA-MRSA), risk factor analysis, monitoring the effectiveness of interventions, or outbreak investigations.

3.1.2. Collection of isolates and microbiological and epidemiological data

Results of BSI isolates were collected in all surveillance programmes. Collection of wound (15/18), skin (12/18) and nose, throat or perineum (12/18) isolates also occurred frequently. Eleven programmes reported the inclusion of isolates from other clinical sample types such as cerebrospinal fluid, urine, pus, sputum or all clinical samples (6/11). Active surveillance of MRSA carriage was reported only for Denmark, the Netherlands and Norway. Isolates from outpatients (9/18) and the general community (10/18) were also reported, but systematic active surveillance of these groups was performed only in Denmark, the Netherlands and Norway. Long-term storage of isolates varied, ranging from BSI isolates only

to all submitted isolates. Programmes with an epidemiological focus often lacked routine isolate collection.

Most programmes collected microbiological data such as antimicrobial susceptibilities (14/18) and the presence of virulence factors (11/18). The presence of the Pantón–Valentine leukocidin (PVL) toxin was most commonly tested (8/11). Eleven programmes performed genotyping on all isolates, with *spa* typing as the most common method (6/11). A wide range of genotyping techniques were reported: whole-genome sequencing (WGS) (10/11); *spa* typing (8/11); multilocus sequence typing (MLST) (6/11); pulsed-field gel electrophoresis (PFGE) (3/11); *agr* group typing (Belgium); clonal complex 398 (CC398) subtyping (Denmark); multiple-locus variable number tandem repeat analysis (MLVA) (the Netherlands); multiple-locus variable number tandem repeat fingerprinting (MLVF) (Poland); DNA microarray (Ireland); staphylococcal cassette chromosome *mec* (SCC*mec*) typing (USA); CC8 subtyping (USA); and double-locus sequence typing (DLST) (local Swiss initiative).

Regarding epidemiological metadata, demographic variables were most commonly collected (16/18), followed by clinical information (14/18), MRSA risk factors (6/18) and outbreak metadata (4/18).

3.2. Webinar

The goals, strengths, challenges and future plans of ten MRSA surveillance programmes in eight countries were presented at the ISAC MRSA webinar. Strengths were the robust network of local laboratories and/or hospitals in the Czech Republic, France and Poland, as well as the national surveillance programmes in Belgium, Denmark, Germany, the Netherlands and Switzerland. In Denmark and the Netherlands, strong collaboration between epidemiological and microbiological departments and existing WGS pipelines enhanced MRSA surveillance. However, limited collaboration between epidemiological and microbiological surveillance structures posed a major challenge for Belgium, France, Germany and Switzerland. Representatives of the Czech Republic, Denmark, Germany, the Netherlands, Poland and Switzerland advocated for the implementation of WGS as a default genotyping technique and an accompanying platform to share WGS data. For many surveillance programmes, stability of financial support was a concern.

Based on our results and webinar discussions, the ISAC MRSA Working Group, MRSA Surveillance Worldwide Study Group and the MACOTRA Study Group propose three suggestions to harmonise MRSA surveillance:




Fig. 2. Proposal for harmonised methicillin-resistant *Staphylococcus aureus* (MRSA) surveillance. To harmonise surveillance, we propose (1) the inclusion of all bloodstream infection (BSI) isolates and a representative sample of skin and soft-tissue infection (SSTI) isolates in proportion to MRSA prevalence, (2) integration of microbiological and epidemiological data in a single database using standardised report templates and (3) implementation of central biobanks for collection and further characterisation of MRSA isolates. Orange flags depict the main challenges in harmonised surveillance. WGS, whole-genome sequencing.

- inclusion of all BSI cases and a representative number of skin and soft-tissue infection (SSTI) cases in proportion to MRSA prevalence;
- integration of microbiological and epidemiological data; and
- implementation of central biobanks at the national level for the collection and further characterisation of MRSA strains using common nomenclature allowing international comparisons.

The challenges and our proposal for harmonised surveillance are summarised in Fig. 2.

4. Discussion

Our study presents an overview of existing MRSA surveillance programmes in various parts of the world with an emphasis on European countries. It demonstrates the great diversity of MRSA surveillance programmes, both in surveillance structure as well as in microbiological and epidemiological data collection. Factors potentially driving this diversity are the primary goals of surveillance, the population size, MRSA prevalence and laboratory capacity. To improve the work of these systems, a harmonised approach for surveillance programmes is needed. We propose the inclusion of SSTI cases in addition to all BSI cases. BSI cases represent the most life-threatening MRSA infections. Because these cases are clearly defined, they provide high-quality data for surveillance. Most surveillance programmes already include BSI cases.

MRSA BSIs are predominantly endogenous infections, preceded by carriage and/or non-invasive infections [15,16]. For this reason, it is desirable to include non-BSI cases in surveillance as well. SSTIs represent the majority of *S. aureus* infections and are often acquired in the community. Inclusion of SSTIs in surveillance likely increases the probability of detecting emerging clones, which may also have significant public-health impact. We recommend including a representative number of SSTI cases in proportion to BSI

cases and MRSA prevalence to limit selection bias. This proportion will depend on the number of estimated MRSA BSI cases within the country, considering the expected volume and thus feasibility. A clear definition of SSTI such as presented in the CDC/NHSN Patient Safety Component Manual must be used to prevent misclassification [17].

Integration of microbiological and epidemiological data should be improved to enhance data quality [4,12]. Completion of a standardised epidemiological metadata report for each submitted case is essential. In addition to demographic data (i.e. age, sex and place of residence), the sampling date and site and classification of the isolate as being from infection or colonisation are necessary. Also required is information on relevant risk factors for MRSA acquisition to assign the patient/carrier to a defined risk group or to identify new risk factors.

Implementation of a central MRSA biobank at the national level is needed to collect isolates corresponding to the obtained epidemiological data. Typically, this biobank would be maintained by a reference laboratory, which can provide genotyping, antimicrobial susceptibility testing, and testing for virulence genes on a well-defined sample of isolates. We advocate for the use of WGS as the routine genotyping technique along with common nomenclature allowing international comparisons, and incorporating detailed phylogenetic data for local, national and international comparisons. Furthermore, we recommend repeating the structured survey undertaken by Grundmann et al. to provide an update of MRSA epidemiology at the European level [18].

We advocate that professional microbiological societies support guideline development for harmonisation. Due to its focus, aims, international representation and goals, ISAC could take the lead in this process. These guidelines should include BSI/SSTI definitions and a report template for epidemiological metadata. Additionally, a

feasible ratio of BSI/SSTI cases for inclusion should be determined in collaboration with programme representatives. Furthermore, we recommend the development of an international repository for standardised surveillance data, including WGS data. Other suggestions for the harmonisation of antimicrobial resistance surveillance should be considered [4,12,19,20], such as the alignment of surveillance goals and standardised methodology for data collection, data analysis and data sharing.

Although many countries expend substantial effort and resources on MRSA surveillance, the stability of financial support is a general concern. This should be recognised in guideline development, as national health budgets will greatly influence the opportunities for harmonisation of surveillance programmes.

Inclusion bias may have limited the generalisability of our study results. Nevertheless, we were able to highlight the diversity of surveillance programmes, and our webinar enabled MRSA surveillance experts to discuss their differences directly. This guided the development of our proposal for the harmonisation of MRSA surveillance programmes.

In conclusion, current MRSA surveillance programmes rely upon heterogeneous data collection, which hampers international epidemiological monitoring and research. For harmonisation of MRSA surveillance, we suggest including SSTI cases in proportion to collected BSI cases, improving the integration of microbiological and epidemiological data, implementing central biobanks for the collection and further characterisation of MRSA isolates, and genotyping of a structured sample of these isolates, preferably using WGS.

Acknowledgments

The authors thank the MRSA Surveillance Worldwide Study Group for their co-operation. The MRSA Surveillance Worldwide Study Group consists of Arjana Tambic Andrasevic, Valérie O. Baede, Dominique S. Blanc, Michael Borg, Grainne Brennan, Boudewijn Catry, Aurélie Chabaud, Michael Z. David, Joanna Empel, Hege Enger, Amy Gargis, Runa Gokhale, Marie Hallin, Marina Ivanova, Amelie Jouzeau, Andreas Kronenberg, Kuntaman Kuntaman, Anders Rhod Larsen, Katrien Latour, Jodi A. Lindsay, Bruno Pichon, Dewi Santosaningsih, Isaac See, Harald Seifert, Leo M. Schouls, Stefania Stefani, François Vandenesch, Margreet C. Vos, Guido Werner, Dorota Żabicka and Helena Žemličková. The authors are thankful to Fee Johnstone, ISAC Executive Assistant, for organisational support, and to Oz Golbasi and Cem Tunçel of Delta Medical Communications for technical assistance. Additionally, the authors thank the MACOTRA Study Group for fruitful discussions. The MACOTRA Study Group consists of Valérie O. Baede, Anaïs Barray, Sake J. de Vlas, Anneke S. de Vos, Arya Gupta, Antoni P.A. Hendrickx, Gwenan M. Knight, Mirjam E.E. Kretzschmar, Gérard Lina, Jodi A. Lindsay, Jean-Philippe Rasigade, Leo M. Schouls, Mehri Tavakol, Margreet C. Vos, Willem J.B. van Wamel and Adam A. Witney.

Funding

This study was supported by JPIAMR 3rd call, *AMR Transmission dynamics* and Dutch ZonMw [grant no. 547001006]. The webinar was financially supported by the International Society of Antimicrobial Chemotherapy (ISAC).

Competing interests

None declared.

Ethical approval

Not required.

Supplementary materials

Supplementary material associated with this article can be found, in the online version, at [doi:10.1016/j.ijantimicag.2022.106538](https://doi.org/10.1016/j.ijantimicag.2022.106538).

References

- [1] Cassini A, Högberg LD, Plachouras D, Quattrocchi A, Hoxha A, Simonsen GS, et al. Attributable deaths and disability-adjusted life-years caused by infections with antibiotic-resistant bacteria in the EU and the European Economic Area in 2015: a population-level modelling analysis. *Lancet Infect Dis* 2019;19:56–66. doi:10.1016/S1473-3099(18)30605-4.
- [2] Enright MC, Robinson DA, Randle G, Feil EJ, Grundmann H, Spratt BG. The evolutionary history of methicillin-resistant *Staphylococcus aureus* (MRSA). *Proc Natl Acad Sci U S A* 2002;99:7687–92. doi:10.1073/pnas.122108599.
- [3] Stefani S, Chung DR, Lindsay JA, Friedrich AW, Kearns AM, Westh H, et al. Methicillin-resistant *Staphylococcus aureus* (MRSA): global epidemiology and harmonisation of typing methods. *Int J Antimicrob Agents* 2012;39:273–82. doi:10.1016/j.ijantimicag.2011.09.030.
- [4] Diallo OO, Baron SA, Abat C, Colson P, Chaudet H, Rolain J-M. Antibiotic resistance surveillance systems: a review. *J Glob Antimicrob Resist* 2020;23:430–8. doi:10.1016/j.jgar.2020.10.009.
- [5] European Centre for Disease Prevention and Control (ECDC) Surveillance of antimicrobial resistance in Europe 2018. Annual report of the European Antimicrobial Resistance Surveillance Network (EARS-Net). Stockholm, Sweden: ECDC; 2019.
- [6] European Committee on Antimicrobial Susceptibility Testing (EUCAST). <https://eucast.org/> [accessed 23 June 2021].
- [7] Song J-H, Hsueh P-R, Chung DR, Ko KS, Kang C-I, Peck KR, et al. Spread of methicillin-resistant *Staphylococcus aureus* between the community and the hospitals in Asian countries: an ANSORP study. *J Antimicrob Chemother* 2011;66:1061–9. doi:10.1093/jac/dkr024.
- [8] Ashley EA, Recht J, Chua A, Dance D, Dhorda M, Thomas NV, et al. An inventory of supranational antimicrobial resistance surveillance networks involving low- and middle-income countries since 2000. *J Antimicrob Chemother* 2018;73:1737–49. doi:10.1093/jac/dky026.
- [9] World Health Organization (WHO)/Pan American Health Organization (PAHO). *Latin American Network of Antimicrobial Resistance Surveillance – ReLAVRA*. <https://www.paho.org/en/topics/antimicrobial-resistance/latin-american-network-antimicrobial-resistance-surveillance> [accessed 13 December 2021].
- [10] Diekema DJ, Pfaller MA, Shortridge D, Zervos M, Jones RN. Twenty-year trends in antimicrobial susceptibilities among *Staphylococcus aureus* from the SEN-TRY Antimicrobial Surveillance Program. *Open Forum Infect Dis* 2019;6(Suppl 1):S47–53. doi:10.1093/ofid/ofy270.
- [11] Seifert H, Blondeau J, Dowdzicki MJ. In vitro activity of tigecycline and comparators (2014–2016) among key WHO ‘priority pathogens’ and longitudinal assessment (2004–2016) of antimicrobial resistance: a report from the T.E.S.T. study. *Int J Antimicrob Agents* 2018;52:474–84. doi:10.1016/j.ijantimicag.2018.07.003.
- [12] Tacconelli E, Sifakis F, Harbarth S, Schrijver R, van Mourik M, Voss A, et al. Surveillance for control of antimicrobial resistance. *Lancet Infect Dis* 2018;18:e99–106. doi:10.1016/S1473-3099(17)30485-1.
- [13] Grundmann H, Klugman KP, Walsh T, Ramon-Pardo P, Sigauque B, Khan W, et al. A framework for global surveillance of antibiotic resistance. *Drug Resist Updat* 2011;14:79–87. doi:10.1016/j.drup.2011.02.007.
- [14] . *Healthcare-Associated Infections–Community Interface (HCAI). Emerging Infections Program (EIP) Network Report: invasive Staphylococcus aureus*. CDC; 2017. p. 2020.
- [15] Wertheim HFL, Vos MC, Ott A, van Belkum A, Voss A, Kluytmans JAJW, et al. Risk and outcome of nosocomial *Staphylococcus aureus* bacteraemia in nasal carriers versus non-carriers. *Lancet* 2004;364:703–5. doi:10.1016/S0140-6736(04)16897-9.
- [16] von Eiff C, Becker K, Machka K, Stammer H, Peters G. Nasal carriage as a source of *Staphylococcus aureus* bacteremia. *N Engl J Med* 2001;344:11–16. doi:10.1056/NEJM200101043440102.
- [17] US Centers for Disease Control and Prevention (CDC). *National Healthcare Safety Network (NHSN). Patient Safety Component Manual*. CDC; 2021. <https://www.cdc.gov/nhsn/psc/index.html> [accessed 23 June 2021].
- [18] Grundmann H, Schouls LM, Aanensen DM, Pluister GN, Tami A, Chlebowicz M, et al. The dynamic changes of dominant clones of *Staphylococcus aureus* causing bloodstream infections in the European region: results of a second structured survey. *Euro Surveill* 2014;19:20987. doi:10.2807/1560-7917.ES2014.19.49.20987.
- [19] World Health Organization (WHO) Global antimicrobial resistance surveillance system (GLASS) report: early implementation 2020. Geneva, Switzerland: WHO; 2020. Licence: CC BY-NC-SA 3.0 IGO.
- [20] World Health Organization (WHO) GLASS Emerging antimicrobial resistance reporting framework (GLASS-EAR), Geneva, Switzerland: WHO; 2018. Licence: CC BY-NC-SA 3.0 IGO.