

# Current Opinion in Otolaryngology and Head & Neck Surgery

## Transitions in tracheostomy care: from childhood to adulthood

--Manuscript Draft--

<b>Manuscript Number:</b>	MOO310302R1
<b>Full Title:</b>	Transitions in tracheostomy care: from childhood to adulthood
<b>Article Type:</b>	Review Article
<b>Corresponding Author:</b>	Louise Edwards, BSc(Hons) University of Central Lancashire Huntingdon, Cambridgeshire UNITED KINGDOM
<b>Corresponding Author Secondary Information:</b>	
<b>Corresponding Author's Institution:</b>	University of Central Lancashire
<b>Corresponding Author's Secondary Institution:</b>	
<b>First Author:</b>	Louise Edwards, BSc(Hons)
<b>First Author Secondary Information:</b>	
<b>Order of Authors:</b>	Louise Edwards, BSc(Hons) Jackie McRae
<b>Order of Authors Secondary Information:</b>	

## Transitions in tracheostomy care: from childhood to adulthood

Louise Edwards<sup>a,b</sup> and Jackie McRae<sup>c,d</sup>

Corresponding author:

Ms Louise Edwards MRCSLT

<sup>a</sup>Professional Doctorate in Health student, University of Central Lancashire, UK

<sup>b</sup>Clinical Educator / Speech & Language Therapist, Atos Medical, UK

Tel: +44 (0) 7946637794

Email: [imedwards2@uclan.ac.uk](mailto:imedwards2@uclan.ac.uk)

Co-author:

Jackie McRae PhD FRCSLT

<sup>c</sup>Associate Professor, Centre for Allied Health, St George's University of London, UK

<sup>d</sup>Honorary Consultant Speech and Language Therapist, University College Hospitals NHS

Foundation Trust, UK

Email: [jmcrac@sgul.ac.uk](mailto:jmcrac@sgul.ac.uk)

## **Abstract**

**Purpose of review:** The purpose of this review is to explore the evidence around children and young people who require a tracheostomy and transition into adult services, reflecting on the challenges and considerations for clinical practice as these needs increase.

**Recent findings:** There are a lack of data on the incidence and prevalence of children and young people with a tracheostomy transitioning to adult services for ongoing care. There are significant variations in care needs, technology and previous experiences that demand more than a simple handover process. Examples of service models that support the transition of care exist, however these lack specificity for children and young people with a tracheostomy.

**Summary:** Further exploration of the needs of children and young people requiring airway technology is indicated, particularly considering the short and long-term education, health, and social care needs.

**Keywords:** tracheostomy, transition, paediatric, airway, children and young people

## Introduction

A tracheostomy is a medical device made of polyvinyl chloride (PVC) or silicone that, in the paediatric population, is typically inserted as part of a planned, surgical procedure to maintain a patent airway and effective ventilation [1]. A tracheostomy can be placed anytime from birth as an EXIT (ex-utero intrapartum treatment) procedure [2] and across the age continuum to safeguard the airway and respiratory system.

In high income countries, recent advances in healthcare, technology, and clinical expertise have increased survivability of many paediatric conditions that were previously life-limiting through compromised respiratory function. This has resulted in lower thresholds for paediatric tracheostomy insertion as a long-term solution for life maintenance. In low- or middle-income countries, the on-going challenges of managing complex airways in the paediatric population are acknowledged, although detailed data are limited [3-4\*\*].

Indications for short-term tracheostomy placement in children and young people (CYP) due to chronic airway obstruction secondary to infections, are infrequent due to mass immunisation programmes [5\*\*]. Current indications for tracheostomy in children and young people reflect the advances in managing respiratory conditions, particularly associated with prematurity resulting in long-term ventilation, and identification of congenital upper airway obstruction in utero or early infancy [6, 7\*]. Additionally, advancing pharmacological and clinical expertise have enabled survival from significant acquired conditions such as brain injury and trauma [8].

Placement of a tracheostomy, particularly for paediatric populations is associated with significant mortality and morbidity risks, although it is debated whether this relates to factors of the underlying diagnoses [9\*], as management of the device (tracheostomy) remains a significant risk factor for mortality [5\*\*, 10, 11\*].

Whilst clinically, mortality has decreased in CYP with a range of underlying conditions (Table 1), morbidity and complexity have increased with greater survival and need for lifelong care. This requires an effective transition into adult services [9\*, 12], with collaboration between adult and paediatric colleagues to understand the context of lifelong paediatric conditions, whether acquired, developmental or degenerative.

Table 1: Paediatric conditions associated with requiring long-term tracheostomy

### **Developmental, psychological, and social context**

The presence and management of a tracheostomy is likely to be embedded into the life and daily schedule of CYP, with all the considerations of childhood development, inclusion, and identity [13] needed by the time they transition into adult services. Additionally, with reportedly increasing prevalence of such technology needs in CYP, the impact on healthcare systems [14], including transition, is highlighted.

Issues at the point of transition have been highlighted in this population, along with similar populations such as CYP with neurodisabilities [15\*\*]. Psychosocial issues for the child are emphasised as they may grapple with identity and difference in social and education

environments that focus on the piece of equipment (tracheostomy) or underlying disease, rather than the skills and strengths of the young person [16\*].

Acknowledging the individual beyond technology is essential in supporting a safe and successful transition into adulthood, and therefore navigating adult services [17\*].

In order to understand the long-term tracheostomy needs of CYP it is important to understand the evidence around removal of the tracheostomy (decannulation) as a clinical goal and how this affects transition to adult services.

### ***Decannulation***

Decannulation rates in the paediatric population range have been variably cited in the literature from 28-52% [11\*] to 31 - 75% [18\*] although long term data is lacking. The Global Tracheostomy Collaborative (GTC) has reported on a number of initiatives to improve tracheostomy care, including highlighting the importance of multidisciplinary clinics, applicable to the adult as much as paediatric populations [4\*\*]. Their review of paediatric and adult data from 62 institutions across six countries showed the paediatric population had an increase in duration of tracheostomy placement (median 16 vs. 36 days) and reduced decannulation rate (4% vs. 52%). A recent longitudinal study in a single centre in the United Kingdom (UK) identified a 44.4% decannulation rate in 172 children over an eight year period and a mortality rate of 22.1% [19]. They report a median duration of 397 days with a tracheostomy. Another single site review over 11 years reported a 39% decannulation in 105 children with a median tracheostomy duration of 408 days [11\*]. Compared to the adult population, where tracheostomy placement is typically short term [4\*\*] and maintained in the acute setting, for example, due to an acquired impairment, the

goal will be decannulation after weaning from ventilation. However, CYP with a tracheostomy may require an artificial airway for a longer term and into adulthood [15\*\*]. Success, or lack thereof, of decannulation can determine the need for long-term placement of tracheostomy. Whilst there have been significant advances in airway surgical techniques and improving outcomes, decannulation failure remains an issue due to the high frequency of airway abnormalities seen in children with tracheostomy [20\*\*] particularly for populations with persistent lower airway impairments such as malacia or stenosis [21] so careful airway evaluation is needed before decannulation. Additionally, failure of decannulation is also particularly prevalent in at-risk populations, for example Trisomy 21 [22], those of poor socioeconomic backgrounds [23], and where failure to thrive and neurological issues exist [9\*, 11\*] again suggesting the need for long-term tracheostomy.

### **Preparing for transition**

In paediatric populations, the burden of care is acknowledged and the need for resilience highlighted [24\*, 25\*]. In the United Kingdom and indeed in other high-income countries, typically, CYP with a tracheostomy are supported by paediatric teams across primary, secondary, and tertiary healthcare settings, including but not limited to acute Ear, Nose, and Throat (ENT) teams, acute and community paediatricians, Allied Healthcare Professionals (AHP) in health and education, and children's community nursing teams. However, as CYP transition into adult care, there remains a risk of abandonment [26] and isolation [27] with a lack of coordinated services as no equivalent of a paediatrician exists, an '*adultrician*' if you will, in adult services until geriatric services become appropriate. In contrast, there are services that lack paediatric specialists, such as in spinal cord injury, leaving them to be managed by adult services who may not have the skills to support behavioural and social

needs [28]. It is acknowledged that whilst some services exist that support continuing care and prepare for transition, for example in established Long-Term Ventilation teams, this is primarily for a population requiring on-going invasive or non-invasive ventilation [29].

Understanding of the extent of service access needs for all CYP with technology needs is lacking, particularly those not requiring ventilation.

### **Navigating different technologies**

There may be a number of variations in the technology or tasks used with the paediatric tracheostomy population so healthcare staff in adult settings need to be aware of what is acceptable and routine practice. Typically, single lumen tracheostomy tubes are utilised until the child or young person has more adult-like upper airway anatomy, and transitions to a double lumen tube as used in adults. Cuffed tracheostomy tubes used in the adult population may have a different method of cuff management, for example air filled, compared to paediatric cuffed tubes that may be water, air, or less typically, foam filled. In the paediatric population the long-term use of water filled cuffed tracheostomy is not recommended due to risk of tracheal wall damage and difficulties managing cuff pressures [30].

Familiar and regular tasks may be differently approached or expedited in the adult setting, for example, sizing of a suction catheter and suction depths and pressures [31\*]. Families and carers of CYP with a tracheostomy would have undergone significant training and skill development to support transition from an acute care setting to home and ensure 1:1 care [32]. With changing technology and management needs, this training needs revisiting and



revising in the adult setting, to optimise safety and optimise tracheostomy management which may present as a further burden to families and carers, particularly during a period of change. This raises the question as to who provides the training and at what point before a young person transitions and requires a change to tracheostomy management.

As noted, around the age that transition ideally should be discussed, there may be changes to airway and tracheostomy tube requirements and therefore implications for parents and carers and indeed healthcare clinicians previously trained in the management of single lumen tubes. Anecdotally, the change in tubes can be unfamiliar and daunting, requiring updated training and support. Double lumen tubes may come with additional features not typically experienced in paediatric, single lumen tubes, for example, subglottic suctioning ports.

Therefore, the paediatric clinician needs to be versed in the wider range of tubes, specifically double lumen tubes that may be necessary for growing CYP and more commonly used before transition. However, typically due to the underlying aetiology requiring the need for single lumen tubes into adulthood, such as craniofacial anomalies, the adult clinician is also encouraged to appreciate the variation and features of single lumen tubes to inform clinical practice, especially in emergency scenarios.

As with the adult population, the purpose of a one-way valve in paediatrics, where clinically indicated, is to optimise all upper airway functions through laryngeal (re)sensitisation, including voice, cough, bearing down, and swallowing, rather than facilitating tracheostomy weaning. Where indicated, the approach and assessment of one-way valve use may need

consideration when the young person transitions to the adult setting. It is possible that CYP may benefit from the use of open-bias one-way valves [33]. Clinically, these offer less resistance and for some children may offer greater acceptability and tolerance, particularly in the presence of sensory or neurological disabilities. Indeed, anecdotally, weaning in the paediatric population may not involve one-way valve use at all, but otherwise optimise downsizing of the tracheostomy tube and capping off of the tube whilst admitted to an acute care or increasingly within a community setting [20\*\*, 34], taking place over consecutive days and nights as opposed to a 24-hour period for adults. There remains, however, considerable variation and lack of clarity in decannulation protocols globally [20\*\*].

### **Redefining parent and carer roles**

Parents and carers of CYP with a new tracheostomy typically undergo an intensive, competency and confidence-based programme of education within an acute setting to facilitate discharge into the home or community setting [35] and ensure the essential skilled, 1:1 care required to manage the tracheostomy. This education and support can take months or even years, depending on personal, social, and environmental factors. It is acknowledged that for families this can be a significant burden of care that may be lifelong, however parents have reported a sense of resilience and determination to adapt to challenging circumstances and remain hopeful [24\*].

Transition may be more clearly defined for some populations, for example those requiring long-term ventilation, with examples of transition clinics noted around the UK and a recently established UK national peer support and advisory group on paediatric long-term ventilation. The group has highlighted the increasing transition from child to adult services

and plans to explore practice throughout the UK [29]. However, other populations, such as those with a tracheostomy without ventilation, may be less well served and therefore 'lost' without the scaffolding of an *'adultrician'*.

When transitioning to adult services, the role of parent and carer may be less defined and result in challenges at many stages of healthcare. Acknowledging the role of the parent and carer and ensuring space for them in the on-going care of a young person transitioning services is essential. For example, in the paediatric acute setting, parents and carers are often the primary carer, continuing their management of the tracheostomy throughout an acute admission. Furthermore, with preparation there is an opportunity to support, particularly in vulnerable young people, as well as optimise self-efficacy and participation in future planning and care [36\*\*].

The parent and carers come with extensive knowledge and expertise in their CYP's tracheostomy. Acknowledging the expertise and working in collaboration with families and carers to negotiate and navigate health, social and education experiences is essential to ensuring adaptation [26]. Families and carers have built links and attachments with teams over decades and so change and adjustment may take time; ensuring clear and consistent communication is essential.

### **Preparing for adulthood**

According to the UK National Health Service (NHS) Long Term Plan [37], planning for transition into adult services should occur when a child is around 14 years old, clinical evidence suggests this does not always happen and indeed the services within adulthood

may not be available. This guidance is not atypical of other countries' policy on supporting transition [36\*\*], for example, *GOT TRANSITION*® the 'federally funded national resource centre on health care transition' [38] in the United States of America providing resources for healthcare professionals, young people and families and carers on navigating transitions.

Other informative guidance and support exists, for example, *Together for Short Lives* [39\*], a charity supporting CYP with life-limiting and life-threatening conditions, documents a pathway for healthcare professionals, CYP and families alike to navigate the transition journey.

Rather than awaiting arrival or transfer into adult services, the importance of actively planning for transition is emphasised. Within the NHS in the UK, examples of healthcare services collaborating with service users to ensure safe and timely transition are apparent, for example the *Ready, Steady, Go* Programme [40] originating from University Hospitals Southampton and, although under development, available across the UK [41]. This programme starts the process of transition at the age of 11 years and aims to empower the young person to understand their condition so that they can make informed choices about their future care by the age of 18 years. The programme is generic so can be applied to a range of long-term conditions using the same framework. Globally, the importance of transition clinics to optimise outcomes for CYP has been highlighted and encouraged to commence early with involvement of clinicians, family members and young people [36\*\*, 42]. However, a lack of consistency and transparency remains in managing transition into adult services across health, social and education settings.

Understanding the experience of navigating childhood with a tracheostomy is important to inform how a young person may have experienced identity, self-esteem, and education and how this may impact on their transition into adulthood services. Learning from individuals' lived experiences, particularly those that have navigated the transition from paediatric to adult services with a tracheostomy is essential for supporting clinicians' understanding of the challenges and opportunities faced.

### **Future research directions**

Further research is needed to inform the epidemiology and long-term outcomes of CYP with tracheostomy transitioning to adult services. Furthermore, exploration of the health, social and education continuum and care pathways for this population may better inform service, policy, and clinical priorities. Gathering evidence of the lived experience and patient reported outcomes of those transitioning from paediatric to adult services may identify what is personally meaningful and necessary to support the transition and on-going needs into adulthood.

### **Conclusion**

Whilst CYP with tracheostomy represent a relatively low incidence population, the associated technology, underlying diagnoses, and potential for life-long needs make them a highly complex population. The transition period to adulthood requires overlap and collaboration between paediatric and adult health, social, and education services to facilitate seamless and compassionate care. The involvement of the CYP and their family is essential to navigate these transitions and ensure engagement, inclusion, and effective transition outcomes. It is recognised globally, as discussed, collaboration should commence as early as possible in preparation for transition.

## **Key points**

- Children and young people with long term tracheostomy need a safe and effective transition to adult services
- Their ongoing needs include adapting to education and work environments, social and psychological support
- There is an opportunity for paediatric and adult clinicians to share skills and knowledge to optimise the transition pathway

## **Acknowledgements**

None

## **Financial support and sponsorship**

L Edwards:

None

J McRae:

None

## **Conflict of interest**

L Edwards:

Receives a salary from Atos Medical for the role of Clinical Educator (part time) outside the submitted work.

Reviewer (non-financial) for the NCEPOD Balancing the Pressures (2020) document included in the references.

J. McRae:

None

## References and recommended reading

\* of special interest

\*\* of outstanding interest

1. Watters KF. Tracheostomy in infants and children. *Resp Care* [internet] 2017; 62(6):799-825.
2. Porter H, Trivedi A, Marquez M, et al. Changing indications and antenatal prognostic factors for ex-utero intrapartum treatment procedures. *Prenat Diagn* [internet] 2022; 42(11):1420-8.
3. Sandler ML, Ayele N, Ncogoza I, et al. Improving tracheostomy care in resource-limited settings. *Ann Otolaryngol Rhinol* [internet] 2020; 129(2):181-90.
4. Brenner MJ, Pandian V, Milliren CE, et al. Global Tracheostomy Collaborative: data-driven improvements in patient safety through multidisciplinary teamwork, standardisation, education, and patient partnership. *Br J Anaesth* [internet]. 2020;125(1):e104-18.

Important article from the Global Tracheostomy Collaborative emphasising the importance of partnership and utilisation of data to inform clinical practice and optimise safety in tracheostomy.

5. Fuller C, Wineland AM, Richter GT. Update on pediatric tracheostomy: Indications, technique, education, and decannulation. *Curr Otorhinolaryngol Rep* [internet]. 2021;9(2):188-99.

Detailed review of current practice and changes to paediatric tracheostomy, with

emphasis on surgical technique, complications, and long term outcomes including decannulation.

6. Marlow N, Johnson S, Hurst JR. The extremely preterm young adult—state of the art. *Semin Fetal Neonatal Med* [internet] 2022; p. 101365.

7. Barker N, Sinha A, Jesson C, et al. Changes in UK paediatric long-term ventilation

\* practice over 10 years. *Arch Dis Child*. Published Online First: 29 November 2022. doi: 10.1136/archdischild-2021-323562 [cited 23 December 2022].

UK wide study identifying changing clinical practice with paediatric long-term ventilation and implications for future practice and management.

8. Salik I, Das A, Naftchi AF, et al. Effect of tracheostomy timing in pediatric patients with traumatic brain injury. *Int J Pediatr Otorhinolaryngol* [internet] 2022; 12:111414.

9. Hebbar KB, Kasi AS, Vielkind M, et al. Mortality and outcomes of pediatric

\* tracheostomy dependent patients. *Front Pediatr* [internet] 2021; 9:661512.

Retrospective study of 187 tracheostomy dependent children highlighting the impact of underlying neurological issues on success of decannulation.

10. Yilmaz Yegit C, Kilinc AA, Can Oksay S, et al. The ISPAT project: Implementation of a standardized training program for caregivers of children with tracheostomy. *Pediatr Pulmonol* [internet]. 2022;57(1):176-84.

11. Chia AZH, Ng ZM, Pang YX, et al. Epidemiology of pediatric tracheostomy and risk

\* factors for poor outcomes: An 11-year single-center experience. *Otolaryngol Head Neck Surg* [internet]. 2020;162(1):121-128.



Retrospective analysis of 105 children requiring tracheostomy over an 11-year period highlighting failure to thrive and neurological difficulties impacting on success of decannulation.

12. Jarvi S, Richardson G, Flemming K, Fraser LK. Numbers, characteristics, and medical complexity of children with life-limiting conditions reaching age of transition to adult care in England: a repeated cross-sectional study. NIHR Open Research [internet]. 2022;2:27.

13. Crumpton J, Wray J. Children's and adolescents' experiences of living with respiratory assistance: A systematic review of qualitative studies. Int J Pediatr Otorhinolaryngol [internet] 2019;127:109658.

14. Roland D, Wolfe I, Klaber RE, Watson M. Final warning on the need for integrated care systems in acute paediatrics. Arch Dis Child [internet] 2022;107:e9.

15. NCEPOD (The National Confidential Enquiry into Patient Outcome and Death).

\*\* *Balancing the Pressures*. 2020. London.

Report on a national study reviewing the long-term ventilation services to children and young people (0-25 years) in the UK and recommendations for clinical practice.

16. Belzer LT, Wright SM, Goodwin EJ, et al. Psychosocial considerations for the child with rare disease: A review with recommendations and calls to action. Children [internet] 2022; 9, 933.

A detailed review of the literature around psychosocial issues facing children and young people with rare diseases and their families. The paper highlights recommendations to improve practice and evidence-based care.

17. Bailey K, Lee S, de Los Reyes T, et al. Quality indicators for youth transitioning to adult care: A systematic review. Pediatrics [internet]. 2022;150(1):e2021055033.

A systematic review whilst identifying patient-centred outcomes were key quality indicators, studies lacked involvement of children, young people and their families. The review demonstrated the need to prioritise quality indicators for those navigating chronic illness and involve service users to inform these indicators.

18. Kolb CM, Halbert K, Xiao W, et al. Comparing decannulation failures and successes in \* pediatric tracheostomy: An 18-year experience. *Pediatr Pulmonol* [internet] 2021;56(8):2761-2768.

Retrospective review of 173 decannulation attempts in a cohort of 439 tracheostomies in a single centre over an 18-year period. Shorter period of cannulation and older gestational age informed success of decannulation. The authors acknowledge the need for further research to inform patient selection and timing of decannulation.

19. Roberts J, Powell J, Begbie J, et al. Pediatric tracheostomy: A large single-center experience. *Laryngoscope* [internet] 2020; 130: E375-E380.

20. Verma R, Mocanu C, Shi J, et al. Decannulation following tracheostomy in children: A \*\* systematic review of decannulation protocols. *Pediatr Pulmonol* [internet] 2021; 56(8), 2426-2443.

A systematic review of paediatric decannulation protocols across 24 studies and 1395 children suggesting that large variability in protocols exists with urgent need for further research to optimise guidance. Polysomnography is highlighted as a useful tool in assessing readiness for decannulation.

21. Kou YF, Teplitzky T, Johnson RF, Chorney SR. Assessment of socioeconomic disadvantage in laryngotracheoplasty outcomes among pediatric patients. *Int J Pediatr Otorhinolaryngol* [internet] 2022;162:111326.

22. Blumenthal D, Leonard J, Habib A, et al. Decannulation in Trisomy 21 patients undergoing laryngotracheal reconstruction. *Int J Pediatr Otorhinolaryngol* [internet] 2022;111407.
23. Liao K, Chorney SR, Brown AB, et al. The impact of socioeconomic disadvantage on pediatric tracheostomy outcomes. *Laryngoscope* [internet]. 2021; 131: 2603-2609.
24. Flynn A, Whittaker K, Donne AJ, et al. Holding their own and being resilient: \* narratives of parents over the first 12 months of their child having tracheostomy. *Compr Child Adolesc Nurs* [internet] 2021; 44:4, 255-265.

This longitudinal narrative study explores the experiences of 12 parents at three time points over a 12-month period following their child's tracheostomy. Stories of resilience are emphasised across the journey from hospital to home, as well as influencing outlook.

25. Gursoy TR, Eyuboglu TS, Aslan AT, et al. The associations between the clinical course \* of children with tracheostomy and their mothers' depression, burnout, burden, and self-esteem [published online ahead of print, 2022 Jul 11]. *J. Clin. Nurs* [internet] 2022;10.1111/jocn.16450.

A cross-sectional study exploring mothers' psychological experiences during the care of their child with a tracheostomy. Frequent and long stay hospital admissions were associated with maternal depression and burnout, emphasising the need for preventative psychological support for families.

26. Li L, Carter N, Ploeg J, et al. The role of family adaptation in the transition to adulthood for youth with medical complexity: a qualitative case study protocol. *J Transl Med* [internet]. 2021;3(1): 20210005.

27. Hoskin J. Troubling norms? Adults and teenagers with a life-limiting impairment in Denmark and England talk about their lives, support and future plans. *Eur. J. Spec. Needs Educ* [internet] 2021; 36:3, 329-343.
28. Botchway EN, Knight S, Muscara F, et al. Rehabilitation models of care for children and youth living with traumatic brain and/or spinal cord injuries: A focus on family-centred care, psychosocial wellbeing, and transitions. *Neuropsychol Rehabil* [internet] 2022;32(4):537-59.
29. Everitt L, Wilkinson S, Evans H. 1014 The national paediatric Long-Term Ventilation (LTV) advisory group - A new initiative. *Arch Dis Child* [internet] 2022; 08;107:A243-A244.
30. Tweedie DJ, Cooke J, Stephenson KA, et al. Paediatric tracheostomy tubes: recent developments and our current practice. *J Laryngol Otol* [internet] 2018;132(11):961-968.
31. National Tracheostomy Safety Project. [www.tracheostomy.org.uk](http://www.tracheostomy.org.uk). [Cited 23 \* December 2022].
- Practical and useful organisation and website including guidance, protocols, training, and resources for adult and paediatric tracheostomy knowledge and skill development.
32. Powell J, Buckley HL, Agbeko R, et al. Tracheostomy trends in paediatric intensive care. *Arch Dis Child* [internet] 2021;106(7):712-4.
33. Greene ZM, Davenport J, Fitzgerald S, et al. Tracheostomy speaking valve modification in children: A standardized approach leads to widespread use. *Pediatr Pulmonol* [internet] 2019;54(4):428-35.
34. Canning J, Mills N, Mahadevan M. Pediatric tracheostomy decannulation: when can decannulation be performed safely outside of the intensive care setting? A 10 year

review from a single tertiary otolaryngology service. *Int J Pediatr Otorhinolaryngol* [internet] 2020; 133: 109986.

35. Chiang J, Karim A, Hoffman A, et al. Tough transitions: Family caregiver experiences with a pediatric long-term ventilation discharge pathway. *Pediatr Pulmonol* [internet] 2021;56(10):3380-8.

36. Fremion E, Cowley R, Berens J, et al. Improved health care transition for young adults \*\* with developmental disabilities referred from designated transition clinics. *J Pediatr Nurs* [internet] 2022; 1;67:27-33.

The study, using logistic regression model analysis, identifies the association between access to a healthcare transition clinic prior to transition into adulthood and improved satisfaction and preparedness reporting by young adults with developmental or intellectual disabilities.

37. The NHS long term plan. 2019. [www.longtermplan.nhs.uk](http://www.longtermplan.nhs.uk) [Cited 30 December 2022].

38. GOT TRANSITION®. [www.gottransition.org](http://www.gottransition.org) [Cited 30 December 2022].

39. Together for Short Lives.

\* [www.togetherforshortlives.org.uk/changing\\_lives/supporting-care-professionals/care-pathways/](http://www.togetherforshortlives.org.uk/changing_lives/supporting-care-professionals/care-pathways/) [Cited 30 December 2022].

This charity provides free access to a suggested care pathway to support children and young people with life-threatening and life-limiting illnesses transitioning to adult services.

40. Ready, Steady Go Programme. <https://www.readysteadygo.net/> [Cited 3 January 2023].

41. Nagra A, McGinnity PM, Davis N, Salmon AP. Implementing transition: Ready Steady Go. *Arch Dis Child Educ Pract Ed* [internet]. 2015; 100(6): 313-320.

42. Teed M, Bekx A, Paul M, Younker L. Health care transition for children with medical complexity: Challenges and lessons learned. *J Pediatr Nurs* [internet] 2021;61:275-9.

Table 1: Paediatric conditions associated with requiring long-term tracheostomy

Congenital conditions	Craniofacial conditions, e.g., Apert's syndrome Laryngeal atresia Congenital heart disease
Acquired conditions (traumatic or non-traumatic)	Spinal cord injury Acquired brain injury Caustic soda ingestion Tumour Stroke
Progressive neuromuscular conditions (where non-invasive ventilation is unsupported)	Muscular dystrophy Duchenne's muscular dystrophy Spinal muscular atrophy
Respiratory conditions	Chronic ventilation needs Chronic lung disease Laryngotracheobronchomalacia