



Management of subcutaneous abscesses: prospective cross-sectional study (MAGIC)

White Rose Surgical Collaborative and MAGIC Collaborators

Correspondence to: Stephen J. Chapman, Room 7.16, Clinical Sciences Building, St James's University Hospital, University of Leeds, Leeds LS2 9JT, UK (e-mail: s.chapman@leeds.ac.uk; [X@SJ_Chapman](https://twitter.com/SJ_Chapman))

Members of the White Rose Surgical Collaborative and MAGIC Collaborators are co-authors of this study and are listed under the heading Collaborators.

Introduction

Subcutaneous abscesses are pus-filled cavities within the dermis and subdermal layers caused by a local bacterial infection, most commonly *Staphylococcus aureus*¹. They are a common presentation, with an incidence of 433 per 100 000 presentations in community practice, leading to 236 per 100 000 admissions to hospital each year². Surgical drainage is the conventional approach used to treat abscesses in the hospital setting. This aims to achieve complete evacuation of pus, thorough irrigation, and healing by secondary intention.

In the UK, no widely established guidance exists for the management of subcutaneous abscesses. The results of a survey of general surgeons in 2018 highlighted stark disparities in the everyday management of abscesses, such as in the choice of anaesthesia, the role of cavity packing, and the role of swabs for culture; in particular, uncertainty surrounding the use of general or local anaesthesia during surgical drainage was shown to exist³. A cohort study at a large teaching hospital later demonstrated the safety of a protocolized care pathway for drainage under local anaesthesia when applied to the majority (84%) of patients with uncomplicated abscesses⁴. The choice of anaesthesia remains of interest due to its potential impact on health resources, costs, the environment, and emergency care capacity.

Another uncertainty is the role of antibiotics after surgical drainage. In the previous survey, the majority of respondents indicated that antibiotics were only indicated sometimes (39.3%) or in rare circumstances (44.3%)³. This was in contrast to a recently published practice recommendation, advocating for the routine use of antimicrobial therapy after surgical drainage, specifically trimethoprim/sulfamethoxazole or clindamycin⁵. This recommendation was made owing to the results of a systematic review published in 2018, which demonstrated a modest reduction in abscess recurrence with antibiotics across 14 RCTs. In contrast, the review revealed an increase in the risk of gastrointestinal side effects, such as diarrhoea, and also raised concerns about other possible harms, such as from antimicrobial resistance^{5,6}.

The aim of this study was to explore the management of subcutaneous abscesses in emergency surgical practice in the UK. In doing so, the study aimed to describe the UK experience of local anaesthesia as the primary anaesthetic modality for

surgical drainage and the experience of antibiotic therapy as an adjuvant treatment after surgery.

Methods

Study governance

The study protocol was made prospectively available online. The Health Research Authority Decision Tool and a National Health Service (NHS) Research & Innovation Service confirmed that NHS Research Ethics Approval was not required. The study was registered with governance departments at each participating hospital, including confirmation of Caldecott Guardian approval for the safe transfer of non-identifiable data via the Research Electronic Data Capture (REDCap) platform⁷. The study is reported in line with the STROBE checklist⁸.

Study design

A prospective, multicentre, observational study was delivered across NHS hospitals in the UK by the White Rose Surgical Collaborative. No change to routine clinical practice was made. The study opened for data inclusion on 1 September 2022 and closed on 1 December 2022, followed by a 60-day follow-up interval of routinely collected data. The inclusion interval was split into 1-month data collection intervals (*Table S1*), with participating hospital sites asked to contribute to at least one interval. A hospital-specific, 10-item survey exploring standards of management at each hospital was piloted and disseminated to local lead investigators.

Study setting

Any secondary or tertiary care NHS hospital providing an acute emergency general surgery service in the UK was eligible to contribute. No limits on case volume or local policy for the management of acute subcutaneous abscesses were applied. Included patients were identified from any relevant acute care setting, including accident and emergency departments, acute surgical assessment and ambulatory units, and inpatient wards.

Eligibility criteria

Any adult (greater than or equal to 18 years old) presenting with an acute subcutaneous abscess was eligible for inclusion. The

Received: March 09, 2024. Revised: May 01, 2024. Accepted: June 09, 2024

© The Author(s) 2024. Published by Oxford University Press on behalf of BJS Foundation Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

abscess location had to be on the posterior neck, anterior or posterior thorax, abdomen, groin, buttock, natal cleft, or proximal limbs. All other abscesses, including peri-anal, breast, distal limb, and facial abscesses, were excluded, as these were considered to demand specialty-specific treatment not provided by an emergency general surgical practice. The remaining exclusion criteria comprised abscesses associated with a previous surgical site, abscesses with known body cavity communication, and lesions or ulcers considered to be of vascular aetiology.

Outcome measures

The key outcomes of interest were abscess recurrence and return to hospital. Recurrence was defined as any recurrent abscess or surgical-site infection at the same anatomical site. Return to hospital was defined as any re-presentation, irrespective of whether this led to an inpatient admission. Both outcomes were measured using routinely collected data up to 60 days from the index treatment. For patients managed with antibiotics, the index treatment was the time of first antibiotics. For patients

managed with surgery, the index treatment was the time of surgery.

Explanatory variables

Explanatory variables were collected to provide a risk-adjusted estimate of outcomes. These included patient-related factors, such as age (years), sex, BMI (kg/m²), smoking status, and relevant co-morbidities (diabetes mellitus, hidradenitis suppurativa, and immunosuppressive therapeutics or conditions), and treatment-related factors, such as pre-hospital and hospital treatment (surgical drainage, needle aspiration, antibiotics, or no treatment). For patients undergoing surgery, anaesthetic and surgical variables were also collected, such as type of anaesthesia (general, local, or other), surgical setting (operating room, emergency department, or ward/treatment area), and operator grade.

Statistical analysis

Patient and treatment characteristics for the entire cohort are presented descriptively as rates, mean(s.d.) values, or median (interquartile range (i.q.r.)) values. Effect estimates are presented as OR (95% c.i.) values. For the subgroup of patients undergoing surgical drainage, multivariable regression models were constructed for each of the clinical outcomes (abscess recurrence and return to hospital) to assess whether either of these was independently associated with: the use of postoperative antibiotics; the use of local anaesthetic; and packing of the abscess cavity. Explanatory variables were entered into the models using a stepwise approach. In addition, to examine

Table 1 Patient characteristics

Patient characteristic	Category	Value
Age (years)	18–25	303 (18.6)
	26–35	351 (21.5)
	36–45	299 (18.3)
	46–55	285 (17.5)
	56–65	194 (11.9)
Sex	>65	199 (12.2)
	Male	871 (53.4)
Smoking status*	Female	760 (46.6)
	Current smoker	490 (30.9)
	Previous smoker	193 (12.2)
BMI (kg/m ²)*, mean(s.d.)	Never smoked	904 (57.0)
	–	29.3(7.1)
Previous abscess	–	–
	No	1002 (61.5)
Hidradenitis suppurativa	Yes	628 (38.5)
	No	1502 (92.1)
Immunosuppression	Yes	129 (7.9)
	No	1567 (96.1)
Diabetes mellitus	Yes	64 (3.9)
	No	1377 (84.4)
	Yes—diet controlled	38 (2.3)
	Yes—tablet controlled	139 (8.5)
Abscess position	Yes—insulin controlled	77 (4.7)
	Abdominal wall—anterior	136 (8.3)
	Abdominal wall—posterior	53 (3.2)
	Axilla	111 (6.8)
	Buttock	199 (12.2)
	Groin	190 (11.6)
	Lower limb—proximal	62 (3.8)
	Pilonidal	331 (20.3)
	Thorax—anterior	100 (6.1)
	Thorax—posterior	260 (15.9)
Maximum abscess diameter (cm)*, mean(s.d.)	Upper limb—proximal	118 (7.2)
	Other†	71 (4.4)
Timing of hospital presentation	–	4.2(2.3)
	Daytime (08.00–16.59)	1189 (72.9)
	Evening (17.00–21.59)	288 (17.7)
Pre-hospital treatment	Night (22.00–07.59)	154 (9.4)
	No pre-hospital treatment	717 (44.0)
Antibiotics	Antibiotics	867 (53.2)
	Surgery	29 (1.8)
	Other	18 (1.1)

Values are n (%) unless otherwise indicated. *Smoking status based on 1587 responses (n = 44 missing), BMI based on 1378 responses (n = 253 missing), and maximum abscess diameter based on 1458 responses (n = 146 missing). †Other comprises perineal abscesses (n = 7) and posterior neck abscesses (n = 64).

Table 2 Treatment characteristics

Treatment characteristic	Category	Value
Primary treatment	Surgical drainage	1351 (82.8)
	Needle aspiration	30 (1.8)
	Antibiotics only	210 (12.9)
	No treatment	31 (1.9)
Timing of surgery†	Other*	9 (0.6)
	First hospital presentation	917 (67.9)
Preoperative antibiotics†	Planned return presentation	434 (32.1)
	No	634 (47.0)
Surgical setting†	Yes	716 (53.0)
	Operating room	922 (68.2)
	Emergency department	20 (1.5)
Anaesthetic type†	Ward/treatment area	409 (30.3)
	General anaesthetic	874 (64.7)
	Local anaesthetic	441 (32.6)
Primary operator†	Other‡	36 (2.7)
	Consultant	75 (5.6)
	Junior surgical trainee	541 (40.1)
	Nurse practitioner	50 (3.7)
Type of incision†	Senior surgical trainee	682 (50.6)
	Cruciate	33 (2.5)
	Elliptical	238 (18.4)
Abscess packed†	Linear	1024 (79.1)
	No	213 (15.8)
Postoperative antibiotics†	Yes	1136 (84.2)
	No	648 (48.0)
	Yes—≥3 days total	579 (42.9)
	Yes—<3 days total	123 (9.1)

Values are n (%). Junior surgical trainee relates to postgraduate year 1–4 or junior clinical fellow and senior surgical trainee relates to postgraduate year 5–10 or senior clinical fellow. *Other comprises chlorhexidine/saline washout (n = 4), silver nitrate (n = 2), cold spray alone for symptomatic relief (n = 1), and surgery was planned, but patient declined or did not attend (n = 2). †Data relate to patients undergoing surgical drainage (n = 1351). ‡Other comprises spinal (n = 22), regional plexus block (n = 4), cold spray alone (n = 6); Entonox, and other inhalational gas alone (n = 4).

Table 3 Abscess recurrence within 60 days

Characteristic	Category	Univariable OR (95% c.i.), P	Multivariable OR (95% c.i.), P	
Postoperative antibiotics	No	Reference	Reference	
	Yes—≥3 days total	0.99 (0.56–1.76), 0.980	0.97 (0.53–1.74), 0.906	
	Yes—<3 days total	0.62 (0.15–1.82), 0.446	0.64 (0.15–1.91), 0.478	
Anaesthetic type	General anaesthetic	Reference	Reference	
	Local anaesthetic	1.18 (0.65–2.08), 0.569	1.13 (0.60–2.04), 0.698	
Age (years)	18–25	Reference	–	
	26–35	1.32 (0.54–3.41), 0.544	–	
	36–45	1.26 (0.48–3.37), 0.638	–	
	46–55	1.41 (0.57–3.68), 0.464	–	
	56–65	0.81 (0.24–2.48), 0.717	–	
	>65	0.83 (0.25–2.54), 0.746	–	
Sex	Male	Reference	–	
	Female	0.84 (0.48–1.47), 0.553	–	
Smoking status	Current smoker	Reference	Reference	
	Previous smoker	0.37 (0.11–0.99), 0.073	0.38 (0.11–1.02), 0.080	
	Never smoked	0.62 (0.35–1.11), 0.105	0.67 (0.37–1.23), 0.191	
BMI (kg/m ²), mean(s.d.)	–	1.05 (1.01–1.08), 0.006	1.04 (1.01–1.08), 0.015	
Hidradenitis suppurativa	No	Reference	–	
	Yes	3.02 (1.33–6.21), 0.004	2.64 (1.13–5.62), 0.016	
Immunosuppression	No	Reference	–	
	Yes	0.99 (0.16–3.35), 0.985	–	
Diabetes mellitus	No	Reference	–	
	Yes	1.19 (0.55–2.31), 0.638	–	
Abscess position	Abdominal wall—anterior	Reference	–	
	Abdominal wall—posterior	0.67 (0.03–4.73), 0.726	–	
	Axilla	0.67 (0.09–3.54), 0.651	–	
	Buttock	1.39 (0.41–5.45), 0.605	–	
	Groin	2.11 (0.66–8.02), 0.225	–	
	Lower limb—proximal	1.52 (0.20–8.16), 0.640	–	
	Pilonidal	1.76 (0.61–6.34), 0.331	–	
	Thorax—anterior	1.57 (0.36–6.85), 0.536	–	
	Thorax—posterior	0.51 (0.12–2.22), 0.354	–	
	Upper limb—proximal	1.68 (0.43–7.00), 0.452	–	
	Other	0.51 (0.03–3.57), 0.553	–	
	Maximum abscess diameter (cm), mean(s.d.)	–	0.98 (0.86–1.10), 0.776	–
	Presence of cellulitis	No	Reference	–
Yes		1.05 (0.56–1.88), 0.874	–	
Timing of hospital presentation	Daytime (08.00–16.59)	Reference	–	
	Evening (17.00–21.59)	0.81 (0.35–1.68), 0.605	–	
	Night (22.00–07.59)	0.89 (0.30–2.11), 0.806	–	
Timing of surgery	First hospital presentation	Reference	–	
	Planned return presentation	0.99 (0.53–1.75), 0.961	–	
Surgical setting	Operating room	Reference	–	
	Emergency department	4.54 (0.67–18.5), 0.059	–	
	Ward/treatment area	1.19 (0.64–2.14), 0.564	–	
Primary operator	Consultant	Reference	–	
	Junior surgical trainee	0.73 (0.27–2.56), 0.577	–	
	Nurse practitioner	0.59 (0.08–3.16), 0.548	–	
	Senior surgical trainee	0.54 (0.20–1.90), 0.273	–	
Abscess packed	No	Reference	–	
	Yes	0.38 (0.21–0.72), 0.002	0.38 (0.20–0.74), 0.003	

Analyses performed on cases without any missing data ($n = 1047$). Junior surgical trainee relates to postgraduate year 1–4 or junior clinical fellow and senior surgical trainee relates to postgraduate year 5–10 or senior clinical fellow.

whether variation existed between centres, multivariable models containing only patient-level variables along with corresponding multi-level models treating the admitting hospital as a random effect were generated for postoperative antibiotics, anaesthetic type, and abscess packing as outcomes. Residual and standard deviations of the random effects were examined and likelihood ratio tests were performed to identify differences in the models at the centre level. This generated P values to determine whether statistically significant centre-level variation existed for each outcome independent of case mix. Analyses were performed using SPSS® (IBM, Armonk, NY, USA; version 22) and R statistical computing software (R Foundation for Statistical Computing, Vienna, Austria; version 4.3.3), with $P < 0.050$ indicative of statistical significance.

Results

Summary of recruitment

A total of 1637 records were collected from 54 NHS hospitals in the UK. After removal of duplicates, 1631 were considered in the analyses. The median number of records submitted per hospital was 27 (i.q.r. 15–44.5) during a median of 2 (i.q.r. 1–2) data collection intervals.

Hospital-specific survey

A response to the survey was received from 54 of 54 hospitals (100%). A defined clinical pathway or protocol for the management of subcutaneous abscesses was in place at 14 hospitals (14 of 54; 25.9%). For surgery taking place in the

Table 4 Return to hospital within 60 days

Characteristic	Category	Univariable OR (95% c.i.), P	Multivariable OR (95% c.i.), P
Postoperative antibiotics	No	Reference	Reference
	Yes—≥3 days total	1.11 (0.79–1.55), 0.547	0.99 (0.70–1.41), 0.976
	Yes—<3 days total	0.78 (0.39–1.44), 0.448	0.75 (0.37–1.41), 0.398
Anaesthetic type	General anaesthetic	Reference	Reference
	Local anaesthetic	0.97 (0.68–1.36), 0.841	1.00 (0.70–1.42), 0.995
Age (years)	18–25	Reference	–
	26–35	1.14 (0.69–1.91), 0.614	–
	36–45	0.84 (0.48–1.48), 0.551	–
	46–55	0.94 (0.55–1.63), 0.832	–
	56–65	1.07 (0.59–1.90), 0.830	–
	>65	0.79 (0.42–1.46), 0.454	–
Sex	Male	Reference	–
	Female	0.88 (0.64–1.22), 0.459	–
Smoking status	Current smoker	Reference	–
	Previous smoker	0.86 (0.50–1.43), 0.564	–
	Never smoked	0.82 (0.58–1.18), 0.278	–
BMI (kg/m ²), mean(s.d.)	–	1.03 (1.01–1.06), 0.002	1.03 (1.01–1.06), 0.003
Hidradenitis suppurativa	No	Reference	–
	Yes	1.85 (1.04–3.16), 0.028	–
Immunosuppression	No	Reference	Reference
	Yes	1.92 (0.90–3.82), 0.073	1.95 (0.90–3.93), 0.073
Diabetes mellitus	No	Reference	–
	Yes	1.37 (0.90–2.04), 0.135	–
Abscess position	Abdominal wall—anterior	Reference	–
	Abdominal wall—posterior	0.53 (0.17–1.41), 0.232	–
	Axilla	0.47 (0.19–1.05), 0.075	–
	Buttock	0.51 (0.25–1.02), 0.059	–
	Groin	0.69 (0.35–1.36), 0.282	–
	Lower limb—proximal	1.04 (0.40–2.56), 0.928	–
	Upper limb—proximal	0.75 (0.35–1.56), 0.446	–
	Pilonidal	0.62 (0.34–1.13), 0.116	–
	Thorax—anterior	0.60 (0.26–1.34), 0.228	–
	Thorax—posterior	0.70 (0.38–1.28), 0.241	–
	Other	0.57 (0.21–1.39), 0.237	–
Maximum abscess diameter (cm), mean(s.d.)	–	1.02 (0.96–1.09), 0.477	–
Presence of cellulitis	No	Reference	Reference
	Yes	1.43 (1.01–2.00), 0.042	1.37 (0.96–1.94), 0.083
Timing of hospital presentation	Daytime (08.00–16.59)	Reference	–
	Evening (17.00–21.59)	0.77 (0.48–1.20), 0.265	–
	Night (22.00–07.59)	0.96 (0.54–1.61), 0.881	–
Timing of surgery	First hospital presentation	Reference	–
	Planned return presentation	0.90 (0.63–1.27), 0.560	–
Surgical setting	Operating room	Reference	–
	Emergency department	0.50 (0.03–2.63), 0.506	–
	Ward/treatment area	1.06 (0.74–1.50), 0.762	–
Primary operator	Consultant	Reference	–
	Junior surgical trainee	0.65 (0.33–1.35), 0.221	–
	Nurse practitioner	0.65 (0.22–1.79), 0.412	–
	Senior surgical trainee	0.68 (0.35–1.39), 0.261	–
Abscess packed	No	Reference	–
	Yes	0.80 (0.52–1.24), 0.304	–

Analyses performed on cases without any missing data ($n = 1047$). Junior surgical trainee relates to postgraduate year 1–4 or junior clinical fellow and senior surgical trainee relates to postgraduate year 5–10 or senior clinical fellow.

operating room, dedicated abscess trays were used in 31 hospitals (31 of 54; 57.4%), of which 26 hospitals (26 of 31; 83.9%) used trays holding reusable instruments. For surgery taking place in an ambulatory unit, dedicated abscess trays were used in 11 hospitals (11 of 54; 20.4%), of which only 1 hospital (1 of 11; 9.1%) used trays holding reusable instruments.

Patient characteristics

Of 1631 patients, a small majority were male (871 of 1631; 53.4%) and the most common age group was 26–35 years (351 of 1631; 21.5%). A total of 30.9% were smokers (490 of 1587), 15.6% were diabetic (254 of 1631), and 38.5% had a previous history of subcutaneous abscesses (628 of 1631). At presentation, the most common abscess positions were pilonidal (331 of 1631; 20.3%),

posterior thorax (260 of 1631; 15.9%), and buttock (199 of 1631; 12.2%), with a mean(s.d.) maximum abscess diameter of 4.2(2.3) cm. In half of cases, patients were treated with at least one course of antibiotics before presentation to hospital (867 of 1631; 53.2%) (Table 1).

Treatment characteristics

A total of 1351 patients underwent surgical drainage (1351 of 1631; 82.8%), with the remainder treated with needle aspiration (30 of 1631; 1.8%) or antibiotics alone (210 of 1631; 12.9%). Surgery was performed most commonly in the operating room (922 of 1351; 68.2%) or an ambulatory treatment area (409 of 1351; 30.0%). General anaesthesia was the most common anaesthetic modality (874 of 1351; 64.7%), followed by local anaesthesia (441




Fig. 1 Variation in the use of postoperative antibiotics by centre

The graph depicts the variation in the use of postoperative antibiotics versus the rate of recurrence. Circle size is proportionate to the number of participants included by each centre. The P value, generated using a likelihood ratio test, confirms statistically significant variation in the use of postoperative antibiotics at the centre level (see the Methods section).




Fig. 2 Variation in the use of local anaesthesia by centre

The graph depicts the variation in the use of local anaesthesia versus the rate of recurrence. Circle size is proportionate to the number of participants included by each centre. The P value, generated using a likelihood ratio test, confirms statistically significant variation in the use of local anaesthesia at the centre level (see the Methods section).




Fig. 3 Variation in the use of abscess packing by centre

The graph depicts the variation in the use of abscess packing versus the rate of recurrence. Circle size is proportionate to the number of participants included by each centre. The P value, generated using a likelihood ratio test, confirms statistically significant variation in the use of abscess packing at the centre level (see the Methods section).

of 1351; 32.6%). The majority of abscess cavities were packed (1136 of 1351; 84.2%) and over half of the patients undergoing surgery received a short (less than 3 days) (123 of 1351; 9.1%) or longer (greater than or equal to 3 days) (579 of 1351; 42.9%) course of postoperative antibiotics (Table 2).

Abscess recurrence

Recurrence occurred in 69 of 1351 patients (5.1%) undergoing surgical drainage and 17 of 280 patients (6.1%) undergoing any other treatment approach. For patients undergoing surgery, a higher BMI (OR 1.04, 95% c.i. 1.01 to 1.08; $P=0.015$) and the presence of hidradenitis suppurativa (OR 2.64, 95% c.i. 1.13 to 5.62; $P=0.016$) were associated with a higher rate of recurrence within 60 days. Packing at the end of surgery (OR 0.38, 95% c.i. 0.21 to 0.72; $P=0.003$) was associated with a lower rate of recurrence compared with not packing. Neither the use of local anaesthesia as the primary anaesthetic modality (OR 1.13, 95% c.i. 0.60 to 2.04; $P=0.698$) or withholding postoperative antibiotics (OR 0.97, 95% c.i. 0.53 to 1.74; $P=0.906$) was independently associated with a higher rate of recurrence (Table 3).

Return to hospital

Return to hospital within 60 days occurred in 215 of 1351 patients (15.9%) undergoing surgical drainage, including one patient with

pneumonia (general anaesthetic), one patient with a deep collection (local anaesthetic), one patient with an acute coronary event (local anaesthetic), two patients with diabetic ketoacidosis (both general anaesthetic), and four patients with a haematoma (all general anaesthetic). Return to hospital occurred in 58 of 280 patients (20.7%) undergoing non-operative treatment approaches, including one due to a deep collection. The remainder of returns were due to other planned or non-emergency visits. For patients undergoing surgery, a higher BMI (OR 1.03, 95% c.i. 1.01 to 1.06; $P=0.003$) and a prior history of abscesses (OR 1.45, 95% c.i. 1.04 to 2.01; $P=0.029$) were associated with more returns. Neither the use of local anaesthesia as the primary anaesthetic modality (OR 1.00, 95% c.i. 0.70 to 1.42; $P=0.995$) or withholding postoperative antibiotics (OR 0.99, 95% c.i. 0.70 to 1.41; $P=0.976$) was independently associated with a higher rate of return (Table 4).

Between-centre variation in practice

There was wide variation between centres in the use of postoperative antibiotics (0–100%), local anaesthesia (0–69.6%), and abscess packing (0–100%), as shown in Figs 1–3. To explore this further, multivariable models (comprising only patient-level variables) and multi-level models (comprising patient-level and centre-level variables) were compared and are reported in full in

Table S2. The multi-level models were significantly different from their respective multivariable models for anaesthetic type ($P < 0.001$), postoperative antibiotics ($P < 0.001$), and abscess packing ($P < 0.001$), suggesting a centre-level effect on each aspect of management. Overall, the results showed statistically significant variation in the choice of patient management strategies at the centre level, independent of the variation in case mix.

Discussion

This study reveals wide variation in the management of acute subcutaneous abscesses in the UK. In particular, it reveals disparity in the choice of anaesthesia, the role of cavity packing, and the role of antibiotics. The data challenge the necessity of general anaesthesia as a default approach for surgical drainage, instead highlighting broad opportunities that exist in ambulatory or clinic-based settings with the use of local anaesthesia or other modalities that involve patients being awake. In the context of an observational study, the data show no clear indication of benefit of routine antimicrobial therapy after surgical drainage. The study provides a snapshot of practice specific to the UK, which should guide future research and other local quality improvement initiatives.

There is little evidence to guide the choice of anaesthesia during surgical drainage. A systematic review of six studies published between 2008 and 2014 showed that local and topical preparations appeared safe and effective, but only three of these focused on adult populations⁹. Despite a lack of robust evidence, local anaesthetic preparations are commonly used, as demonstrated by the present data. Indeed, one-third of all drainage procedures were undertaken using local anaesthesia alone and, in some hospitals, this was true for over half of their patients. Possible advantages of local anaesthesia in ambulatory settings include reduced admissions, fewer bed days, improved operating room flow, a quicker return to work and activities for patients, and reduced financial and environmental costs for health systems.

Whilst the use of antibiotics after surgical drainage is not routine in the UK, their use was advocated in a practice recommendation published in 2018⁵. This drew on a systematic review of 14 RCTs, demonstrating modest benefits with oral agents, such as trimethoprim/sulfamethoxazole or clindamycin. The data demonstrated a significant reduction in recurrence at 1 month (OR 0.48, 95% c.i. 0.30 to 0.77 (moderate quality)), as well as a reduction in re-hospitalization (OR 0.55, 95% c.i. 0.32 to 0.94 (moderate quality)). Conversely, the review showed that antibiotics increased gastrointestinal side effects (trimethoprim/sulfamethoxazole: OR 1.28, 95% c.i. 1.04 to 1.58 (moderate quality)); and clindamycin: OR 2.29, 95% c.i. 1.35 to 3.88 (high quality)), including the incidence of diarrhoea⁶. In the present study, approximately half of all patients received antibiotics after surgery, with many receiving antibiotics for greater than or equal to 3 days. The modest benefits conferred by these antibiotics should be weighed carefully against potential harms of side effects and the societal impact of antimicrobial resistance⁵.

Abscess cavity packing is undertaken to prevent premature closure of the incision and, in turn, to reduce the risk of recurrence. The present data reveal a significant association between packing and reduced recurrence at 60 days. This contrasts with the results of other high-quality studies, such as the PPAC2 RCT, which demonstrated no difference in recurrence rates along with improved pain scores for patients whose

perianal abscess cavity was not packed¹⁰. The results of the present study should be interpreted carefully, as no data are available on the duration of packing and whether this was done solely to achieve haemostasis after surgery. The data should, however, discourage extrapolation of previous results drawn from different populations and disease groups. The need for packing should remain a consideration after surgical drainage of subcutaneous non-perianal abscesses.

Limitations of the study are recognized. First, as an observational study of routinely collected data, it is likely that some events were missed, leading to under-reporting of key outcomes. This may have occurred if, for example, patients re-presented to community settings, alternative hospitals, or not at all¹¹. Second, the study was unable to explore other outcomes that are probably important for patients and healthcare systems, such as pain, quality of life, cosmesis, and health economics. These remain an important target for future research. Third, owing to limitations of follow-up procedures, some return-to-hospital events may have been planned clinical reviews in hospital, particularly for the non-operative group; it was not possible to delineate these from emergency returns.

It is envisaged that the present data will initiate local discussions between relevant stakeholders around the management of subcutaneous abscesses. The role of local anaesthesia is a key consideration, particularly the opportunity it may present to expand ambulatory pathways and to reduce the environmental impact of the operating room. The role of antibiotics after surgical drainage is also a key consideration. The data should initiate local discussions between surgical and microbiology stakeholders with respect to the balance between treatment outcomes, side effects, and the risk of antimicrobial resistance. Local quality improvement initiatives are encouraged and the present data provide a baseline to guide such activities in the future.

Collaborators

White Rose Surgical Collaborative and MAGIC Collaborators

S. J. Chapman, J.R. Burke, J. Helliwell, M. Kowal, K. Boag, M. Ayres, L. McMenemy, T. Ho, J. Carr-Wilkinson, A. Peckham-Cooper (Leeds Teaching Hospitals NHS Trust); J. Spackman (Chesterfield Royal Hospital NHS Foundation Trust); A. Rashid, K. Eng, V. Boodhoo (Sheffield Teaching Hospitals NHS Foundation Trust); T. Williams (Hull York Medical School); J. Bundred (Leeds Teaching Hospitals NHS Trust); A. Nessa, G. Ramsay (Aberdeen Royal Infirmary); G. Greenlees, L. Whittaker (Airedale General Hospital); N. Pinsker, J. Phillips (Barnsley Hospital); M. Abu Sayed, A. Kausar (Royal Blackburn Teaching Hospital); J. Spackman, R. Ardley (Chesterfield Royal Hospital NHS Foundation Trust); A. Rees, D. Vimalachandran (Countess of Chester Hospital); L. Henderson, D. Watt (Crosshouse Hospital); V. Miu, R. George (Doncaster Royal Infirmary); C. English, B. Stubbs (Dorset Country Hospital); O. Adepoju, L. Nicol (Dr Gray's Hospital); L. Rampersad, M.S.J. Wilson (Forth Valley Royal Hospital); M. Chauhan, S. Shirol (Frimley Park Hospital); M. Abdelkarim, V. Ochieng (Glan Clwyd Hospital); A. Golder, D. Mansouri (Glasgow Royal Infirmary); D. Liu, T. Walker (Gloucestershire Hospitals NHS Foundation Trust); M. El Boghdady, K. Adams (Guys and St. Thomas Hospital); H. Standing, C. McNaught (Harrogate District Hospital); W. Ramadan, S. Harris (The Hillingdon Hospitals NHS Foundation Trust); A. Ghoneima, A. Saha (Huddersfield Royal Infirmary); X. Y. Choo, M. Alwahid (Inverclyde Royal Hospital);

G. Karagiannidis, J. Pitt (Ipswich Hospital NHS Trust); G. James, V. Kanakala (James Cook University Hospital); B. Down, G. Bond-Smith (John Radcliffe Hospital); S. Lamidi, A. Raicu (Kettering General Hospital); A. Hamadani, H. Shanti (Kings College Hospital); D. Sapre, E. Gemmill (Kings Mill Hospital); W. Y. Chung, L. B. Samarakoon (Leicester Royal Infirmary); S. Ingram-Walpole, C. Macutkiewicz (Manchester Royal Infirmary); S. Oo, R. Harries (Morrison Hospital); I. El-Kharbotly, G. Bashir (Newham University Hospital); A. Adesuyi, A. Arain (North Devon District Hospital); N. Abbud, N. Shah (Northern General Hospital); A. Shalaby, B. Appleton (Princess of Wales Hospital); B. Dolaszynska (Queen Elizabeth Hospital, Gateshead); N. Fazili, D. Naumann (Queen Elizabeth Hospital Birmingham); G. Chillarge, N. Smart (Royal Devon and Exeter Hospital); R. Barnett, M. Mulla (Royal Glamorgan Hospital); T. Pelly, K. Hancorn (Royal London Hospital); J. Honey, G. Branagan (Salisbury District General Hospital); C. Ezeme, R. Newton (St. Richard's Hospital); S. Jones, A. Peckham-Cooper (Leeds Teaching Hospitals NHS Trust); S. Hodge, S. Mehmood (Stepping Hill Hospital); M. Bhardwaj, A. Liaqat (Stoke Mandeville Hospital); S. Wallace-King, W. Carr (Sunderland Royal Hospital); A. Rahman, K. Carswell (Surrey and Sussex Healthcare NHS Trust); A. S. Opaluwa, S. Hyder (The Queen Elizabeth NHS Foundation Trust, Kings Lynn); Y. Akula, S. Robertson (University Hospitals Coventry and Warwickshire); C. J. Fairfield, F. Maxwell (University Hospital Wishaw); S. Abdelrahman, S. Abbott (Walsall Manor Hospital); H. Fowler, N. Pranesh (Warrington and Halton Teaching Hospitals NHS Foundation Trust); B. Manku, A. Baker (Warwick Hospital); M. Mehta, S. Shirol (Wexham Park Hospital); E. Wates, R. Fernandes R (William Harvey Hospital); M. I. K. Malik, S. Duff (Wythenshawe Hospital); N. Lawrie, G. Neelankavil Davis, D. Dorkeh (Aberdeen Royal Infirmary); E. Moran (Airedale General Hospital); N. Aruparayil, H. Allott, J. Creed, K. Lehovsky (Barnsley Hospital); A. Sallam, C. Abdulrahman, S. Puranik, P. Suwichacherdchu (Royal Blackburn Teaching Hospital); O. M. O'Connor, S. Kaul, R. Ghosh (Chesterfield Royal Hospital NHS Foundation Trust); B. Davies (Countess of Chester Hospital); L. Mamer, E. McCarron (Crosshouse Hospital); H. Pendergast, J. Corkerry, M. Dowiedar (Doncaster Royal Infirmary); T. Knowles, K. Tin, L. Demicoli (Dorset Country Hospital); P. Osei-Bonsu, J. Balfour, A. Varma (Dr Gray's Hospital); J. Sikora (Forth Valley Royal Hospital); P. Belaguthi, R. Lee, L. Berger, T. Baron, S. Myat, C. Rowlands, O. Oyekan, L. Irabor Ezoba, S. Tontus (Frimley Park Hospital); T. Panos, N. Usmani, A. Abdelhamid, S. Kamil, A. Maraqa, W. Chantima, M. Dyer, K. Thane, S. Yap (Glan Clwyd Hospital); S. Tompkins, A. Asurlekar, P. Polz, B. Clark, D. Low, K. Robertson (Glasgow Royal Infirmary); O. Hamid (Gloucestershire Hospitals NHS Foundation Trust) M. Pipe, N. Gadiyar (Guys and St. Thomas Hospital); Z. L. Hoo, Y. W. Wong (Harrogate District Hospital); F. Sheikh, V. Satish (The Hillingdon Hospitals NHS Foundation Trust); M. Kronberga, S. Suresan, M. Kattakayam, A. Ed Sabbagh, D. Saleh, H. Akhtar, S. Gurung, S. Bakare (Huddersfield Royal Infirmary); E. Ross, M. Shahid, A. Tweedie, C. M. Moen (Inverclyde Royal Hospital); M. El-Farran, T. Khaliq (Ipswich Hospital NHS Trust); B. Peter (James Cook University Hospital); M. Shams, J. Fung, T. Theivendrapillai, J. Bundred, A. Ang, R. Ahmed (John Radcliffe Hospital); A. Sherafat, K. Annamalai, A. Ali (Kettering General Hospital); V. Kalogianni, A. Hassan (Kings College Hospital); W. Aziz, B. Srikumar, N. Minhas, R. Parks, S. Hanif, A. Farooq, M. Salem, A. Ali (Kings Mill Hospital); G. Davalos (Leicester Royal Infirmary); A. Hardy, Y. Zhang (Manchester Royal Infirmary); K. Ngai, H. Baker,

J. Ling-Soffe, M. Mcmillan, D. Thompson (Morrison Hospital); A. Dkhakhni, D. Sapountzis (Newham University Hospital); A. Li, H. Al-Siraj (North Devon District Hospital); S. Khan, A. Kuligowska, I. Ogunwonyi, R. Broadbent, H. George, A. Rashid (Northern General Hospital); I. Zarad, M. Osman, M. Abdelsalam (Princess of Wales Hospital); A. Gupta (Queen Elizabeth Hospital, Gateshead); S. Moodalbyle, C. Lewis (Queen Elizabeth Hospital Birmingham); O. Barrington M, Shams (Royal Devon and Exeter Hospital); C. Jayasekara, K. Lam (Royal Glamorgan Hospital); R. Hwang, L. Xu, A. Ishak (Royal London Hospital); W. Protheroe, T. Stylianou (Salisbury District General Hospital); E. O. Oladeji, H. Anuff, E. Husain (St. Richard's Hospital); J. Helliwell, J. Holmes, N. Shatwell (Leeds Teaching Hospitals NHS Trust); M. Kishima M (Stepping Hill Hospital); F. Issa, R. Vashisht, G. Bhardwaj (Stoke Mandeville Hospital); A. Thaker, F. Bandari (Sunderland Royal Hospital); B. Robertson-Jones, M. Mostafa, T. Nahiyani (Surrey and Sussex Healthcare NHS Trust); N. VinodKumar, B. Altaf, A. Sharma, O. Khalil, A. Mensah, E. Ozekhome-Mike (The Queen Elizabeth NHS Foundation Trust, Kings Lynn); F. Ghafur, S. Edensor, W. Hodges (University Hospitals Coventry and Warwickshire); A. Omran, W. H. Younas, Y. Harris (University Hospital Wishaw); E. Spurring, F. Osunlusi, C. Okoronkwo, S. Shah, E. L. Yan, R. Hajdok (Walsall Manor Hospital); C. Anderton, F. Costigan, H. Selvachandran, B. David (Warrington and Halton Teaching Hospitals NHS Foundation Trust); A. O'Sullivan, C. Spreafico, M. Phillips (Warwick Hospital); S. Arora, A. Ahmed, M. Ghotmi, R. Mohamed (Wexham Park Hospital); M. Kawabata, A. Gardner, I. Atherley, A. Chowdhury, J. Kalliyath (William Harvey Hospital); K. Manor, M. Bamgboye, L. Walker, J. Law, E. Reynolds, Z. Bholah, R. Castelhao (Wythenshawe Hospital).

Funding

The authors have no funding to declare.

Acknowledgements

The authors are grateful to Catherine Moriarty (Leeds Teaching Hospitals NHS Trust) for providing access to and support with the Research Electronic Data Capture (REDCap) platform during data collection. The authors are grateful to the Association of Surgeons of Great Britain and Ireland (ASGBI), the North West Research Collaborative (NWRC), and the CRAMSurg group for disseminating the study protocol to participating sites. The authors are also grateful to the National Institute of Health and Care Research Surgical MedTech Co-operative for their assistance with disseminating the findings of this work.

Author contributions

S. J. Chapman and J. R. Burke are responsible for idea conception. All collaborators participated in data collection. Local Leads coordinated the study with oversight from the Steering Committee. The Writing Group drafted and finalized the article and all individuals agreed to its submission. The corresponding author attests that all listed collaborators meet required criteria. J. Bundred is the statistical guarantor. S. J. Chapman and J. R. Burke are the overall guarantors

Disclosure

The authors declare no conflict of interest.

Supplementary material

Supplementary material is available at BJS online.

Data availability

Requests for data sharing will be considered by the study management group upon receipt of a statistical analysis plan.

References

1. Moran GJ, Krishnadasan A, Gorwitz RJ, Fosheim GE, McDougal LK, Carey RB *et al.* Methicillin-resistant *S. aureus* infections among patients in the emergency department. *N Engl J Med* 2006;**355**:666–674
2. Shallcross LJ, Hayward AC, Johnson AM, Petersen I. Evidence for increasing severity of community-onset boils and abscesses in UK general practice. *Epidemiol Infect* 2015;**143**: 2426–2429
3. Thomas O, Ramsay A, Yiasemidou M, Hardie C, Ashmore D, Macklin C *et al.* The surgical management of cutaneous abscesses: a UK cross-sectional survey. *Ann Med Surg (Lond)* 2020;**60**:654–659
4. Elkadi HH, Dodd E, Poulton T, Bolton W, Burke JR, Tiernan J *et al.* The effect of anaesthetic use on healing in sub cutaneous abscess management: a retrospective before and after cohort study. *Ann R Coll Surg Engl* 2023;**105**:241–246
5. Vermandere M, Aertgeerts B, Agoritsas T, Liu C, Burgers J, Merglen A *et al.* Antibiotics after incision and drainage for uncomplicated skin abscesses: a clinical practice guideline. *BMJ* 2018;**360**:k243
6. Wang W, Chen W, Liu Y, Siemieniuk RAC, Li L, Martínez JPD *et al.* Antibiotics for uncomplicated skin abscesses: systematic review and network meta-analysis. *BMJ Open* 2018;**8**:e020991
7. Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic data capture (REDCap)—a metadata-driven methodology and workflow process for providing translational research informatics support. *J Biomed Inform* 2009;**42**:377–381
8. von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP. Strengthening the reporting of observational studies in epidemiology (STROBE) statement: guidelines for reporting observational studies. *BMJ* 2007;**335**:806–808
9. Poulton T, Hogston SP, Bolton WS, Burke JR, Jayne DG. Anaesthetic management of subcutaneous abscesses: current status. *Br J Anaesth* 2020;**125**:e227–e229
10. Newton K, Dumville J, Briggs M, Law J, Martin J, Pearce L *et al.* Postoperative Packing of Perianal Abscess Cavities (PPAC2): randomized clinical trial. *Br J Surg* 2022;**109**:951–957
11. Wounds UK. *Best Practice Statement: Post-Operative Wound Care—Reducing the Risk of Surgical Site Infection*. London: Wounds UK, 2020