

1 **Title page**

2 Title: Identifying modifiable factors that influence walking in patients undergoing surgery for  
3 neurogenic claudication. A prospective longitudinal study

4

5 Authors:

6 Suzanne McIlroy<sup>1, 2\*</sup>, Lindsay Bearne<sup>3</sup>, John Weinman<sup>4</sup>, Sam Norton<sup>2</sup>

7 Affiliations:

8 1 Physiotherapy Department, King's College Hospital, London, UK.

9 2 Health Psychology Section, King's College London, London, UK

10 3 Population Health Research Institute, City St George's, University of London, London, UK

11 4 School of cancer and pharmaceutical science, King's College London, UK

12

13 Corresponding author: [Suzanne.mcilroy@nhs.net](mailto:Suzanne.mcilroy@nhs.net)

14

1 **Title:** Identifying modifiable factors that influence walking in patients undergoing surgery for  
2 neurogenic claudication. A prospective longitudinal study

3 **Abstract**

4 Neurogenic claudication, caused by lumbar spinal stenosis, is the most common reason for spinal  
5 surgery in older adults, aiming to improve pain and walking. However, most people do not increase  
6 walking post-operatively. This study aimed to identify modifiable physical and psychosocial factors  
7 that could be targeted with rehabilitation. A prospective longitudinal study recruited 97 adults, aged  
8 >50 years, awaiting surgery for neurogenic claudication. Walking measures (six-minute walk test,  
9 daily step count, self-rated maximum walking distance) were assessed pre-surgery and 12-weeks  
10 post-surgery. Modifiable variables, mapped to a behaviour change model (COM-B; e.g. falls, lower  
11 limb performance, fear of movement, illness perceptions), were evaluated using mixed-effects  
12 regression models. All walking measures demonstrated statistically significant improvements  
13 ( $p < .001$ ). However, 50% did not achieve minimum clinically important differences. The strongest  
14 correlation with post-operative walking was pre-operative walking. Cross-sectionally, lower limb  
15 performance (b: .75; 95%CI .64, .86 to b: .35; 95%CI .19, .52), pre-surgery history of falls (b: -.29; 95%CI  
16 -.44, -.13), fear of falling (b: -.55; 95%CI -.69, -.41 to b: -.32; 95%CI -.48, -.15), fear of movement (b: -.48;  
17 95%CI -.63, -.33 to -.22; 95%CI -.40, -.03), coherence of condition (b: -.23; 95%CI -.41, -.05 to b: -.17;  
18 95%CI -.33, -.01) and perceived personal control (b: .26; 95%CI .09, .43 to b: .14; 95%CI .02, .31), were  
19 significantly associated with pre-surgical walking ( $p < .05$ ). Most pre-surgical variables were not  
20 longitudinally associated with change in walking post-surgery. Six-weeks post-surgery fear of falling  
21 (b: -.35; 95%CI -.57, -.13 to b: -.18; 95%CI -.33, -.02), fear of movement (b: -.32; 95%CI -.53, -.11 to b: -.19;  
22 95%CI -.33, -.05), and emotional response (b: -.24; 95%CI -.38, -.11 to b: -.22; 95%CI -.41, -.03) were  
23 significantly associated with less improvement in 12-weeks. Prehabilitation and post-operative  
24 rehabilitation targeting walking, balance, and psychosocial factors is recommended to optimise post-  
25 surgical walking.

26

27 **Keywords:**

28 Neurogenic claudication; Walking; Rehabilitation; Fear of movement; Fear of falling; Prognosis

29

## 1 Introduction

2 Neurogenic claudication (NC) affects approximately 10% of the general population with incidence  
3 increasing with age<sup>1,2</sup>. It is caused by lumbar spinal stenosis (LSS), a degenerative condition that  
4 leads to narrowing around and compression of the nerves and blood vessels within the lumbar  
5 spine. NC is characterised by bilateral leg pain, paraesthesia, and/or weakness often accompanied by  
6 low back pain. It is exacerbated by standing and can cause substantial walking restriction<sup>3</sup>, greater  
7 than that experienced by people with hip or knee osteoarthritis<sup>4</sup>. Additionally, it can lead to a  
8 substantial reduction in quality of life, comparable to that caused by stroke and heart disease<sup>5</sup>.

9 Neurogenic claudication is the most common cause for lumbar surgery in older adults and the  
10 number of procedures performed annually is increasing worldwide<sup>6</sup>. Surgery aims to reduce pain and  
11 improve walking<sup>7-9</sup>. Yet, after surgery for NC, approximately 40% of people have ongoing pain and  
12 walking restriction<sup>10</sup> and the majority of people do not increase their daily step count and physical  
13 activity<sup>11,12</sup>. Walking is an accessible and acceptable form of physical activity in older people<sup>13</sup> and is  
14 associated with many health benefits including greater function and lower morbidity and mortality<sup>14</sup>.  
15 Thus, if people post-surgery are not increasing their walking, they remain at risk of the consequences  
16 of inactivity.

17 Identifying the factors associated with post-operative walking in patients undergoing surgery for NC  
18 is important because they could be targeted with rehabilitation to improve outcomes. To date, most  
19 of the research has focused on routinely collected and biological factors, for example sex, age, body  
20 mass index (BMI), comorbidities, and radiological parameters<sup>15-19</sup>, as potential predictor variables  
21 whereas there has been limited investigation into the modifiable psychosocial variables of walking  
22 behaviour<sup>20</sup>. A systematic review of pre-operative factors (n=34 studies, 9,973 patients) identified  
23 moderate quality of evidence that pre-operative walking capacity was positively associated with  
24 post-operative walking capacity; and spondylolisthesis and severity of stenosis was not found to be  
25 associated with post-operative walking capacity<sup>20</sup>. There was weak or inconclusive evidence that  
26 other factors were associated with post-operative walking: higher BMI, smoking, and previous  
27 lumbar surgery were negative prognostic factors; and higher income and better self-rated health  
28 were associated with better outcomes. However, age, sex, symptom severity and duration, fear  
29 avoidance, and social support were not found to be prognostic factors. Conflicting evidence  
30 emerged regarding comorbidities; musculoskeletal conditions and diabetes were identified as  
31 negative prognostic factors, while cardiovascular and respiratory comorbidities did not appear to  
32 influence post-operative walking capacity<sup>20</sup> Crucially, other than one study investigating fear

1 avoidance beliefs<sup>21</sup>, modifiable psychosocial variables of behaviour that may be addressed with  
2 rehabilitation, had not been studied.

3 Factors in the early post-operative period may also be important variables associated with long-term  
4 outcome. Studies in mixed lumbar-surgical populations have identified factors such as pain self-  
5 efficacy, fear of movement, illness perceptions and social support as being important<sup>22-25</sup>. However,  
6 these have not been thoroughly investigated in patients with NC.

7 Walking is a complex, multifaceted behaviour<sup>26</sup>, behaviour change models can be used to  
8 comprehensively understand the behaviour and identify barriers and enabling factors, particularly  
9 those that might be most amenable to intervention. The Behaviour Change Wheel (BCW), a  
10 synthesis of 19 behaviour change models<sup>27</sup>, provides a systematic framework for understanding the  
11 enabling factors of a target behaviour (e.g. walking) which can then be used to design an  
12 intervention (e.g. rehabilitation programme) targeting these factors. At the core of the BCW is the  
13 COM-B model. This specifies three drivers of behaviour: Capability (physical and psychological),  
14 Opportunity (social and physical), and Motivation (reflective and automatic). It further theorises that  
15 both Capability and Opportunity influence Motivation, making Motivation the central mediator  
16 within the COM-B model. The comprehensive coverage of the COM-B allows researchers to analyse  
17 the salient determinants specific to the population and behaviour of interest. Identifying the salient  
18 determinants is a crucial step in developing interventions and the COM-B and the BCW have been  
19 successfully utilised to develop interventions to increase physical activity in other populations<sup>28,29</sup>,  
20 demonstrating its practical utility.

21 To date, no study has used a behavioural model to investigate the modifiable factors associated with  
22 walking in people undergoing surgery for NC. The aims of this study were to evaluate change in  
23 walking capacity and performance from pre-surgery to 12-weeks post-surgery; and to identify  
24 potentially modifiable physical and psychosocial determinants of walking capacity and performance  
25 in people undergoing surgery for NC that could be targeted with rehabilitation.

## 26 Materials and methods

27 Study design: A prospective multi-site observational study was conducted. The *Strengthening the*  
28 *Observational Report on Epidemiology* (STROBE) guidelines<sup>30</sup> were used to inform design and  
29 reporting of the study. The study was performed in accordance with the Declaration of Helsinki,  
30 written informed consent was provided by all participants. Ethical approval was obtained from the  
31 East Midlands - Nottingham 1 Research Ethics Committee (20/EM/0307). The study protocol was  
32 registered at Open Science Framework (DOI 10.17605/OSF.IO/BHQJZ).

1 Participants: Participants were recruited from three NHS hospital trusts in England. Participants were  
2 eligible if they were  $\geq 50$  years old and due to have decompressive surgery for degenerative LSS with  
3 symptoms of NC. Neurogenic claudication symptoms were defined as leg or buttock pain and/or  
4 tingling, numbness or heaviness, made worse when standing or walking, and/or eased by sitting or  
5 bending forward. Symptoms may be present with or without low back pain<sup>3,31</sup>. Exclusion criteria  
6 consisted of LSS caused by tumour, fracture or significant deformity ( $>15^\circ$  lumbar scoliosis;  $\geq$  grade II  
7 spondylolisthesis); patients requiring emergency surgery;  $>1$  level fusion surgery; or if they reported  
8 other conditions that were the primary cause of walking restriction. Patients with less than a week  
9 before their scheduled date for surgery were excluded as there was insufficient time for baseline  
10 data collection. Conversational level English or willingness to use an interpreter was also required.

11 The target sample size was set at 122 participants and calculated based on the objective to  
12 determine the factors that are associated with 12-week six-minute walk distance (6MWD) after  
13 controlling for clinical-demographic confounding variables. Specifically, 97 participants providing  
14 80% power ( $\alpha=.05$ ) to detect a continuous predictor variable that explained an additional 5% of  
15 the variance in a linear regression model, including five control variables that combined explained  
16 35% of the variance (i.e.  $R^2_{\text{change}}=.05$ , equivalent to  $d=.5$ ). This number was inflated to account for  
17 20% attrition resulting in the target of 122.

18 Procedures: Potentially eligible participants were invited to find out more about the study either in  
19 person during their surgical clinic appointment or via the telephone. They were invited to attend  
20 two, one hour assessments, one prior to their surgery and one 12-weeks following their surgery. If  
21 interested, they were posted the participant information sheet, consent form and a paper-based  
22 baseline questionnaire pack consisting of questions to collect demographic information and the self-  
23 reported measures. At the assessments, the questionnaire packs were checked for completion and  
24 the objective measures completed. Participants also completed a self-reported questionnaire pack  
25 at six weeks post-surgery which was returned by post.

26 Tests and measures:

27 Walking capacity and performance: Walking capacity was assessed using the six-minute walk test.  
28 Participants were asked to walk as far as possible around two cones, placed 10m apart in a straight  
29 corridor, in 6 minutes. The total distance walked (in metres) in 6 minutes was recorded. The six-  
30 minute walk test is reliable and responsive to change in older people with long-term conditions<sup>32</sup>.  
31 The minimal clinically important difference (MCID) for the 6MWD has been calculated to be 50m<sup>33</sup>.  
32 Walking performance was measured in mean steps/day using a valid and reliable triaxial  
33 accelerometer (ActivPal3™, PAL Technologies Ltd., Glasgow, UK) which uses information about

1 acceleration and thigh position to determine body posture, stepping, and cadence<sup>4,12</sup>. Each  
2 participant was fitted with an accelerometer on the mid-anterior thigh with a waterproof dressing.  
3 Initiation of the recording started at midnight of the day of assessment and participants were  
4 requested to wear it continuously for the following seven days. Three-dimensional acceleration data  
5 were collected over 60 second epochs. Accelerometer data with at least 14 hours of wear time per  
6 24-hour period and a minimum of 5 days of wear time was considered valid<sup>34</sup>. A minimum of 7000  
7 steps/day is recommended for older adults to achieve health benefits<sup>11,35</sup>. The MCID for NC has not  
8 been defined, in its absence we have utilised the MCID for people with peripheral artery disease:  
9 558 steps<sup>36</sup>. Maximum self-rated walking distance was assessed by asking “what is the maximum  
10 distance (in meters) you can walk at your usual pace on a flat surface before you have to stop?”<sup>37</sup>.  
11 No MCID has been defined, for the purpose of this study we set it at 250m. This was based on the  
12 MCID of the self-paced walking test which measures how far people with NC can walk for  
13 ≤30minutes before requiring a rest<sup>38</sup>.

14 Clinical and demographic details: Clinical and demographic details collected included: age, body  
15 mass index (kg/m<sup>2</sup>), sex, ethnicity, education, employment, indices of deprivation, social support,  
16 co-morbidities, smoking history, falls history. We collected the Oswestry Disability Index (ODI)<sup>39</sup> to  
17 assess back-pain related disability. The EuroQol five dimension, 5-level questionnaire (EQ5D-5L)<sup>40</sup>  
18 was used to assess quality of life. Average severity of back pain and average leg pain when resting  
19 and when walking over the last week was collected using an 11-item numerical rating scale<sup>41</sup>.

20 Candidate predictor variables: Candidate predictor variables were selected by mapping measures  
21 and constructs onto the COM-framework<sup>27</sup>. There is no standardised method or measurement tool  
22 to capture all the components of the COM-B therefore, to ensure we considered a range of  
23 potentially modifiable factors across the domains of the framework, that may be suitable to target in  
24 future rehabilitation. We considered variables that had weak or inconclusive evidence from the  
25 previous systematic review e.g. fear of movement<sup>20</sup>, factors associated with walking in older people  
26 e.g. balance<sup>42,43</sup> and other conditions resulting in walking restriction e.g. illness perceptions in  
27 peripheral artery disease<sup>44</sup>. We included at least one measure for each of the six COM-B  
28 components, balancing the need for comprehensive assessment with minimising participant burden,  
29 sample size requirements, and risks of multiple testing. This approach ensured that the selected  
30 measures were both scientifically robust and feasible within the constraints of the study design.

31 *Walking capacity (physical and psychological skills or knowledge required to perform the behaviour):*  
32 Lower limb performance was measured with the Short Physical Performance Battery. This valid and  
33 reliable measure comprises three standing balance tests, gait speed over 2.44m and time to

1 complete 5 sit-stands. A performance score between 0-12 is calculated and higher scores indicate  
2 better performance<sup>45</sup>. Maximum grip strength (kg) was assessed using a hand held Jamar Plus+  
3 Dynamometer<sup>46</sup>. Grip strength is widely used in the elderly to assess strength, lower scores are is  
4 associated with sarcopenia and frailty<sup>46</sup>. Participants perception that their condition is  
5 understandable, meaningful, and manageable was assessed using the coherence question from the  
6 Brief Illness Perceptions Questionnaire (B-IPQ). The BIPQ is a reliable and valid nine-item scale used  
7 to assess the cognitive and emotional representations of illness<sup>47</sup>. Ability to plan exercise was  
8 assessed using the action planning domain of the validated Self-Regulation questionnaire<sup>48</sup>. Higher  
9 scores indicated better ability to plan exercise.

10 *Walking opportunity (social and physical environments that may enable or constrain the behaviour):*  
11 Physical environment and suitability for walking was assessed using the valid and reliable self-  
12 reported physical activity-related environmental factors (ALPHA) scale<sup>49</sup>. A higher score indicates  
13 higher walkability. The social support and exercise survey was used to assess how much support  
14 family and friends provide to exercise<sup>50,51</sup>. Total scores range between -16 and 88 with higher scores  
15 indicating greater support.

16 *Walking motivation (reflective and automatic processes driving the behaviour):* Patient health  
17 questionnaire 4 (PHQ4) was used as an ultra-brief screening tool for depression and anxiety<sup>52</sup>. It  
18 requires participants to rate how often they have been bothered by thoughts and feelings on 4-  
19 items on a four-point Likert-type scale. Higher scores indicate greater severity. Beliefs about  
20 consequences, timeline, personal control, treatment control, identity, emotional representation,  
21 and illness concern were assessed with the B-IPQ<sup>47</sup>. Fear of falling was assessed using the valid and  
22 reliable Short Falls Efficacy Scale International (SFESI)<sup>53</sup>. A higher score indicates greater fear of  
23 falling. Fear of movement was assessed with the valid and reliable Tampa Scale of Kinesiophobia<sup>54,55</sup>.  
24 The total score of the scale range from 17- 68. A higher score indicates greater fear of movement.  
25 Ability to monitor and regulate exercise behaviour was assessed using the action control domain of  
26 the Self-Regulation questionnaire<sup>48</sup>. Higher scores indicated better ability to regulate behaviour.

27 Analysis:

28 Descriptive statistics were used to summarise participants' demographic and clinical characteristics.  
29 Means and standard deviations, or medians and interquartile ranges (IQR) for continuous variables  
30 are reported depending on skew; and frequencies and percentages for categorical variables. To  
31 compare walking capacity and performance and change in clinical characteristics pre- to post-  
32 operatively, we conducted two-tailed, paired t-tests. To determine factors associated with post-  
33 surgical walking capacity and performance, we assessed bivariate correlations and estimated

1 multivariable linear mixed-effects regression coefficients, adjusted for key putative confounders.  
2 Regression models were estimated separately for each candidate predictor variable with walking  
3 measures at each post-surgery assessment as the outcome. A random-intercept accounted for the  
4 repeated measures nature of the data and a predictor-by-time interaction term was estimated to  
5 allow the estimation of different effects at each assessment. Models, controlled for age, gender,  
6 ethnicity, obesity, smoking and baseline score of the outcome. The maximum likelihood estimation  
7 approach allowed for the inclusion of all participants with at least one post-surgery assessment of  
8 the outcome, under the *missing at random* assumption. Predictor variables were the baseline scores  
9 of the COM-B domain factors described above plus the change from baseline to 12-weeks where this  
10 was recorded. To aid comparisons across predictor variables, standardised regression coefficients  
11 with 95% confidence intervals were estimated and presented in a forest plot. Given the exploratory  
12 nature of the study, statistical significance was set at 0.05 and no adjustment for multiple testing  
13 was made. Magnitude of regression coefficients was defined as strong  $\geq .5$ ; moderate  $\geq .3$ ; and weak  
14  $\geq .1$ <sup>56</sup>. This approach combined with the consideration of the distribution of the candidate predictor,  
15 to ensure they do not have floor/ceiling effects and are thus potentially modifiable, is recommended  
16 for selecting target variables for behavioural interventions<sup>57</sup>. All statistical analyses were conducted  
17 using STATA version 17 (Stata Corp, College Station Texas, USA).

## 18 Results

19 Between April 2021 and July 2022, 288 patients were screened for eligibility. Of these, 221 met the  
20 inclusion criteria and 134 consented and underwent baseline assessment. Mean (SD) age was 70.2  
21 (8.6) years, and 69 (51.5%) were female. As 17 people did not have surgery, in total 117 participants  
22 were recruited (Figure 1, study flow). This was slightly below the target of 122, but as attrition was  
23 lower than anticipated, power remained at 80%. In total, 109 participants (93%) completed 12-week  
24 follow-up assessments, although objective measures were only collected on 97 (84%) participants.  
25 This was due to concerns about attending hospitals due to the perceived risk of COVID-19 infection  
26 or transport issues.

27 Table 1 presents baseline demographics and clinical details on all participants. The most common  
28 surgical procedure was laminectomy (n=91), and most participants had one level operated on  
29 (n=80). Median post-operative length of stay was one night (IQR 1 to 2; range 0-31). Thirteen (11%)  
30 had a hospital recorded complication (post-operative haematoma and required further surgery=3,  
31 dural tear=2; wound ooze=2; acute urinary retention=1, urinary infection=1, delirium=1, heart failure  
32 one-month post-operative=1, COVID-19=1, hospital acquired pneumonia=1). Twenty three (20%)



1 received peri-operative (inpatient) physiotherapy and 43 (37%) received post-operative  
2 physiotherapy following discharge.

3 Change in walking measures after surgery:

4 Table 2 and figure 2 illustrate the walking capacity and performance measures at baseline and  
5 follow-up. Changes in disability, quality of life and pain measures are presented in Supplementary  
6 Material 1. All walking measures demonstrated a statistically significant change post-operatively:  
7 mean increase in 6MWD was 61.38m ( $\pm 72.57$ ) and daily step count 582.31 ( $\pm 1720.3$ ), and median  
8 increase in maximum walking distance was 400m (IQR 15 to 1200) (all  $p < .001$ ). Percentage achieving  
9 MCID were 49% for 6MWD; 58% for maximum walking distance; and 40% for daily step count. There  
10 was no statistically significant difference between the number of patients walking  $\geq 7000$  steps/day  
11 at baseline and follow-up assessment ( $p = .250$ ).

12 Factors associated with change in walking:

13 The pre-operative walking score for each of the walking measures explained 67% of the variance of  
14 the residualised change score in walking improvement for the 6MWD ( $R^2 = .674$ ) and 75% of the  
15 variance in step count ( $R^2 = .753$ ) and 38% of the variance of the self-rated maximum walking distance  
16 ( $R^2 = .380$ ) (Supplementary Material 2).

17 The pre-operative and post-operative factors associated with pre-operative walking and change in  
18 post-operative walking are illustrated in the forest plots (figure 3) and tables 3-6. The unadjusted  
19 correlations between the candidate predictor variables and walking measures are reported in  
20 Supplementary Material 3.

21 Walking capability:

22 In the analysis of the pre-operative capability factors, all the pre-operative capability variables were  
23 significantly associated cross sectionally with at least one of the pre-operative walking measures  
24 ( $p < .05$ , table 3). Specifically, all variables were significantly associated with pre-operative 6MWD and  
25 number of falls, ability to plan exercise, and lower limb physical performance were significantly  
26 associated with all three walking measures. Lower limb physical performance had the strongest  
27 cross-sectional association with pre-operative walking measures, with a moderate to strong  
28 association. Pre-operative ability to plan exercise, and pre-operative grip strength were positively  
29 but weakly associated with all pre-operative walking measures. Pre-operative coherence and history  
30 of a fall had weak negative association with pre-operative 6MWD.

1 None of the pre-operative variables were significantly longitudinally associated with change in  
2 walking at 12-weeks post-operative (table 3). Of the 6-weeks post-operative capability measures,  
3 only history of a fall post-operatively was statistically significant associated with change in walking at  
4 12-weeks although the effect size was weak (table 6).

5 Walking opportunity:

6 In the analysis of the pre-operative opportunity factors, both social support to exercise, and physical  
7 environment suitability for walking, were significantly, yet weakly, positively associated cross  
8 sectionally with the pre-operative 6MWD but not the pre-operative step count or maximum walking  
9 distance (table 4).

10 Neither of the pre-operative or post-operative opportunity variables were significantly longitudinally  
11 associated with change in walking at 12-weeks post-operative (table 4 and 6).

12 Walking motivation:

13 In the analysis of the pre-operative motivation variables, the ability to self-regulate exercise, beliefs  
14 about the consequences of the condition, identity, fear of falling, and fear of movement were  
15 significantly associated cross-sectionally with all three walking measures. Fear of falling had a  
16 moderate-strong, negative cross-sectional association with pre-operative walking measures, pre-  
17 operative fear of movement, and consequences had moderate negative associations with pre-  
18 operative walking measures. Pre-operative identity (symptom severity) and ability to self-regulate  
19 exercise had weak negative associations with pre-operative walking measures. Pre-operative  
20 emotional response had a weak negative association with pre-operative 6MWD, and perceived  
21 personal control had a weak positive association with pre-operative step count. Longitudinally, pre-  
22 operative perceived personal control, emotional response and fear of movement were significantly,  
23 albeit weakly, associated with change in maximum walking distance at 12-weeks post-operatively  
24 (table 5).

25 Of the post-operative motivation measures collected 6-weeks post-operatively consequences,  
26 identity, illness concern, emotional response, and fear of falling were significantly associated with  
27 changes in all walking measures at 12-weeks post-operatively ( $<.05$ , table 6). Distress and fear of  
28 movement were significantly associated with change in 6MWD and maximum walking distance. The  
29 strengths of the associations were moderate-weak for fear of movement and fear of falling, and  
30 weak for the remaining variables.

31

1

## 2 Discussion

3 This study aimed to identify the salient, modifiable physical and psychosocial factors associated with  
4 walking in people undergoing surgery for NC. There were statistically significant changes in walking  
5 outcomes after surgery, but these changes were not clinically meaningful for many participants. In  
6 addition, there was no statistically significant change in the number of people walking at least the  
7 recommended 7000 steps/day for older adults<sup>35</sup>. This means that their health may be at risk of the  
8 consequences of inactivity. Consistent with previous studies, the strongest predictor of change in  
9 walking after surgery was pre-operative walking<sup>20</sup>, due to the strength of the association, there was  
10 little scope for the other candidate pre-operative variables to be statistically important. Therefore,  
11 scrutiny of variables associated with pre-operative walking was required. Biopsychosocial factors  
12 from all domains of the COM-B framework were associated with pre-operative walking but only a  
13 few pre-operative measures, from the motivation domain, were longitudinally associated with  
14 change in post-operative walking.

15 The pre-operative factors associated with change in walking at 12-weeks post-operatively were  
16 associated with self-reported maximum walking distance but not objective walking measures. This  
17 discrepancy may be because maximum walking distance assesses an individuals' perception of their  
18 walking capacity, and this aligns more closely with the motivation domain of the COM-B model. It  
19 may also be due to the known discordance between subjective and objective measures of physical  
20 activity<sup>58</sup>. While motivation is theorised as the central mediator in COM-B, our study identified  
21 associations between these factors, but they do not necessarily imply causation.

22 A greater number of candidate predictor variables collected at 6-weeks post-operatively were  
23 longitudinally associated with changes in walking at 12-weeks compared to those collected pre-  
24 operatively. Additionally, factors were associated with all walking measures. Since most healing and  
25 recovery occur within the first 6–12 weeks post-surgery<sup>59,60</sup>, these findings likely reflect the early  
26 effects of surgery, as improvements and differences in outcomes between individuals with good and  
27 poor recovery become apparent. Thus, early post-operative variables are crucial for understanding  
28 post-operative changes, alongside pre-operative factors.

29 Fear of falling pre-operatively and at 6-weeks post-operatively was associated with pre-operative  
30 walking capacity and change in walking respectively. 57% of participants reported a fall within the  
31 last year, this is approximately twice the prevalence of community dwelling older adults<sup>61</sup> and higher  
32 than a non-surgical cohort of older people with NC where prevalence of falling was found to 40%<sup>1</sup>.

1 As falls are a leading cause of injury related deaths in older people, in addition to being associated  
2 with walking, balance and fear of falling is an important area for rehabilitation for patients prior and  
3 after lumbar surgery. Additionally, physical performance tests (grip strength and lower limb physical  
4 performance (individual items and composite score), physical capability) were associated cross-  
5 sectionally with pre-operative and post-operative walking but not with change in walking post-  
6 operatively. Our findings are consistent with previous studies of lumbar surgical patients<sup>62,63</sup> and  
7 thus suggest that physical capability needs to be targeted with rehabilitation pre-operatively to  
8 maximise pre-operative walking.

9 Beliefs regarding the consequences of pain can result in fear of movement and fear-avoidance  
10 behaviours and ultimately less treatment benefit<sup>64</sup>. Pre-operative fear of movement has been found  
11 to be associated with reduced quality of life and increased pain and disability 6-12 months after  
12 lumbar disc surgery<sup>65,66</sup> and 2 years after lumbar surgery for mixed indications<sup>67</sup>. High fear of  
13 movement 6-weeks after lumbar surgery has been found to have a small mediating effect on step  
14 count 12-months after lumbar laminectomy<sup>68</sup>. Our findings which show that pre-operative fear of  
15 movement is associated with pre-operative walking, and that post-operative fear of movement to be  
16 associated with less improvement in post-operative walking are consistent, with the previous  
17 studies. Therefore, our study adds to the growing body of evidence that fear of movement needs to  
18 be addressed in patients undergoing surgery for NC to optimise surgical outcomes.

19 Further beliefs and cognitions were assessed by the Brief Illness Perception Questionnaire<sup>47</sup>, with  
20 most of the constructs associated with walking pre-operatively cross-sectionally, and 6-week post-  
21 operative measures longitudinally, albeit weakly. This questionnaire is based on the self-regulatory  
22 model by Leventhal, which proposes that illness representations influence coping behaviours  
23 through a continuous appraisal loop<sup>69</sup>. There is consistent evidence that illness representations are  
24 correlated with health and treatment outcomes, including low back pain, although have not been  
25 explicitly studied in lumbar surgery<sup>70</sup>. Thus, by identifying pertinent beliefs we may understand how  
26 people undergoing surgery for NC perceive their condition and treatment which in turn may help  
27 explain adopted coping mechanisms and treatment responses and outcome to surgery and should  
28 be explored and addressed with people undergoing surgery for NC.

29

30 The results of this study have important clinical implications for pre-operative and postoperative  
31 care. The findings can be used for risk stratification and to guide rehabilitation. Where previous  
32 interventions have investigated prehabilitation *or* post-operative rehabilitation<sup>71-74</sup> we propose that  
33 patients require rehabilitation before *and* after surgery. Prehabilitation is required to optimise

1 patient's pre-operative walking capacity and performance, and this should be provided in  
2 combination with behaviour change strategies aiming to increase walking and psychological  
3 preparation for surgery. Evidence for prehabilitation and psychological preparation is mixed<sup>71,72,75</sup>. A  
4 recent meta-analysis of prehabilitation programmes for patients undergoing lumbar surgery (15  
5 studies, predominantly for lumbar fusion surgery) suggested that there was low to very low certainty  
6 evidence that there were no additional benefits of psychoeducational interventions on post-  
7 operative physical functioning and pain<sup>71</sup>. Although a randomised controlled trial of 197 patients  
8 with LSS demonstrated that a 9-week multi-modal prehabilitation programme delivered with a  
9 behavioural approach, increased pre-operative walking capacity, and was associated with physical  
10 activity levels one-year post-surgery<sup>72</sup>. This suggest that a complex, systematically developed, multi-  
11 modal intervention, targeting the salient factors identified in this study, specifically for this surgical  
12 population, requires development.

13 Our results indicate that post-operative rehabilitation targeting ongoing fear of falling and  
14 movement is also required. Indeed, only a small proportion of our cohort received any form of post-  
15 operative rehabilitation. Archer et al.<sup>74</sup> demonstrated that in patients with high pre-operative fear of  
16 movement undergoing lumbar laminectomy, 6 weekly sessions of post-operative cognitive-  
17 behavioural-based physical therapy resulted in clinically meaningful improvements in pain, disability,  
18 and physical performance tests 6-month after surgery. This adds credibility to our recommendations  
19 and provides a promising direction for improving outcomes in people with NC.

20 With our results demonstrating that greater walking and physical capacity are associated with  
21 greater improvement after surgery, it could be argued that earlier surgical intervention, before  
22 significant functional decline, might be beneficial. International guidelines for NC recommend a  
23 stepped approach to care, prioritising non-surgical treatments as the first-line approach with  
24 surgical decompression reserved for patients with persistent, severe, or debilitating leg pain<sup>76,77</sup>. The  
25 natural history of NC indicates that rapid progression to severe neurological deficits is uncommon,  
26 and long-term studies show that one-third to half of patients improve without surgery<sup>78-80</sup>.

27 Prognostic indicators for spontaneous improvement remain unclear although, patients with very  
28 tight dural sac areas (<0.5 cm<sup>2</sup>) and severe symptoms are less likely to improve significantly<sup>78-80</sup>.

29 Given these considerations, treatment decisions regarding whether to seek surgical opinion, should  
30 adhere to shared decision-making principles, incorporating patient preferences and clinical  
31 circumstances. Regardless of the chosen treatment pathway, healthcare professionals should  
32 emphasise walking and physical activity as essential components of care.

33

1 Strength and limitations:

2 Strengths of this study include the use of objective and self-rated measures of walking capacity and  
3 performance which has ensured a comprehensive assessment of different walking constructs<sup>81</sup>. Our  
4 focused approach of only including patients with NC, reduced heterogeneity in the study population,  
5 ensuring that the potentially modifiable factors are identified in a manner maximising the chance of  
6 generalisability. To our knowledge, this is the first study within lumbar surgery to use a behaviour  
7 change framework to select a comprehensive battery of candidate predictor variables with a focus  
8 on modifiable factors and therefore the results can be used to inform rehabilitation. Due to  
9 constraints in the study design, it was not feasible to investigate an exhaustive list of candidate  
10 predictor variables, meaning some factors influencing post-operative walking may not have been  
11 identified. However, as pre-operative walking explains a substantial proportion of the change in  
12 post-operative walking, it is unlikely that other factors will substantially explain walking outcomes.  
13 Further limitations include the relatively short follow-up period of three months. However, previous  
14 studies have demonstrated that outcomes between three and 12-months post-surgery are relatively  
15 stable<sup>59</sup>. Not all patients were able to attend for their post-operative objective measures due to  
16 COVID-19 restrictions and some accelerometers were lost in the post resulting in some missing data  
17 however, sufficient data was collected to achieve planned statistical power. Finally, whilst we  
18 identified associations between variables and walking, we are not able to identify the causal  
19 mechanism. Future work should consider whether the same factors are causal mediators.

20 Conclusions

21 Using a comprehensive battery of biopsychosocial measures, mapped to a behaviour change  
22 framework, we aimed to identify modifiable factors associated with walking in people undergoing  
23 surgery for NC. Surgery resulted in statistically significant improvements in walking capacity and  
24 performance, however, for approximately 50% of the sample the changes were not clinically  
25 meaningful. The strongest predictor of change in walking after surgery was pre-operative walking,  
26 the greater the walking before surgery, the greater the improvement after surgery. Several physical  
27 and psychosocial factors were associated with pre-operative walking. The results indicate that  
28 prehabilitation targeting walking, balance and psychosocial factors combined with post-operative  
29 rehabilitation targeting ongoing fear of falling and movement, and unhelpful beliefs may be required  
30 to optimise surgical outcomes in patients undergoing surgery for neurogenic claudication.

31

32 References

1 1 Williamson, E. *et al.* The Prevalence of Back and Leg Pain and the Cross-sectional Association  
2 With Adverse Health Outcomes in Community Dwelling Older Adults in England. *Spine*  
3 (*Philadelphia, Pa. 1976*) **46**, 54-61, doi:10.1097/BRS.0000000000003719 (2021).

4 2 Jensen, R. K., Jensen, T. S., Koes, B. & Hartvigsen, J. Prevalence of lumbar spinal stenosis in  
5 general and clinical populations: a systematic review and meta-analysis. *European spine*  
6 *journal* **29**, 2143-2163 (2020).

7 3 Tomkins-Lane, C. *et al.* ISSLS Prize Winner: Consensus on the Clinical Diagnosis of Lumbar  
8 Spinal Stenosis: Results of an International Delphi Study. *Spine (Phila Pa 1976)* **41**, 1239-  
9 1246, doi:10.1097/BRS.0000000000001476 (2016).

10 4 Winter, C. C. *et al.* Walking ability during daily life in patients with osteoarthritis of the knee  
11 or the hip and lumbar spinal stenosis: a cross sectional study. *BMC Musculoskelet Disord* **11**,  
12 233, doi:10.1186/1471-2474-11-233 (2010).

13 5 Fanuele, J. C., Birkmeyer, N. J. O., Abdu, W. A., Tosteson, T. D. & Weinstein, J. N. The impact  
14 of spinal problems on the health status of patients - Have we underestimated the effect?  
15 *Spine* **25**, 1509-1514, doi:10.1097/00007632-200006150-00009 (2000).

16 6 Deyo, R. A. *et al.* Trends, major medical complications, and charges associated with surgery  
17 for lumbar spinal stenosis in older adults. *JAMA* **303**, 1259-1265, doi:10.1001/jama.2010.338  
18 (2010).

19 7 Lurie, J. & Tomkins-Lane, C. Management of lumbar spinal stenosis. *Bmj* **352** (2016).

20 8 Anderson, D. B. *et al.* Clinical assessment and management of lumbar spinal stenosis: clinical  
21 dilemmas and considerations for surgical referral. *The Lancet Rheumatology* (2024).

22 9 Katz, J. N., Zimmerman, Z. E., Mass, H. & Makhni, M. C. Diagnosis and management of  
23 lumbar spinal stenosis: a review. *Jama* **327**, 1688-1699 (2022).

24 10 Weinstein, J. N. *et al.* Surgical versus nonoperative treatment for lumbar spinal stenosis four-  
25 year results of the Spine Patient Outcomes Research Trial. *Spine (Phila Pa 1976)* **35**, 1329-  
26 1338, doi:10.1097/BRS.0b013e3181e0f04d (2010).

27 11 Aubry, C. *et al.* Accelerometry-based physical activity, disability and quality of life before and  
28 after lumbar decompression surgery from a physiotherapeutic perspective: An observational  
29 cohort study. *North American Spine Society Journal (NASSJ)* **8**, 100087 (2021).

30 12 Smuck, M. *et al.* Objective measurement of function following lumbar spinal stenosis  
31 decompression reveals improved functional capacity with stagnant real-life physical activity.  
32 *Spine J* **18**, 15-21, doi:10.1016/j.spinee.2017.08.262 (2018).

33 13 Mobily, K. E. Walking among older adults. *World Leisure Journal* **56**, 130-140 (2014).

34 14 Ramsey, K. A., Meskers, C. G. & Maier, A. B. Every step counts: synthesising reviews  
35 associating objectively measured physical activity and sedentary behaviour with clinical  
36 outcomes in community-dwelling older adults. *The Lancet Healthy Longevity* **2**, e764-e772  
37 (2021).

38 15 McIlroy, S., Jadhakhan, F., Bell, D. & Rushton, A. Prediction of walking ability following  
39 posterior decompression for lumbar spinal stenosis. *European Spine Journal* **30**, 3307-3318  
40 (2021).

41 16 Aalto, T. J. *et al.* Preoperative predictors for postoperative clinical outcome in lumbar spinal  
42 stenosis: systematic review. *Spine (Phila Pa 1976)* **31**, E648-663,  
43 doi:10.1097/01.brs.0000231727.88477.da (2006).

44 17 Takenaka, H. *et al.* Predictors of walking ability after surgery for lumbar spinal canal stenosis:  
45 a prospective study. *Spine Journal* **19**, 1824-1831 (2019).

46 18 Held, U. *et al.* Prognostic function to estimate the probability of meaningful clinical  
47 improvement after surgery - Results of a prospective multicenter observational cohort study  
48 on patients with lumbar spinal stenosis. *Plos One* **13**, doi:10.1371/journal.pone.0207126  
49 (2018).

1 19 Budithi, S., Dhawan, R., Cattell, A., Balain, B. & Jaffray, D. Only walking matters-assessment  
2 following lumbar stenosis decompression. *Eur Spine J* **26**, 481-487, doi:10.1007/s00586-016-  
3 4881-x (2017).

4 20 McIlroy, S. *et al.* Pre-operative prognostic factors for walking capacity after surgery for  
5 lumbar spinal stenosis: a systematic review. *Age and Ageing* **50**, 1529-1545 (2021).

6 21 Burgstaller, J. M. *et al.* The Influence of Pre- and Postoperative Fear Avoidance Beliefs on  
7 Postoperative Pain and Disability in Patients With Lumbar Spinal Stenosis: Analysis of the  
8 Lumbar Spinal Outcome Study (LSOS) Data. *Spine (Phila Pa 1976)* **42**, E425-E432,  
9 doi:10.1097/BRS.0000000000001845 (2017).

10 22 Archer, K. R. *et al.* The effect of fear of movement beliefs on pain and disability after surgery  
11 for lumbar and cervical degenerative conditions. *Spine* **36**, 1554-1562 (2011).

12 23 Coronado, R. A. *et al.* Bouncing back after lumbar spine surgery: early postoperative  
13 resilience is associated with 12-month physical function, pain interference, social  
14 participation, and disability. *The spine journal* **21**, 55-63, doi:10.1016/j.spinee.2020.07.013  
15 (2021).

16 24 Balck, F., Jeszenszky, C., Günther, K.-P., Kirschner, S. & Linke, M. The impact of illness  
17 perception on functionality, pain, stiffness, and activity of daily living after total hip  
18 replacement surgery. *Journal of Psychosomatic Research* **155**, 110749 (2022).

19 25 Rosenberger, P. H., Jokl, P. & Ickovics, J. Psychosocial factors and surgical outcomes: An  
20 evidence-based literature review. *Journal of the American Academy of Orthopaedic Surgeons*  
21 **14**, 397-405, doi:10.5435/00124635-200607000-00002 (2006).

22 26 Delgado-Ortiz, L. *et al.* Listening to the patients' voice: a conceptual framework of the  
23 walking experience. *Age and Ageing* **52**, afac233 (2023).

24 27 Michie, S., van Stralen, M. M. & West, R. The behaviour change wheel: a new method for  
25 characterising and designing behaviour change interventions. *Implement Sci* **6**, 42,  
26 doi:10.1186/1748-5908-6-42 (2011).

27 28 Hall, J. *et al.* A co-production approach guided by the behaviour change wheel to develop an  
28 intervention for reducing sedentary behaviour after stroke. *Pilot and feasibility studies* **6**, 1-  
29 13 (2020).

30 29 Webb, J., Foster, J. & Poulter, E. Increasing the frequency of physical activity very brief  
31 advice for cancer patients. Development of an intervention using the behaviour change  
32 wheel. *Public Health* **133**, 45-56, doi:10.1016/j.puhe.2015.12.009 (2016).

33 30 von Elm, E. *et al.* The Strengthening the Reporting of Observational Studies in Epidemiology  
34 (STROBE) statement: guidelines for reporting observational studies. *Lancet* **370**, 1453-1457,  
35 doi:10.1016/S0140-6736(07)61602-X (2007).

36 31 de Schepper, E. I. T. *et al.* Diagnosis of Lumbar Spinal Stenosis An Updated Systematic  
37 Review of the Accuracy of Diagnostic Tests. *Spine* **38**, E469-E481,  
38 doi:10.1097/BRS.0b013e31828935ac (2013).

39 32 Steffen, T. M., Hacker, T. A. & Mollinger, L. Age- and gender-related test performance in  
40 community-dwelling elderly people: Six-Minute Walk Test, Berg Balance Scale, Timed Up &  
41 Go Test, and gait speeds. *Phys Ther* **82**, 128-137, doi:10.1093/ptj/82.2.128 (2002).

42 33 Takenaka, H. *et al.* Responsiveness and minimal clinically important difference of the 6-  
43 minute walk distance in patients undergoing lumbar spinal canal stenosis surgery. *Clinical*  
44 *Spine Surgery* **35**, E345-E350 (2022).

45 34 Edwardson, C. L. *et al.* Considerations when using the activPAL monitor in field-based  
46 research with adult populations. *Journal of sport and health science* **6**, 162-178 (2017).

47 35 Tudor-Locke, C. *et al.* How many steps/day are enough? For older adults and special  
48 populations. *International journal of behavioral nutrition and physical activity* **8**, 1-19 (2011).

49 36 Gardner, A. W., Montgomery, P. S., Wang, M. & Shen, B. Minimal clinically important  
50 differences in daily physical activity outcomes following supervised and home-based  
51 exercise in peripheral artery disease. *Vascular Medicine* **27**, 142-149 (2022).



1 37 Tew, G. *et al.* Feasibility and validity of self-reported walking capacity in patients with  
2 intermittent claudication. *Journal of vascular surgery* **57**, 1227-1234 (2013).

3 38 Carlesso, C., Piva, S., Smith, C., Ammendolia, C. & Schneider, M. Responsiveness of Outcome  
4 Measures in Non-Surgical Patients with Lumbar Spinal Stenosis: A Secondary Analysis from a  
5 Randomized Controlled Trial. *Spine (Phila Pa 1976)*, doi:10.1097/BRS.0000000000003920  
6 (2020).

7 39 Fairbank, J. C. T. & Pynsent, P. B. The Oswestry Disability Index. *Spine* **25**, 2940-2953 (2000).

8 40 Janssen, M. F. *et al.* Measurement properties of the EQ-5D-5L compared to the EQ-5D-3L  
9 across eight patient groups: a multi-country study. *Quality of Life Research* **22**, 1717-1727  
10 (2013).

11 41 Royal College of Physicians, British Geriatrics Society & British Pain Society. Vol. 8 (Royal  
12 College of Physicians, London, 2007).

13 42 Tomkins-Lane, C. C. & Battie, M. C. Predictors of objectively measured walking capacity in  
14 people with degenerative lumbar spinal stenosis. *Journal of Back and Musculoskeletal  
15 Rehabilitation* **26**, 345-352, doi:10.3233/bmr-130390 (2013).

16 43 Jia, H. *et al.* Prevalence, risk factors, and burden of disease for falls and balance or walking  
17 problems among older adults in the US. *Preventive medicine* **126**, 105737 (2019).

18 44 Galea Holmes, M. N., Weinman, J. A. & Bearne, L. M. Are Walking Treatment Beliefs and  
19 Illness Perceptions Associated With Walking Intention and 6-Min Walk Distance in People  
20 With Intermittent Claudication? A Cross-Sectional Study. *Journal of Aging & Physical Activity*  
21 **27** (2019).

22 45 Guralnik, J. M. *et al.* A Short Physical Performance Battery Assessing Lower Extremity  
23 Function: Association With Self-Reported Disability and Prediction of Mortality and Nursing  
24 Home Admission. *Journal of Gerontology* **49**, M85-M94, doi:10.1093/geronj/49.2.M85  
25 (1994).

26 46 Mehmet, H., Yang, A. W. H. & Robinson, S. R. Measurement of hand grip strength in the  
27 elderly: A scoping review with recommendations. *J Bodyw Mov Ther* **24**, 235-243,  
28 doi:10.1016/j.jbmt.2019.05.029 (2020).

29 47 Broadbent, E., Petrie, K. J., Main, J. & Weinmann, J. The Brief Illness Perception  
30 Questionnaire. *Journal of Psychosomatic Research* **60**, 631-637,  
31 doi:10.1016/j.jpsychores.2005.10.020 (2006).

32 48 Sniehotta, F., Schwarzer, R., Scholz, U. & Benjamin Schüz, B. Action planning and coping  
33 planning for long-term lifestyle change: theory and assessment. *European Journal of Social  
34 Psychology* **35**, 565-576, doi:<https://doi.org/10.1002/ejsp.258> (2005).

35 49 Spittaels, H. *et al.* Measuring physical activity-related environmental factors: reliability and  
36 predictive validity of the European environmental questionnaire ALPHA. *Int J Behav Nutr  
37 Phys Act* **7**, 48, doi:10.1186/1479-5868-7-48 (2010).

38 50 Sallis, J. F., Grossman, R. M., Pinski, R. B., Patterson, T. L. & Nader, P. R. The development of  
39 scales to measure social support for diet and exercise behaviors. *Prev Med* **16**, 825-836,  
40 doi:10.1016/0091-7435(87)90022-3 (1987).

41 51 Anderson, P. J., Bovard, R. S., Wang, Z., Beebe, T. J. & Murad, M. H. A survey of social  
42 support for exercise and its relationship to health behaviours and health status among  
43 endurance Nordic skiers. *BMJ Open* **6**, e010259, doi:10.1136/bmjopen-2015-010259 (2016).

44 52 Kroenke, K., Spitzer, R. L., Williams, J. B. & Löwe, B. An ultra-brief screening scale for anxiety  
45 and depression: the PHQ-4. *Psychosomatics* **50**, 613-621, doi:10.1176/appi.psy.50.6.613  
46 (2009).

47 53 Delbaere, K. *et al.* The Falls Efficacy Scale International (FES-I). A comprehensive longitudinal  
48 validation study. *Age Ageing* **39**, 210-216, doi:10.1093/ageing/afp225 (2010).

49 54 Swinkels-Meewisse, E., Swinkels, R., Verbeek, A. L. M., Vlaeyen, J. W. S. & Oostendorp, R. A.  
50 B. Psychometric properties of the Tampa Scale for kinesiophobia and the fear-avoidance

1 beliefs questionnaire in acute low back pain. *Manual Therapy* **8**, 29-36,  
2 doi:10.1054/math.2002.0484 (2003).

3 55 Miller, R. P., Kori, S. H. & Todd, D. D. The Tampa Scale: a measure of kinesiophobia . *The*  
4 *Clinical Journal of Pain* **7**, 51 (1991).

5 56 Cohen, J. Statistical power analysis for the behavioural sciences. Hillsdale, New Jersey: L. Lea  
6 **56**, 102 (1988).

7 57 Crutzen, R., Peters, G.-J. Y. & Noijen, J. Using confidence interval-based estimation of  
8 relevance to select social-cognitive determinants for behavior change interventions.  
9 *Frontiers in public health* **5**, 165 (2017).

10 58 Haskell, W. L. Physical activity by self-report: a brief history and future issues. *Journal of*  
11 *Physical Activity and Health* **9**, S5-S10 (2012).

12 59 Ayling, O. G. *et al.* Clinical outcomes research in spine surgery: what are appropriate follow-  
13 up times?: Presented at the 2018 AANS/CNS Joint Section on Disorders of the Spine and  
14 Peripheral Nerves. *Journal of Neurosurgery: Spine* **30**, 397-404 (2018).

15 60 Zeitlberger, A. M. *et al.* Distance to first symptoms measured by the 6-min walking test  
16 differentiates between treatment success and failure in patients with degenerative lumbar  
17 disorders. *European Spine Journal* **31**, 596-603 (2022).

18 61 Salari, N., Darvishi, N., Ahmadipanah, M., Shohaimi, S. & Mohammadi, M. Global prevalence  
19 of falls in the older adults: a comprehensive systematic review and meta-analysis. *Journal of*  
20 *orthopaedic surgery and research* **17**, 1-13 (2022).

21 62 Master, H. *et al.* Physical Performance Tests Provide Distinct Information in Both Predicting  
22 and Assessing Patient-Reported Outcomes Following Lumbar Spine Surgery. *Spine (Phila Pa*  
23 *1976)* **45**, E1556-E1563, doi:10.1097/BRS.0000000000003665 (2020).

24 63 Kwon, J. W. *et al.* Hand grip strength can predict clinical outcomes and risk of falls after  
25 decompression and instrumented posterolateral fusion for lumbar spinal stenosis. *Spine J*  
26 **20**, 1960-1967, doi:10.1016/j.spinee.2020.06.022 (2020).

27 64 Vlaeyen, J. W., Kole-Snijders, A. M., Boeren, R. G. & Van Eek, H. Fear of movement/(re) injury  
28 in chronic low back pain and its relation to behavioral performance. *Pain* **62**, 363-372 (1995).

29 65 Johansson, A. C., Linton, S. J., Rosenblad, A., Bergkvist, L. & Nilsson, O. A prospective study of  
30 cognitive behavioural factors as predictors of pain, disability and quality of life one year after  
31 lumbar disc surgery. *Disability and Rehabilitation* **32**, 521-529,  
32 doi:10.3109/09638280903177243 (2010).

33 66 den Boer, J. J., Oostendorp, R. A., Beems, T., Munneke, M. & Evers, A. W. Continued  
34 disability and pain after lumbar disc surgery: the role of cognitive-behavioral factors. *Pain*  
35 **123**, 45-52 (2006).

36 67 Kemani, M. K., Hägg, O., Jakobsson, M. & Lundberg, M. Fear of movement is related to low  
37 back disability during a two-year period in patients who have undergone elective lumbar  
38 spine surgery. *World Neurosurgery* **137**, e416-e424 (2020).

39 68 Master, H. *et al.* Role of psychosocial factors on the effect of physical activity on physical  
40 function in patients after lumbar spine surgery. *BMC Musculoskeletal Disorders* **22**, 1-11  
41 (2021).

42 69 Leventhal, H., Phillips, L. A. & Burns, E. The Common-Sense Model of Self-Regulation (CSM):  
43 a dynamic framework for understanding illness self-management. *Journal of behavioral*  
44 *medicine* **39**, 935-946 (2016).

45 70 Breland, J. Y., Wong, J. J. & McAndrew, L. M. Are common sense model constructs and self-  
46 efficacy simultaneously correlated with self-management behaviors and health outcomes: A  
47 systematic review. *Health Psychology Open* **7**, 2055102919898846 (2020).

48 71 Janssen, E. R. *et al.* Current prehabilitation programs do not improve the postoperative  
49 outcomes of patients scheduled for lumbar spine surgery: a systematic review with meta-  
50 analysis. *Journal of Orthopaedic & Sports Physical Therapy* **51**, 103-114 (2021).

- 1 72 Fors, M., Enthoven, P., Abbott, A. & Oberg, B. Effects of pre-surgery physiotherapy on  
2 walking ability and lower extremity strength in patients with degenerative lumbar spine  
3 disorder: Secondary outcomes of the PREPARE randomised controlled trial. *BMC*  
4 *Musculoskeletal Disorders* **20 (1) (no pagination)** (2019).
- 5 73 McGregor, A. H. *et al.* Rehabilitation following surgery for lumbar spinal stenosis. A Cochrane  
6 review. *Spine (Phila Pa 1976)* **39**, 1044-1054, doi:10.1097/BRS.0000000000000355 (2014).
- 7 74 Archer, K. R. *et al.* Cognitive-Behavioral-Based Physical Therapy for Patients With Chronic  
8 Pain Undergoing Lumbar Spine Surgery: A Randomized Controlled Trial. *J Pain* **17**, 76-89,  
9 doi:10.1016/j.jpain.2015.09.013 (2016).
- 10 75 Powell, R. *et al.* Psychological preparation and postoperative outcomes for adults  
11 undergoing surgery under general anaesthesia. *Cochrane Database of Systematic Reviews*  
12 (2016).
- 13 76 Anderson, D. B. *et al.* A critical appraisal of clinical practice guidelines for the treatment of  
14 lumbar spinal stenosis. *The Spine Journal* **21**, 455-464 (2021).
- 15 77 Comer, C. *et al.* Consensus on a standardised treatment pathway algorithm for lumbar spinal  
16 stenosis: an international Delphi study. *BMC Musculoskeletal Disorders* **23**, 550 (2022).
- 17 78 Jensen, R. K., Hartvigsen, L. & Kongsted, A. Pain trajectories over 12 months following  
18 conservative care consultation in patients with lumbar spinal stenosis. *BMC Research Notes*  
19 **17**, 174 (2024).
- 20 79 Minamide, A., Yoshida, M. & Maio, K. The natural clinical course of lumbar spinal stenosis: a  
21 longitudinal cohort study over a minimum of 10 years. *Journal of Orthopaedic Science* **18**,  
22 693-698, doi:10.1007/s00776-013-0435-9 (2013).
- 23 80 Otani, K. *et al.* The change of lumbar spinal stenosis symptoms over a six-year period in  
24 community-dwelling people. *Medicina* **57**, 1116 (2021).
- 25 81 Conway, J., Tomkins, C. C. & Haig, A. J. Walking assessment in people with lumbar spinal  
26 stenosis: capacity, performance, and self-report measures. *Spine J* **11**, 816-823,  
27 doi:10.1016/j.spinee.2010.10.019 (2011).

28

29 Acknowledgements:

30 This study was funded by a research training fellowship awarded to Suzanne McIlroy, provided by  
31 the Dunhill Medical Trust. RTF2006\14

32 This study represents independent research supported by the National Institute for Health Research  
33 (NIHR)/Wellcome King's Clinical Research Facility and the NIHR Biomedical Research Centre and  
34 Dementia Unit at South London and Maudsley NHS Foundation Trust and King's College London. The  
35 views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the  
36 Department of Health and Social Care.

37 The authors wish to thank all those involved in the study especially Tim Noblet, Emily Cowles,  
38 Charlotte Walker, Bethany Hedges, Cleo Dobson, Marion Mueller, Elayne Morris, Laura Warne and  
39 Andrew McCarter. We would also like to thank all the participants of this study.

40

1 Author contributions:

2 Suzanne McIlroy: Conceptualization, data curation, formal analysis, funding acquisition,  
3 investigation, methodology, project administration, writing - original draft

4 Lindsay Bearne; Conceptualization, funding acquisition, methodology, supervision, writing – review  
5 & editing

6 John Weinman: Conceptualization, funding acquisition, methodology, supervision, writing – review  
7 & editing

8 Sam Norton: Conceptualization. data curation, funding acquisition, methodology supervision, writing  
9 – review & editing

10

11 Data availability: De-identified data from this study are not available in a public archive. De-  
12 identified data from this study may be made available, on reasonable request, by emailing the  
13 corresponding author.


14

15 Additional information: The authors declare no competing interests.

16


17 Figures:

1 Figure 1 demonstrating flow of participants through the study




2

- 1 Figure 2 boxplots illustrating the walking capacity and performance measures at baseline
- 2 and follow-up.



- 3
- 4
- 5
- 6 Figure 3 a forest plot illustrating the associations between the pre-operative factors with
- 7 pre-operative walking and change in post-operative walking at 12-weeks post-operative.



\*p<.05, \*\*p<.001; adjusted for age, sex, ethnicity, BMI, smoking

1 Table 1: Clinical and demographic characteristics of study participants

Demographic variable		Mean $\pm$ SD or N (%)		
		All participants	Did not receive surgery	Received surgery
		N=134	N=16	N=118
<b>Age (years)</b>		70.2 (8.6)	72.1 (8.2)	69.9 (8.6)
<b>Sex</b>	Female, N (%)	69 (51.5%)	6 (37.5%)	63 (53.4%)
	Male, N (%)	65 (48.5%)	10 (62.5%)	55 (46.6%)
<b>Body mass index</b>		29.3 (5.2)	30.4 (8.3)	29.2 (4.6)
<b>Education</b>	Up to end secondary school N (%)	89 (66.4%)	12 (75.1%)	77 (65.2%)
	High professional or university N (%)	45 (33.6%)	4 (25.0%)	41 (34.7%)
<b>Employment status</b>	Working, N (%)	29 (21.6%)	1 (6.3%)	28 (23.7%)
	Retired, N (%)	95 (70.9%)	14 (87.5%)	81 (68.6%)
<b>Ethnicity</b>	White British, N (%)	97 (72.4%)	11 (68.8%)	86 (72.9%)
	White other, N (%)	8 (6.0%)	1 (6.3%)	7 (5.9%)
	Asian, N (%)	8 (6.0%)	3 (18.8%)	5 (4.2%)
	Black, N (%)	14 (10.4%)	1 (6.3%)	13 (11.0%)
	Mixed, N (%)	4 (3.0%)	0 (0.0%)	4 (3.4%)
	Other, N (%)	3 (2.2%)	0 (0.0%)	3 (2.5%)
<b>Marital or civil status</b>	In a relationship, N (%)	79 (59.0%)	5 (31.3%)	74 (62.7%)
<b>Deprivation indices</b>		6.0 (4.0-9.0)	4.0 (2.5-8.5)	6.0 (4.0-9.0)
<b>Smoking history</b>	Current smoker, N (%)	19 (14.2%)	1 (6.3%)	18 (15.3%)
	Previous smoker, N (%)	64 (47.8%)	6 (37.5%)	58 (49.2%)
<b>Cumulative comorbidity score</b>		8.0 (4.0)	10.9 (4.4)	7.6 (3.7)
<b>PHQ4 score</b>		2.1 (2.8)	2.6 (3.6)	2.0 (2.7)
<b>Duration NC symptoms</b>	months	48.0 (44.6)	60.6 (60.3)	46.2 (42.0)
<b>Prior lumbar surgery</b>		27 (20.1%)	4 (25.0%)	23 (19.5%)
<b>Surgical procedure</b>	Laminectomy			91
	Laminectomy & discectomy			9
	Laminectomy & excision synovial cyst			3
	Laminectomy & foraminotomy			7
	Laminotomy			6
	TLIF or XLIF			2
	No. of levels decompressed			1.4 (0.6)
<b>Leg pain at rest</b>	NRS	4.2 (2.9)	2.4 (2.1)	4.4 (2.9)
<b>Leg pain when walking:</b>	NRS	7.2 (2.4)	6.3 (2.5)	7.3 (2.3)
<b>Back pain at rest</b>	NRS	4.3 (2.8)	3.6 (2.6)	4.4 (2.9)
<b>Back pain when walking</b>	NRS	6.9 (2.8)	6.8 (3.1)	6.9 (2.8)
<b>Disability</b>	ODI	44.1 (15.4)	46.5 (14.8)	43.7 (15.5)
<b>Fallen within last year</b>		76 (57.1%)	12 (75.0%)	64 (54.7%)
<b>No. of falls in last year</b>		1.0 (0.0-4.0)	2.5 (0.5-5.0)	1.0 (0.0-3.0)

PHQ4: patient health questionnaire; NC: neurogenic claudication; NRS: numerical rating scale; ODI: Oswestry Disability Index



1 Table 2: Walking measures at baseline and follow-up

Variable	Baseline mean (±SD)	12 weeks post-operative mean (±SD)	Mean difference (±SD)	95% confidence intervals of mean difference	P value	Percentage achieving MCID
6MWD (m)	238.46 (112.70)	299.93 (104.12)	61.38 (72.57)	46.76, 76.0	<.001	49%
Log of maximum walking distance †	4.71 (1.71)	6.08 (1.85)	1.37 (1.82)	2.75, 1.74	<.001	58%
Daily step count	4878.97 (2502.30)	5461.28 (3042.50)	582.31 (1720.30)	219.93, 944.70	<.001	40.4%
Walking ≥7000 steps/day	19.38%	24.73%	5.35%	n/a	.250	n/a
<p>Table 2 demonstrating the walking capability and performance measures at baseline and 12-week follow up.                      6MWD: six-minute walk distance; MCID: minimal clinically important difference                      † log of data reported as data were skewed</p>						

2

3

1  
2

**Table 3: Regression coefficients for pre-operative capability variables with pre and post-operative walking measures**

Capability candidate pre-operative predictor variable	Time point	6 min walk distance						Step count						Max walking distance					
		b	se	z	p	95%CI ll	95%CI UL	b	se	z	p	95%CI ll	95%CI UL	b	se	z	p	95%CI ll	95%CI UL
Fallen in last year	Pre-op	-0.289	0.079	-3.644	0.000*	-0.445	-0.134	-0.147	0.085	-1.723	0.085	-0.313	0.020	-0.078	0.091	-0.861	0.389	-0.255	0.100
Fallen in last year	12-weeks	-0.087	0.066	-1.330	0.183	-0.216	0.041	-0.071	0.061	-1.178	0.239	-0.190	0.047	-0.137	0.086	-1.596	0.111	-0.306	0.031
No. of falls	Pre-op	-0.314	0.079	-3.984	0.000*	-0.469	-0.160	-0.167	0.083	-2.018	0.044*	-0.328	-0.005	-0.200	0.088	-2.277	0.023*	-0.372	-0.028
No. of falls	12-weeks	-0.234	0.127	-1.845	0.065	-0.482	0.014	0.008	0.108	0.076	0.939	-0.203	0.219	-0.153	0.173	-0.885	0.376	-0.493	0.186
Ability to plan exercise	Pre-op	0.272	0.085	3.210	0.001*	0.106	0.437	0.234	0.090	2.588	0.010*	0.057	0.411	0.213	0.093	2.279	0.023*	0.030	0.396
Ability to plan exercise	12-weeks	-0.105	0.070	-1.502	0.133	-0.242	0.032	0.068	0.063	1.072	0.284	-0.056	0.191	0.115	0.094	1.224	0.221	-0.069	0.299
Coherence of condition	Pre-op	-0.171	0.083	-2.067	0.039*	-0.334	-0.009	-0.070	0.089	-0.783	0.434	-0.245	0.105	-0.231	0.090	-2.552	0.011*	-0.408	-0.054
Coherence of condition	12-weeks	0.028	0.067	0.420	0.674	-0.103	0.159	0.051	0.064	0.790	0.430	-0.075	0.177	0.156	0.091	1.708	0.088	-0.023	0.335
Lower limb performance: SPPB	Pre-op	0.752	0.055	13.565	0.000*	0.643	0.860	0.354	0.083	4.245	0.000*	0.191	0.518	0.487	0.084	5.790	0.000*	0.322	0.652
Lower limb performance: SPPB	12-weeks	0.140	0.103	1.360	0.174	-0.062	0.342	-0.007	0.070	-0.099	0.921	-0.144	0.131	0.154	0.103	1.489	0.137	-0.049	0.357
Max grip strength	Pre-op	0.229	0.091	2.527	0.011*	0.051	0.407	0.119	0.095	1.259	0.208	-0.067	0.305	0.168	0.101	1.658	0.097	-0.031	0.366
Max grip strength	12-weeks	-0.105	0.163	-0.645	0.519	-0.426	0.215	-0.288	0.147	-1.956	0.051	-0.576	0.001	-0.066	0.225	-0.293	0.769	-0.508	0.375

Regression coefficients for pre-operative variables with pre and post-operative walking measures. Estimates are from separate mixed-effects regression models, adjusting for age, sex, ethnicity, BMI, smoking and pre-operative scores.  
Key: SPPB: short physical performance battery; 12-weeks: 12-weeks post-operative walking \* p ≤.05; \*\*p≤001

3

1  
2  
3

**Table 4: Regression coefficients for pre-operative opportunity variables with pre and post-operative walking measures**

Opportunity candidate pre-operative predictor variable	Time point	6 min walk distance						Step count						Max walking distance					
		b	se	z	p	95% CI ll	95% CI ul	b	se	z	p	95% CI ll	95% CI ul	b	se	z	p	95% CI ll	95% CI ul
Social support for exercise	Pre-op	0.284	0.085	3.349	0.001*	0.118	0.451	0.155	0.088	1.755	0.079	-0.018	0.328	0.127	0.094	1.347	0.178	-0.058	0.312
Social support for exercise	12-weeks	0.001	0.064	0.020	0.984	-0.124	0.127	0.040	0.067	0.602	0.548	-0.091	0.172	0.001	0.089	-0.009	0.993	-0.174	0.173
Physical environment suitability for walking	Pre-op	0.284	0.086	3.295	0.001*	0.115	0.452	0.099	0.093	1.065	0.287	-0.083	0.280	0.180	0.095	1.891	0.059	-0.007	0.366
Physical environment suitability for walking	12-weeks	-0.008	0.073	-0.109	0.913	-0.151	0.135	-0.042	0.068	-0.613	0.540	-0.174	0.091	0.104	0.102	1.020	0.308	-0.096	0.303

Regression coefficients for pre-operative variables with pre and post-operative walking measures. Estimates are from separate mixed-effects regression models, adjusting for age, sex, ethnicity, BMI, smoking and pre-operative scores.  
Key: 12-weeks: 12-weeks post-operative walking \* p ≤.05; \*\*p≤001

4

1

**Table 5: Regression coefficients for pre-operative motivation variables with pre and post-operative walking measures**

Motivation candidate pre-operative predictor variable	Time point	6 min walk distance						Step count						Max walking distance					
		b	se	z	p	95% CI ll	95% Cul	b	se	z	p	95% CI ll	95% Cul	b	se	z	p	95% CI ll	95% Cul
Distress	Pre-op	-0.162	0.083	-1.944	0.052	-0.325	0.001	0.103	0.085	1.222	0.222	-0.062	0.269	-0.115	0.090	-1.268	0.205	-0.292	0.063
Distress	12-weeks	-0.004	0.071	-0.058	0.953	-0.144	0.135	-0.009	0.067	-0.133	0.894	-0.140	0.122	-0.041	0.095	-0.437	0.662	-0.227	0.144
Ability to self-regulate exercise	Pre-op	0.244	0.084	2.921	0.004*	0.080	0.408	0.242	0.090	2.689	0.007*	0.066	0.419	0.248	0.090	2.741	0.006*	0.071	0.425
Ability to self-regulate exercise	12-weeks	-0.086	0.065	-1.308	0.191	-0.214	0.043	-0.001	0.062	-0.014	0.989	-0.122	0.120	0.027	0.093	0.290	0.772	-0.156	0.210
Consequences of condition	Pre-op	-0.358	0.079	-4.498	0.000*	-0.513	-0.202	-0.304	0.084	-3.603	0.000*	-0.469	-0.138	-0.411	0.085	-4.857	0.000*	-0.577	-0.245
Consequences of condition	12-weeks	0.104	0.070	1.487	0.137	-0.033	0.242	0.044	0.065	0.683	0.494	-0.083	0.172	0.084	0.095	0.891	0.373	-0.101	0.270
Timeline beliefs	Pre-op	-0.161	0.087	-1.845	0.065	-0.331	0.010	-0.087	0.090	-0.964	0.335	-0.264	0.090	0.002	0.097	0.019	0.985	-0.188	0.192
Timeline beliefs	12-weeks	0.012	0.065	0.191	0.848	-0.115	0.140	0.041	0.060	0.684	0.494	-0.076	0.158	-0.073	0.091	-0.796	0.426	-0.252	0.106
Perceived personal control	Pre-op	0.143	0.085	1.692	0.091	-0.023	0.309	0.258	0.085	3.036	0.002*	0.091	0.425	0.088	0.092	0.962	0.336	-0.092	0.269
Perceived personal control	12-weeks	0.084	0.069	1.218	0.223	-0.051	0.220	-0.018	0.067	-0.268	0.788	-0.149	0.113	0.189	0.093	2.028	0.043*	0.006	0.372
Perceived treatment control	Pre-op	0.018	0.084	0.216	0.829	-0.147	0.183	-0.013	0.087	-0.146	0.884	-0.184	0.158	-0.082	0.091	-0.899	0.369	-0.259	0.096
Perceived treatment control	12-weeks	0.065	0.069	0.937	0.349	-0.071	0.200	-0.005	0.064	-0.075	0.941	-0.130	0.120	0.159	0.097	1.638	0.101	-0.031	0.349
Identity (symptom severity)	Pre-op	-0.272	0.080	-3.424	0.001*	-0.428	-0.116	-0.214	0.084	-2.543	0.011*	-0.378	-0.049	-0.297	0.086	-3.444	0.001*	-0.465	-0.128
Identity (symptom severity)	12-weeks	0.095	0.061	1.557	0.120	-0.025	0.216	0.050	0.057	0.869	0.385	-0.062	0.161	-0.048	0.089	-0.541	0.589	-0.223	0.126
Illness concern	Pre-op	-0.075	0.083	-0.903	0.366	-0.237	0.088	-0.063	0.084	-0.752	0.452	-0.228	0.102	-0.047	0.089	-0.524	0.600	-0.222	0.129
Illness concern	12-weeks	0.097	0.056	1.712	0.087	-0.014	0.207	0.007	0.052	0.126	0.900	-0.096	0.109	-0.060	0.081	-0.740	0.460	-0.219	0.099
Emotional response	Pre-op	-0.170	0.082	-2.074	0.038*	-0.330	-0.009	-0.022	0.086	-0.253	0.800	-0.190	0.146	-0.125	0.089	-1.408	0.159	-0.300	0.049
Emotional response	12-weeks	-0.077	0.060	-1.269	0.204	-0.195	0.042	-0.005	0.056	-0.094	0.925	-0.115	0.104	-0.171	0.085	-2.014	0.044*	-0.338	-0.005
Fear of falling	Pre-op	-0.549	0.071	-7.747	0.000*	-0.687	-0.410	-0.315	0.082	-3.829	0.000*	-0.477	-0.154	-0.363	0.086	-4.243	0.000*	-0.530	-0.195
Fear of falling	12-weeks	-0.110	0.081	-1.351	0.177	-0.269	0.049	0.033	0.064	0.511	0.609	-0.093	0.159	-0.134	0.101	-1.327	0.184	-0.333	0.064
Fear of movement	Pre-op	-0.483	0.076	-6.345	0.000*	-0.633	-0.334	-0.256	0.088	-2.900	0.004*	-0.428	-0.083	-0.215	0.093	-2.312	0.021*	-0.396	-0.033
Fear of movement	12-weeks	0.055	0.076	0.718	0.473	-0.094	0.203	-0.035	0.061	-0.571	0.568	-0.153	0.084	-0.208	0.090	-2.322	0.020*	-0.384	-0.032

Regression coefficients for pre-operative variables with pre and post-operative walking measures. Estimates are from separate mixed-effects regression models, adjusting for age, sex, ethnicity, BMI, smoking and pre-operative scores. Key: 12-weeks: 12-weeks post-operative walking \* p ≤ 0.05; \*\* p ≤ 0.001

2

1 **Table 6: Regression coefficients for variables collected 6-week post-surgery with change in walking at 12-weeks post-surgery**

2

Post-operative candidate predictor variable	6 min walk distance						Step count						Max walking distance					
	b	se	z	p	95%CI ll	95%CI ul	b	se	z	p	95%CI ll	95%CI ul	b	se	z	p	95%CI ll	95%CI ul
<b>Walking capability</b>																		
Fallen in last 6 weeks	-0.140	0.065	-2.143	0.032*	-0.268	-0.012	0.023	0.061	0.378	0.705	-0.096	0.142	-0.075	0.100	-0.742	0.458	-0.271	0.122
Ability to plan exercise	-0.035	0.070	-0.495	0.621	-0.172	0.102	0.012	0.067	0.179	0.858	-0.120	0.144	-0.029	0.087	-0.333	0.739	-0.199	0.141
Coherence of condition	0.088	0.072	1.211	0.226	-0.054	0.230	0.031	0.069	0.457	0.647	-0.103	0.166	0.057	0.097	0.584	0.559	-0.133	0.246
<b>Walking opportunity</b>																		
Social support to exercise	0.050	0.080	0.619	0.536	-0.108	0.207	0.101	0.089	1.131	0.258	-0.074	0.276	0.164	0.121	1.356	0.175	-0.073	0.401
<b>Walking motivation</b>																		
Distress	-0.252	0.063	-4.020	0.000*	-0.375	-0.129	-0.084	0.065	-1.285	0.199	-0.212	0.044	-0.270	0.092	-2.944	0.003*	-0.449	-0.090
Ability to self-regulate exercise	-0.014	0.075	-0.186	0.852	-0.160	0.132	0.018	0.068	0.261	0.794	-0.116	0.152	0.062	0.096	0.641	0.522	-0.127	0.250
Consequences of condition	-0.133	0.062	-2.162	0.031*	-0.254	-0.012	-0.155	0.056	-2.765	0.006*	-0.265	-0.045	-0.233	0.086	-2.712	0.007*	-0.402	-0.065
Timeline beliefs	-0.192	0.069	-2.789	0.005*	-0.326	-0.057	-0.112	0.066	-1.694	0.090	-0.242	0.018	-0.153	0.099	-1.548	0.122	-0.346	0.041
Perceived personal control	0.045	0.068	0.672	0.501	-0.087	0.178	-0.013	0.062	-0.213	0.831	-0.134	0.108	0.038	0.096	0.394	0.693	-0.151	0.227
Perceived treatment control	0.146	0.067	2.179	0.029*	0.015	0.278	0.041	0.066	0.617	0.537	-0.088	0.169	0.178	0.092	1.933	0.053	-0.002	0.358
Identity (Symptom severity)	-0.243	0.061	-4.000	0.000*	-0.362	-0.124	-0.197	0.058	-3.399	0.001*	-0.310	-0.083	-0.232	0.087	-2.667	0.008*	-0.402	-0.061
Illness concern	-0.227	0.064	-3.576	0.000*	-0.352	-0.103	-0.142	0.061	-2.313	0.021*	-0.262	-0.022	-0.270	0.086	-3.130	0.002*	-0.439	-0.101
Emotional response	-0.241	0.074	-3.246	0.001*	-0.386	-0.095	-0.243	0.067	-3.618	0.000*	-0.375	-0.111	-0.220	0.098	-2.234	0.025*	-0.413	-0.027
Fear of falling	-0.178	0.079	-2.252	0.024*	-0.333	-0.023	-0.191	0.073	-2.639	0.008*	-0.333	-0.049	-0.354	0.112	-3.156	0.002*	-0.573	-0.134
Fear of movement	-0.230	0.086	-2.679	0.007*	-0.398	-0.062	-0.123	0.077	-1.594	0.111	-0.274	0.028	-0.317	0.107	-2.961	0.003*	-0.527	-0.107

Regression coefficients for variables collected 6-weeks post-surgery with change in walking at 12-weeks post-surgery. Estimates are from separate mixed-effects regression models, adjusting for age, gender, ethnicity, BMI, smoking and baseline score. \* p ≤.05; \*\*p≤001

3

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23

**Supplementary material**

Title: Identifying modifiable factors that influence walking in patients undergoing surgery for neurogenic claudication. A prospective longitudinal study

Authors:

Suzanne McIlroy<sup>1, 2\*</sup>, Lindsay Bearne<sup>3</sup>, John Weinman<sup>4</sup>, Sam Norton<sup>2</sup>

Affiliations:

1 Physiotherapy Department, King’s College Hospital, London, UK.

2 Health Psychology Section, King’s College London, London, UK

3 Population Health Research Institute, City St George’s, University of London, London, UK

4 School of cancer and pharmaceutical science, King’s College London, UK

**Contents**

**Supplementary Table S1 Change in disability, pain, quality of life, fear of falling and fear of movement after surgery for neurogenic claudication..... 31**

**Supplementary Figure S1 Box plots demonstrating disability (ODI), quality of life (EQ5D) and leg and back pain at rest and on walking, fear of movement (Tampa) and fear of falling (SFESI) at baseline and follow up. .... 32**

**Supplementary Table S2 Residualised change scores for the walking measures..... 34**

**Supplemental Table S3 Unadjusted correlations between pre-operative predictor variables and walking measures pre and post-operatively..... 35**

**Supplemental Table S4 Correlations between post-operative candidate predictor variables and 12-week walking measures ..... 36**

1


2 **Supplementary Table S1 Change in disability, pain, quality of life, fear of falling and fear of movement after surgery for neurogenic claudication**

Variable	n	Baseline mean ( $\pm$ SD) n=133	12 weeks post-operative mean ( $\pm$ SD)	Mean difference ( $\pm$ SD)	95% confidence intervals of mean difference	P value
Disability (ODI)	107	42.99 (15.08)	27.08 (17.86)	-15.90 (15.45)	-18.87, -12.94	<.001
Leg pain at rest (NRS)	104	4.20 (2.86)	1.55 (2.21)	-2.65 (2.82)	-3.20, -2.10	<.001
Leg pain when walking (NRS)	104	7.29 (2.38)	3.03 (2.99)	-4.26 (3.19)	-4.88, -3.64	<.001
Back pain at rest (NRS)	103	4.22 (2.84)	1.81 (2.45)	-2.42 (2.72)	-2.95, -1.89	<.001
Back pain when walking (NRS)	103	6.84 (2.86)	3.53 (2.86)	-3.33 (3.01)	-3.92, -2.74	<.001
Quality of life (EQ5D-5L)	106	.434 (.230)	.660 (.203)	.226 (.245)	.178, .273	<.001
Fear of falling (SFESI)	106	14.53 (5.14)	11.31 (4.84)	-3.22 (4.37)	-4.06, -2.38	<.001
Fear of movement (Tampa)	100	42.65 (7.27)	38.45 (7.83)	-4.20 (6.92)	-5.57, -2.83	<.001
ODI: Oswestry disability index; NRS: numerical rating scale; EQ5D-5L: EuroQol five dimension, 5-level questionnaire; SFESI: Short Falls Efficacy Scale International; Tampa: Tampa Scale of Kinesiophobia						

3

4

1 **Supplementary Figure S1** Box plots demonstrating disability (ODI), quality of life (EQ5D) and leg and back pain at rest and on walking, fear of movement  
2 **(Tampa)** and fear of falling (SFESI) at baseline and follow up.





3

4



5



1



2



3

4

1 **Supplementary Table S2 Residualised change scores for the walking measures**

Walking measure	6MWD			Step Count			Log of max walk distance		
	B	p	95ll/95ul	b	p	95ll/95ul	b	p	95ll/95ul
_base	0.77	0.000	0.63 to 0.91	0.91	0.000	0.77 to 1.04	0.41	0.000	0.24 to 0.57
age10	-2.08	0.046	-4.12 to -0.04	-2.89	0.004	-4.85 to -0.93	-0.30	0.832	-3.07 to 2.47
age10 # age10	0.13	0.077	-0.01 to 0.28	0.19	0.008	0.05 to 0.34	-0.00	0.980	-0.20 to 0.20
female	-0.09	0.536	-0.36 to 0.19	0.07	0.549	-0.16 to 0.30	-0.49	0.004	-0.83 to -0.16
White British	0.31	0.044	0.01 to 0.61	-0.13	0.360	-0.42 to 0.15	0.30	0.135	-0.09 to 0.68
Obese	-0.03	0.805	-0.31 to 0.24	0.18	0.166	-0.08 to 0.44	-0.37	0.053	-0.75 to 0.00
Current smoker	0.15	0.408	-0.20 to 0.50	-0.29	0.086	-0.63 to 0.04	0.14	0.563	-0.34 to 0.63
Constant	7.64	0.031	0.71 to 14.57	10.61	0.002	3.94 to 17.28	2.34	0.625	-7.15 to 11.83
Observations	96			88			99		
R-squared	0.674			0.753			0.380		
Table demonstrating the residualised change scores for the walking measures Key: 6MWD: 6 minute walk distance; 95ll/95ul: 95% confidence intervals									

2

3

1  
2

**Supplemental Table S3 Unadjusted correlations between pre-operative predictor variables and walking measures pre and post-operatively**

Correlations between pre-operative variables and walking measures at baseline & 12-weeks post-surgery						
Pre-operative variables	Baseline walking measure			Walking measure at 12-weeks post-surgery		
	6MWD (m)	Max walking distance (m)	Step Count	6MWD (m)	Max walking distance (m)	Step Count
6MWD (m)	1.000			0.779***	0.305**	0.427***
Max walking distance (m)	0.487***	1.000		0.374***	0.402***	0.263*
Step Count	0.585***	0.461***	1.000	0.548***	0.414***	0.825***
Sit to Stand Score (SPPB)	0.659***	0.204*	0.275**	0.548***	0.194	0.319**
Standing Balance Score (SPPB)	0.541***	0.222*	0.335***	0.464***	0.242*	0.225*
Gait Speed (SPPB)	0.704***	0.348***	0.375***	0.563***	0.225*	0.262*
SPPB Total score	0.799***	0.325***	0.418***	0.673***	0.277**	0.345***
Maximum grip strength (kg)	0.249**	0.068	0.125	0.369***	0.128	0.168
Walking aid	-0.668***	-0.326***	-0.465***	-0.677***	-0.310**	-0.432***
Fallen in last year	-0.315***	-0.044	-0.141	-0.299**	-0.122	-0.207*
No. of falls	-0.259**	-0.080	-0.142	-0.223*	-0.035	-0.090
Ability to plan exercise	0.282**	0.158	0.210*	0.099	0.156	0.176
Ability to self-regulate exercise	0.257**	0.275**	0.243**	0.072	0.246*	0.190
Physical environment and suitability for walking (ALPHA)	0.317***	0.151	0.116	0.202*	0.103	0.162
Social support & exercise survey	0.299***	0.209*	0.230*	0.255*	0.104	0.317**
Live alone	0.010	0.054	-0.124	-0.259*	-0.153	-0.165
In a relationship	0.039	-0.001	0.079	0.299**	0.232*	0.226*
PHQ4 score	-0.219*	-0.115	0.103	-0.122	-0.004	0.180
EQ5D VAS	0.390***	0.254**	0.328***	0.293**	0.191	0.219*
EQ5D index value	0.435***	0.355***	0.326***	0.306**	0.135	0.209*
Consequences of condition	-0.403***	-0.414***	-0.344***	-0.216*	-0.111	-0.125
Timeline beliefs	-0.126	-0.058	-0.123	-0.061	-0.150	-0.091
Perceived personal control	0.220*	0.175*	0.291**	0.322**	0.232*	0.168
Perceived treatment control	0.062	0.047	0.038	0.158	0.096	0.109
Identity (symptom severity)	-0.332***	-0.358***	-0.247**	-0.079	-0.135	-0.029
Illness concern	-0.079	-0.171	-0.081	0.107	-0.260**	0.025
Coherence of condition	-0.141	-0.098	-0.006	-0.050	0.062	-0.009
Emotional response	-0.214*	-0.210*	-0.044	-0.197	-0.237*	0.025
Fear of falling:	-0.596***	-0.301***	-0.376***	-0.509***	-0.251*	-0.281**
Fear of movement	-0.442***	-0.176	-0.282**	-0.224*	-0.200*	-0.151

\*  $p < 0.05$ , \*\*  $p < 0.01$ , \*\*\*  $p < 0.001$  ALPHA: physical activity-related environmental factors (ALPHA) scale; PHQ4: Patient health questionnaire 4; EQ5D-5L: EuroQol five dimension, 5-level questionnaire; VAS: visual analogue scale; NRS: numerical rating scale; SPPB: Short physical performance battery; 6MWD: six minute walk distance

1

2

**Supplemental Table S4 Correlations between post-operative candidate predictor variables and 12-week walking measures**

<b>Correlations between variables at 6-weeks and 12-weeks post-surgery and walking measures at 12-weeks post-surgery</b>						
Variables	Correlations between variables at 6-weeks and 12-week walking measures			Correlations between variables at 12-weeks post-operative and 12-week walking measures		
	6MWD	Max walking distance	Step Count	6MWD	Max walking distance	Step Count
6MWD (m)	n/a	n/a	n/a	1.000		
Maximum walking distance (m)	0.498***	0.514***	0.273*	0.430***	1.000	
Step Count	n/a	n/a	n/a	0.513***	0.481***	1.000
Sit to Stand Score (SPPB)	n/a	n/a	n/a	0.686***	0.411***	0.471***
Standing Balance Score (SPPB)	n/a	n/a	n/a	0.415***	0.240*	0.259*
Gait Speed (SPPB)	n/a	n/a	n/a	0.655***	0.383***	0.403***
SPPB Total score	n/a	n/a	n/a	0.781***	0.451***	0.499***
Maximum grip (kg)	n/a	n/a	n/a	0.380***	0.205*	0.191
Walking aid	-0.718***	-0.379***	-0.462***			
Fallen in previous 6 weeks	-0.252*	-0.154	-0.103	-0.230*	-0.165	-0.135
No. of falls last 6 weeks	-0.179	-0.081	-0.097	-0.208*	-0.113	-0.117
Ability to plan exercise	0.065	0.079	0.242*	-0.040	0.087	0.176
Ability to self-regulate exercise	0.105	0.161	0.314**	0.060	0.171	0.252*
Social support & exercise survey (ALPHA)	0.158	0.165	0.160	0.220*	0.318**	0.246*
Consequences of condition	-0.328**	-0.382***	-0.280**	-0.497***	-0.395***	-0.284**
Timeline beliefs	-0.401***	-0.321**	-0.262*	-0.223*	-0.224*	-0.105
Perceived personal control	0.328**	0.178	0.166	0.250*	0.233*	0.213*
Perceived treatment control	0.276**	0.230*	0.159	0.165	0.156	0.137
Identity (symptom severity)	-0.383***	-0.366***	-0.304**	-0.323**	-0.407***	-0.308**
Illness concern	-0.336***	-0.357***	-0.147	-0.307**	-0.327***	-0.177
Coherence of condition	0.174	0.110	0.031	0.088	0.042	-0.043
Emotional response	-0.393***	-0.299**	-0.178	-0.391***	-0.233*	-0.142
Total PHQ4	-0.126	-0.003	0.181	-0.109	-0.003	0.187
EQ5D index value	0.555***	0.329***	0.403***	0.569***	0.402***	0.430***
EQ5D VAS	0.433***	0.276**	0.169	0.466***	0.304**	0.261*
Fear of falling	-0.531***	-0.358***	-0.328**	-0.651***	-0.402***	-0.394***
Fear of movement	-0.386***	-0.282**	-0.226*	-0.339***	-0.327***	-0.276**

Hospital recorded complication	-0.183	-0.084	-0.074	n/a	n/a	n/a
Length of stay post-op	-0.401***	-0.225*	-0.303**	n/a	n/a	n/a
Patient reported post-op complication	-0.027	-0.055	-0.110	-0.074	-0.149	-0.049
Received inpatient physiotherapy	0.031	-0.121	0.031	n/a	n/a	n/a
Received outpatient physiotherapy	0.033	0.054	0.241*	-0.106	-0.118	0.071
<p>* <math>p &lt; 0.05</math>, ** <math>p &lt; 0.01</math>, *** <math>p &lt; 0.001</math>  PHQ4: Patient health questionnaire 4; EQ5D-5L: EuroQol five dimension, 5-level questionnaire; VAS: visual analogue scale; NRS: numerical rating scale; SPPB: Short physical performance battery; 6MWD: six minute walk distance; 6MWD: 6 minute walk distance</p>						

1

2