


















# Status and implications of the knowledge, attitudes and practices towards AWaRe antibiotic use, resistance and stewardship among low- and middle-income countries

Zikria Saleem <sup>1</sup>, Catrin E. Moore <sup>2</sup>, Aubrey C. Kalungia <sup>3</sup>, Natalie Schellack <sup>4</sup>, Olayinka Ogunleye <sup>5,6</sup>, Audrey Chigome<sup>7</sup>, Kona Chowdhury <sup>8</sup>, Freddy Eric Kitutu <sup>9,10,11</sup>, Amos Massele <sup>12</sup>, Nishana Ramdas<sup>7</sup>, E. Sam Orubu<sup>13,14</sup>, Aislinn Cook <sup>2,15</sup>, Felix Khuluza<sup>16</sup>, Trust Zaranyika<sup>17</sup>, Elisa Funicello <sup>2</sup>, Giulia Lorenzetti<sup>2</sup>, Miriam Nantamu<sup>2</sup>, Ayuska Parajuli <sup>18,19</sup>, Amanj Kurdi <sup>7,20,21,22</sup>, Hellen Nabayiga<sup>23</sup>, Ammar Abdulrahman Jairoun <sup>24,25,26</sup>, Mainul Haque <sup>27,28</sup>, Stephen M. Campbell <sup>7,29</sup>, Dena Van Der Bergh <sup>30</sup>, Brian Godman <sup>2,7,21,25\*</sup> and Mike Sharland<sup>2</sup>

<sup>1</sup>Department of Pharmacy Practice, Faculty of Pharmacy, Bahauddin Zakariya University, Multan 60800, Pakistan; <sup>2</sup>Centre for Neonatal and Paediatric Infection, Institute for Infection and Immunity, City St George's, University of London, London SW17 0RE, UK; <sup>3</sup>Department of Pharmacy, School of Health Sciences, University of Zambia, Lusaka P.O. Box 50110, Zambia; <sup>4</sup>Department of Pharmacology, Faculty of Health Sciences, University of Pretoria, Pretoria 0084, South Africa; <sup>5</sup>Department of Pharmacology, Therapeutics and Toxicology, Lagos State University College of Medicine, Ikeja, Lagos 100271, Nigeria; <sup>6</sup>Department of Medicine, Lagos State University Teaching Hospital, Ikeja 100271, Nigeria; <sup>7</sup>Department of Public Health Pharmacy and Management, School of Pharmacy, Sefako Makgatho Health Sciences University, Ga-Rankuwa 0208, South Africa; <sup>8</sup>Department of Pediatrics, Gonoshasthaya Somaj Vittik Medical College, Dhaka 1344, Bangladesh; <sup>9</sup>Sustainable Pharmaceutical Systems (SPS) Unit, School of Health Sciences, Makerere University, PO Box 7072, Kampala, Uganda; <sup>10</sup>Department of Women's and Children's Health, International Maternal and Child Health (IMCH), Uppsala University, SE-751 85 Uppsala, Sweden; <sup>11</sup>Department of Pharmacy, Makerere University School of Health Sciences, Kampala, Uganda; <sup>12</sup>Department of Clinical Pharmacology and Therapeutics, Hurler Kairuki Memorial University, 70 Chwaku Road Mikocheni, Dar Es Salaam P.O. Box 65300, Tanzania; <sup>13</sup>Department of Biomedical Engineering, Boston University College of Engineering, Boston, MA, USA; <sup>14</sup>Institute for Health System Innovation & Policy, Boston University, Boston, MA, USA; <sup>15</sup>Health Economics Research Centre, Nuffield Department of Population Health, University of Oxford, Oxford OX1 2JD, UK; <sup>16</sup>Pharmacy Department, Formerly College of Medicine, Kamuzu University of Health Sciences (KUHeS), Blantyre P.O. Box 278, Malawi; <sup>17</sup>Faculty of Medicine and Health Sciences, University of Zimbabwe, Harare P.O. Box MP167, Zimbabwe; <sup>18</sup>HERD International, Lalitpur, Nepal; <sup>19</sup>Public Health Research Society, Kathmandu, Nepal; <sup>20</sup>Department of Clinical Pharmacy, College of Pharmacy, Hawler Medical University, Kurdistan Regional Governorate, Erbil, Iraq; <sup>21</sup>Strathclyde Institute of Pharmacy and Biomedical Science (SIPBS), University of Strathclyde, Glasgow G4 0RE, UK; <sup>22</sup>College of Pharmacy, Al-Kitab University, Kirkuk 36015, Iraq; <sup>23</sup>Management Science Department, Strathclyde Business School, University of Strathclyde, 199 Cathedral Street, Glasgow G4 0QU, UK; <sup>24</sup>Health and Safety Department, Dubai Municipality, Dubai, United Arab Emirates; <sup>25</sup>Center of Medical and Bio-Allied Health Sciences Research, Ajman University, Ajman 346, United Arab Emirates; <sup>26</sup>Department of Clinical Sciences, College of Pharmacy and Health Sciences, Ajman University, Ajman 346, United Arab Emirates; <sup>27</sup>Unit of Pharmacology, Faculty of Medicine and Defence Health, Universiti Pertahanan Nasional Malaysia (National Defence University of Malaysia), Kem Sungai, Besi, 57000 Kuala Lumpur, Malaysia; <sup>28</sup>Karnavati Scientific Research Center, Karnavati School of Dentistry, Karnavati University, Gandhinagar 382 422, Gujarat, India; <sup>29</sup>School of Health Sciences, University of Manchester, Manchester, UK; <sup>30</sup>Division of Infectious Diseases and HIV Medicine, Department of Medicine, Groote Schuur Hospital, University of Cape Town, Cape Town, South Africa

\*Corresponding author. E-mail: Brian.Godman@strath.ac.uk

**Background:** There are concerns globally with rising rates of antimicrobial resistance (AMR), particularly in low- and middle-income countries (LMICs). AMR is driven by high rates of inappropriate prescribing and dispensing of antibiotics, particularly Watch antibiotics. To develop future interventions, it is important to document current knowledge, attitudes and practices (KAP) among key stakeholder groups in LMICs.

**Methods:** We undertook a narrative review of published papers among four WHO Regions including African and Asian countries. Relevant papers were sourced from 2018 to 2024 and synthesized by key stakeholder group,

country, WHO Region, income level and year. The findings were summarized to identify pertinent future activities for all key stakeholder groups.

**Results:** We sourced 459 papers, with a large number coming from Africa (42.7%). An appreciable number dealt with patients' KAP (33.1%), reflecting their influence on the prescribing and dispensing of antibiotics. There was marked consistency of findings among key stakeholder groups across the four WHO Regions, all showing concerns with high rates of prescribing of antibiotics for viral infections despite professed knowledge of antibiotics and AMR. There were similar issues among dispensers. Patients' beliefs regarding the effectiveness of antibiotics for self-limiting infectious diseases were a major challenge, although educational programmes did improve knowledge. The development of the AWaRe (Access, Watch and Reserve) system, including practical prescribing guidance, provides a future opportunity for the standardization of educational inputs.

**Conclusions:** Similar KAP regarding the prescribing and dispensing of antibiotics across LMICs and stakeholder groups presents clear opportunities for standardization of educational input and practical training programmes based on the AWaRe system.

## Introduction

Antimicrobial resistance (AMR) increases morbidity, mortality and healthcare costs, and is now seen as a serious public health concern.<sup>1-4</sup> If not addressed, AMR could cause or exacerbate the next pandemic.<sup>5,6</sup> Whereas AMR is ubiquitous, the burden is disproportionately higher in low- and middle-income countries (LMICs), which currently account for 80% of an estimated 10 million deaths annually attributable to AMR.<sup>7-10</sup> AMR levels will continue increasing in LMICs unless addressed, with current high rates of AMR exacerbated by the growing use of antibiotics from the WHO's Watch list, with resultant implications for increasing rates of MDR bacteria.<sup>11,12</sup> The classification of antibiotics into Access, Watch and Reserve (AWaRe) groups in the WHO Essential Medicines List was part of global initiatives to reduce inappropriate use of Watch and Reserve antibiotics with their greater potential for selection of antibiotic resistance (ABR).<sup>11,13,14</sup> Alongside this, across LMICs including African countries, there is continued inappropriate prescribing and dispensing of antibiotics generally, including among children, exacerbated by limited knowledge and awareness of AMR.<sup>11,15-21</sup>

A number of global and regional initiatives have been introduced in recent years to try to minimize rising rates of ABR and reduce its consequences.<sup>22-25</sup> These include the WHO's Global Action Plan (GAP) to reduce AMR, translated into National Action Plans (NAPs) across member states.<sup>22,26</sup> Other initiatives include the classification of antibiotics into AWaRe groups as well as the WHO AWaRe antibiotic book guiding optimal use of antibiotics in 35 infections in adults and children for both the primary care and hospital settings, published in 2022.<sup>14,27,28</sup> However, there are currently major concerns with the implementation of NAPs, especially among LMICs, including among African and Asian countries, given the demands on available personnel and resources.<sup>29-33</sup> Similar concerns exist regarding national and local implementation of the guidance included within the WHO AWaRe book.<sup>34-36</sup>

The greatest use of antibiotics in humans in LMICs is in primary care, which can account for up to 90% of total antibiotic use in humans and consists of mainly older multiple sourced oral Access and Watch antibiotics.<sup>37</sup> Consequently, strategies aimed at reducing inappropriate antibiotic use, and consequently ABR, among patients in LMICs should principally be focused on primary care. Primary care includes hospital outpatients as well as community pharmacies and drugstores in addition to primary care

clinics in the community. These strategies should be aimed at reducing considerable rates of inappropriate prescribing and dispensing of antibiotics that currently takes place in primary care across LMICs, especially for self-limiting conditions such as minor upper respiratory tract infections (URTIs).<sup>9,15-17,38</sup> Physicians and other prescribers in LMICs often prescribe antibiotics for self-limiting conditions given severe time pressures, and the resultant insufficient time to fully diagnose patients, combined with pressure from patients to prescribe antibiotics.<sup>9,15,39,40</sup> Within primary care in LMICs, nurses and other healthcare professionals (HCPs), including clinical officers, are also often involved alongside physicians in the prescribing of antibiotics in public healthcare clinics (PHCs).<sup>41-44</sup> Similar to physicians, there have been concerns with their knowledge and prescribing practices regarding antibiotics.<sup>45-47</sup>

Many people across LMICs also self-purchase antibiotics without a prescription for themselves or their children in both the formal and informal sectors despite legislation.<sup>16,17,38</sup> High rates of self-purchasing are driven by a number of factors.<sup>16,17,48</sup> Key factors include high patient copayments, limited knowledge of antibiotics, long waiting times to see HCPs in PHCs, shortages of pertinent medicines in clinics, as well as the convenience and availability of community pharmacies, drugstores and drug sellers especially in rural areas.<sup>16,17,49-51</sup> There are though appreciable concerns regarding the knowledge and communication skills of pharmacists and their assistants, as well as among informal drug sellers, with respect to ABR and AMR.<sup>17,52-54</sup> However, pharmacy personnel can play a key public health role in improving antibiotic use in the community, which builds on their role during the recent COVID-19 pandemic.<sup>55-57</sup> The increased role of community pharmacists is reflected by some countries, including LMICs, allowing pharmacists to dispense certain antibiotics without a prescription.<sup>17,58,59</sup> The lack of knowledge regarding antibiotics, their use, ABR and AMR, as well as key aspects surrounding antimicrobial stewardship, among physicians, pharmacists, nurses and other HCPs, usually starts in college or university and can continue post qualification unless adequately addressed.<sup>17,60-62</sup> This is a particular concern in LMICs compared with high-income countries such as the UK.<sup>62,63</sup>

As mentioned, patients and clients also play an appreciable role in LMICs regarding the overuse and misuse of antibiotics through requesting or suggesting antibiotics to HCPs, including

pharmacists, for often self-limiting conditions.<sup>15,64–67</sup> Requests for antibiotics are often driven by limited knowledge of antibiotics and ABR.<sup>15,65,66</sup> Educational campaigns can be effective in LMICs to improve the knowledge of patients and reduce their requests for antibiotics for self-limiting viral conditions, although there are challenges.<sup>68–70</sup> There can also be language barriers to full comprehension, as well as a lack of a specific word or language about infection and its treatment in LMICs, which can make it difficult for pharmacists and their assistants to fully convey concepts such as the actions of antibiotics, ABR and AMR, unless they are aware of possible language issues in advance in their deliberations with patients.<sup>70–72</sup>

Antimicrobial stewardship programmes (ASPs) are being increasingly encouraged across sectors, including primary care, to improve future antibiotic use.<sup>73–77</sup> There have been challenges with developing ASPs in LMICs because of resource and personnel issues.<sup>77,78</sup> However, this is changing across LMICs and sectors, enhanced by groups such as the British Society of Antimicrobial Chemotherapy, the WHO, the Fleming Fund and the Open University providing high-quality training on AMR and ASPs as well as improving connectivity, with published ASPs in primary care also providing exemplars.<sup>68,78–87</sup> There is also growing use of hub and spoke models as part of ASPs, and this will continue.<sup>88</sup>

We are also seeing the growing use of quality indicators across LMICs, including adherence to current guidelines, to improve antibiotic use across sectors.<sup>81,89–92</sup> The use of quality indicators to improve future antibiotic use is likely to accelerate with the recent publication of the WHO AWaRe book providing treatment guidance for a range of common infections.<sup>14,93,94</sup> There can though be challenges with the introduction, implementation and monitoring of quality indicators in LMICs.<sup>95</sup>

Consequently, given rising AMR rates across LMICs, there is a need to ascertain the current situation regarding key stakeholder knowledge, attitudes and practices (KAP) towards antibiotics, AMR and ASPs in primary care. Current KAPs are important as misinformation or poor attitudes and knowledge towards antibiotics and AMR, including lack of knowledge regarding local AMR patterns and the implications for antibiotic dosing regimens, as well as concerns with language, will further increase AMR unless addressed.<sup>70,96,97</sup> We recognize it is challenging to change attitudes as well as prescribing and dispensing practices; however, this is changing.<sup>17,71,81,82</sup> Changes in attitudes and practices are important to achieve the United Nations General Assembly (UNGA-AMR) goals to increase the use of Access antibiotics to 70% across sectors, and combined with other activities to reduce AMR.<sup>98</sup>

As a result, the objective of this narrative review is to document current KAP among all key stakeholder groups in LMICs where AMR is a particular concern. Subsequently, to use the findings to provide future direction to key policy makers, academics, HCPs and others to improve future antibiotic use in primary care in LMICs, including potential behavioural interventions.<sup>83</sup> Instigating activities to improve antibiotic use in LMICs builds on the WHO GAP initiative in 2015 to raise awareness and encourage best practices among HCPs, policy makers and the public through undertaking surveys using instruments provided by the WHO.<sup>99</sup> In this study, we used a narrative review to understand KAP surrounding AMR and the use of antibiotics in four critical WHO Regions during the past 7 years (2018–2024).

## Methodology

### Study design

A narrative review approach was undertaken for this study, similar to other reviews of KAP studies and their implications among LMICs.<sup>65</sup> The rationale for our adopted approach was to allow for a broader scope of sourced papers given that information contained within identified papers may be part of a broader paper, whose insight and findings may be missed when undertaking a systematic review. Examples include, for instance, knowledge and attitudes towards antibiotics, ABR and ASPs among patients visiting community pharmacies or drugstores for antibiotics to treat their infection without initially seeking help from physicians or nurses.<sup>17,38,100</sup> In addition, parts of studies can be included, for example, when assessing point-of-care testing for infections,<sup>101</sup> or adherence rates to current standard treatment guidelines (STGs),<sup>102,103</sup> alternatively, evaluating the impact of ASPs in ambulatory care where pertinent.<sup>104</sup> The greater flexibility and coverage of possible papers also helps provide robust future direction to all key stakeholder groups. LMICs were chosen since, as mentioned, they face the greatest challenges with ABR and AMR, particularly for African and Asian countries.<sup>1,105</sup>

This narrative review builds on previously published systematic reviews assessing KAPs of key stakeholder groups, which have included studies published appreciably earlier than 2018.<sup>61,70,96,106–108</sup> As a result, this review focuses on more up-to-date findings, not just on antibiotic use but also surrounding ABR, AMR and AMS in one comprehensive review. We have adopted similar approaches previously when documenting, as well as suggesting, activities to improve the care of patients with infectious disease across sectors in both Africa and Asia to minimize AMR.<sup>17,29,91,109</sup> Consequently, we believed this approach was more suitable for this comprehensive review, and summarizing the resulting implications, as the principal aim of this paper is to suggest potential ways forward for all key stakeholder groups to improve their KAP surrounding antibiotics, AMR and ASPs, besides acknowledging the difficulties involved with improving future prescribing and dispensing of antibiotics in LMICs. The suggestions will be based on the considerable experiences of the co-authors, similar to other narrative reviews undertaken by the group.<sup>17,91,109,110</sup>

A number of databases were initially searched by the corresponding author (B.G.), including Google Scholar and PubMed/MEDLINE for published papers between January 2018 and October 2024. Search terms included ambulatory care; antibiotics; antibiotic prescribing; primary health facilities; primary healthcare centres; community pharmacists; patients; antimicrobial stewardship; antimicrobial stewardship programs; and KAP among the targeted African and Asian countries within the four key WHO Regions. These are the WHO African, Eastern Mediterranean, South-East Asia and Western Pacific Regions.<sup>111</sup> The references of sourced papers, especially those published from January 2019 to October 2024, were also examined for potential additional references that could be included in the narrative review to enhance its comprehensive nature. The co-authors from across the four WHO Regions also provided additional relevant papers to enhance the review. As a measure of quality, only the sourced papers that are cited in PubMed are included in the tables.




Figure 1. Synopsis of the literature review methodology.

Sourced papers concerning KAP regarding antibiotics, ABR, AMR and AMS in primary care, including hospital outpatients, were divided into the four key WHO Regions for comparative purposes (Figure 1). After this, the results were further divided into individual countries in each WHO Region according to their current World Bank economic status (Figure 1). The current economic status includes three groups: low-, low-middle- and upper-middle-income countries.<sup>112</sup> This breakdown has been undertaken since there can be differences in utilization patterns and challenges between income levels due to many issues including access and affordability of care.<sup>15</sup> The main stakeholders within each country have been further divided into four groups for comparative purposes: (i) prescribers including physicians and nurses, (ii) dispensers including pharmacists and drug sellers, (iii) students including both healthcare and non-healthcare students, and (iv) patients and the public.

**Inclusion and exclusion criteria**

Within the LMICs, given the high numbers of papers reviewed, we concentrated on countries in Africa, the Middle East and Asia in the four WHO Regions where concerns with antibiotic utilization patterns and ABR levels are at a critical point.<sup>1,7,11</sup>

Only papers dealing with primary care, including hospital outpatients, and published from 2018 onwards until November 2024 were eligible for inclusion in this review. Primary care was selected as the focus for this review because it currently accounts for up to 90% of antibiotic use in humans in LMICs.<sup>37</sup> The year 2018 was taken as the early cut-off for sourced papers, with NAPs only launched from 2017 onwards following the WHO GAP in 2015; however, we recognized ongoing concerns with their implementation.<sup>22,29,33,113</sup>

We are aware that the extensive social restriction measures introduced to combat COVID-19, which included closure of universities, had a profound impact on the education of HCPs across LMICs.<sup>114–116</sup> However, we have not separated out papers dealing specifically with COVID-19 and its implications, principally treating COVID-19 as another viral infectious disease as patients with COVID-19 were often inappropriately treated with antibiotics, especially across LMICs.<sup>117–119</sup>

Exclusion criteria included studies discussing KAPs of key HCPs treating inpatients in hospitals as well as LMICs outside the four designated WHO Regions. We also excluded papers published before 2018, those exclusively dealing with animals/veterinary scientists, and any paper not cited in PubMed (for inclusion in the tables).

**Data analysis and narrative synthesis of findings**

The data analysis process followed a systematic approach to identify, extract and organize relevant information from the studies, including information about the studies’ characteristics and outcomes of interest. Activities included creating tables containing a summary of the objectives, methodology and key findings by each key stakeholder group (prescribers, dispensers, students and patients/public) as well as by country and region. Furthermore, the documented synopsis for each cited paper included the country, its economic status and the study design as well as the number of participants. Details of any questionnaire design were also included where documented as a key element of the paper and its findings, especially how the questionnaire was developed including any pilot studies as well as any validation. The publication year along with the first author was also included in the tables, with the papers listed in year sequence to assess changes over time where pertinent (Figure 1). Key findings were also documented as part of the narrative synthesis led by the principal author (B.G.). Key findings included any differences between perceived knowledge of antibiotics, ABR, AMR and AMS, among the stakeholder groups versus actual practices, especially in terms of prescribing, dispensing or using antibiotics for essentially self-limiting conditions, driving up ABR.

The emerging themes from each stakeholder group across the LMICs and the four WHO Regions were subsequently synthesized into major themes for comparisons within and across the four WHO Regions. No attempt was made to combine the findings and produce any summary statistics apart from documenting the number of sourced papers by country and WHO Region in view of the recognized heterogeneity of the sourced papers in terms of the populations studied and the methodology. However, key reported quantitative descriptive statistics or



measures of effect, such as prevalence rates or mean/median scores related to specific study outcomes or thematic areas, were summarized. This approach was employed to capture the anticipated variability across different studies and contexts, particularly given the diverse methodologies and populations examined across LMICs, as well as key differences between stated knowledge regarding antibiotics and AMR and actual practice. The use of descriptive statistics, including ranges where pertinent, allows for a broader representation of the findings, accounting for regional and contextual differences while providing an aggregated view of key outcomes, including KAP towards antibiotic use, ABR and AMS. Where relevant, the figures are discussed in the context of each theme to highlight variations and commonalities across the different studies and regions.

The consolidated themes were subsequently used to structure a range of future potential interventions to improve KAP in all key stakeholder groups, summarizing key activities to instigate and improve antibiotic use in primary care.

### Ethical considerations

Ethical approval was not needed for this review as this study did not involve direct contact with humans as noted in similar studies.<sup>9,17,82,109</sup>

## Results

The general findings will be discussed first before synthesizing the findings by each stakeholder group and region.

### General

A total of 459 papers were sourced across the LMICs that met the inclusion criteria. These have been divided into the four key stakeholder groups across the four WHO Regions (Tables S1 to S16, available as [Supplementary data](#) at JAC-AMR Online).

An appreciable number of sourced papers came from the African Region in all four stakeholder categories, representing 42.7% of all published studies, with most coming from Ghana, Nigeria, South Africa and Tanzania (Table 1). The number of papers is perhaps not surprising, with the greatest burden of infectious diseases worldwide, including HIV and AIDS, acute respiratory diseases, malaria and TB, currently in Africa.<sup>120,121</sup> Alongside this burden, there are high and growing rates of AMR among African countries.<sup>105,120–122</sup> The appreciable number of publications emanating from China (8.1% of total publications) may reflect high rates of inappropriate antibiotic prescribing and dispensing in the country, encouraged by limited patient knowledge, despite recent government initiatives, along with high rates of AMR.<sup>123–127</sup> This may be changing; however, there is a continued need to encourage appropriate prescribing and dispensing of antibiotics to reduce AMR in the country.<sup>128–130</sup>

The considerable number of papers documenting the KAP of the patients/public (33.1% of the total number of publications within each stakeholder group—ascertained by documenting the number of papers cited in Tables S13 to S16 by the total number of cited studies in Tables S1 to S16) reflects their appreciable role in influencing the prescribing and dispensing of antibiotics in LMICs (Table S20).

Differences have been seen between high-income countries and LMICs, especially with respect to antibiotic use patterns, including the self-purchasing of antibiotics, which is appreciably greater in LMICs, as well as educational campaigns among patients.<sup>11,15,48,131–135</sup> However, concerns about the knowledge of physicians with respect to antibiotics and AMR have been seen across all countries irrespective of income level.<sup>136</sup> Initially, educational campaigns among patients to improve their knowledge were principally undertaken in high-income countries.<sup>15,137,138</sup> However, this is now changing.<sup>68,139–143</sup>

The findings from the four key stakeholder groups across the four WHO Regions are summarized in Tables S17 to S20. The consolidated findings have subsequently been expanded to provide additional background to potential interventions that can be implemented by key stakeholders going forward. This includes documenting concerns and issues with the KAP of each stakeholder group across the four Regions, alongside any appreciable differences between stated knowledge regarding antibiotics and ABR and actual practice. The latter is seen as particularly important when it comes to suggesting future activities.

### Summary of key findings among each stakeholder group across the four WHO Regions

#### Prescribers

Overall, reported rates of antibiotic prescribing in primary care were broadly comparable across the LMICs in the four WHO Regions independent of World Bank income levels (Table S17).

High rates of prescribing of antibiotics were noted among physicians (range: 23%–97%) as well as nurse practitioners and clinical officers (range: 35.8%–69%) across the four WHO Regions. However, in some of the LMICs, a higher rate of prescribing of antibiotics was seen among physicians compared with other HCWs.<sup>46,144</sup> Private health professionals and informal providers also tended to have higher rates of antibiotic prescribing (range: 26.2%–64.3%) across the four WHO Regions. This though was not always the case.<sup>35,42</sup>

Antibiotics were often prescribed for self-limiting conditions, including acute respiratory infections (up to 97%), across the four WHO Regions. This was exacerbated by the lack of time that prescribers typically have with patients in PHCs coupled with patient demands for antibiotics across the LMICs. However, there were exceptions among some countries. This was particularly the case in the Eastern Mediterranean Region where a number of prescribers believed antibiotics were not effective against viral infections, with the body able to fight mild infections without the need for antibiotics.<sup>68,145–147</sup> However, this was not always the case in this Region, especially where a diagnosis or indication was lacking.<sup>52,148,149</sup>

There were also variable adherence rates to current guidelines among LMICs in the four WHO Regions. This is important with adherence to guidelines increasingly seen as an important marker of the quality of care provided.<sup>9,89,90</sup> Where recorded, compliance rates varied across the WHO African (45.1%–94.9%), Eastern Mediterranean (32.5%–69.5%), South-East Asia (31.9%–98.5%) and Western Pacific Regions (29.3%–93.37%). As a result, in some LMICs and Regions, there were high rates of prescribing of Watch antibiotics.<sup>144,150–152</sup> This included the African Region,

**Table 1.** Breakdown of published studies by WHO Regions and stakeholder groups<sup>a</sup>

WHO Region	Prescribers	Dispensers	Students	Patients/public
<b>African Region</b>				
Countries	23	13	12	18
Published studies	90 (40.7%)	40 (18.1%)	27 (12.2%)	64 (29.0%)
Top countries with published studies	• South Africa: 23 studies • Tanzania: 12 studies	• South Africa: 7 studies • Ethiopia and Tanzania: 5 studies	• Nigeria: 7 studies • Uganda: 4 studies	• Ghana: 11 studies • Ethiopia, Nigeria, Tanzania: 7 studies each
<b>Eastern Mediterranean Region</b>				
Countries	6	9	7	12
Published studies	19 (18.8%)	28 (27.7%)	20 (19.8%)	34 (33.7%)
Top countries with published studies	• Jordan: 5 studies • Pakistan: 4 studies	• Pakistan: 13 studies • Egypt: 4 studies	• Pakistan: 7 studies • Jordan: 5 studies	• Pakistan: 9 studies • Jordan: 7 studies
<b>South-East Asia Region</b>				
Countries	8	6	5	6
Published studies	29 (26.6%)	34 (31.2%)	17 (15.6%)	29 (26.6%)
Two top countries	• India and Nepal: 8 studies each	• Bangladesh: 12 studies • Nepal: 11 studies	• Bangladesh, India, Nepal and Sri Lanka: 4 studies each	• Bangladesh: 11 studies • India: 7 studies
<b>Western Pacific Region</b>				
Countries	5	4	2	5
Published studies	27 (30.7%)	8 (9.1%)	10 (11.4%)	43 (48.8%)
Two top countries	• China: 13 studies • Vietnam 5 studies	• China: 3 studies • Vietnam: 3 studies	• China: 6 studies • Malaysia: 4 studies	• China: 20 studies • Malaysia and Vietnam: 8 studies each

<sup>a</sup>Percentage values in brackets of the total number of papers from each Region for each stakeholder group. The total number of documented studies at 518 is more than 459 as some studies included different countries and stakeholder groups.

where up to 55.3% of antibiotics prescribed in one country and sector were from the Watch group;<sup>153</sup> in the South-East Asia Region up to 54.9% of antibiotics in India in one study were from the Watch group;<sup>154</sup> and in the Western Pacific Region the rate was up to 31% or more.<sup>155</sup>

There were also concerns with knowledge regarding antibiotics, ABR, AMR and AMS among prescribers across the four WHO Regions irrespective of World Bank income levels. However, there was variability regarding the extent of concerns for these key issues both within and across the Regions. Typically, qualified physicians had greater knowledge of antibiotics, ABR and AMR than other prescribers, including unqualified prescribers. Terminology was also an issue in some countries.

Another key identified concern was that although in some LMICs there was good knowledge of antibiotics, ABR and AMR among prescribers, this typically did not always translate into actual prescribing practices with, as mentioned, typically excessive prescribing of antibiotics across the four WHO Regions. Excessive prescribing of antibiotics includes prescribing them essentially for viral infections such as URTIs as well as those from the Watch list where Access antibiotics or no antibiotics initially are more appropriate. For instance, in the African Region in Sierra Leone in one study, physicians demonstrated sound knowledge and attitudes regarding antibiotics and ABR; however, 68% of them believed that the prescribing of antibiotics may speed up recovery from a cold or cough.<sup>156</sup> In Nigeria, 49.2% of prescribers had good

knowledge of ABR and AMR; however, 75.7% admitted prescribing antibiotics for sore throats.<sup>157</sup> Alongside this in South Africa, 95.8% of prescribers in one study believed ABR is a major problem; however, 66.5% felt pressure from patients to prescribe antibiotics for their presenting infectious disease irrespective of their potential effectiveness for the presenting infectious disease and the need.<sup>158</sup>

In the Eastern Mediterranean Region in Palestine in one study, despite only 9.2% of participating physicians believing antibiotics can be used to treat viral infections, 60.8% would prescribe antibiotics for acute sinusitis, and 43.4% for a 17-year-old complaining of a sore throat or nonproductive cough.<sup>159</sup> In Egypt in one study, 98.7% of participating physicians believed bacteria can become resistant to antibiotics; however, 61.4% believed that broad-spectrum antibiotics are preferred for treating URTIs.<sup>146</sup> In the South-East Asia Region in one study in India, although over 90% of prescribers claimed their antibiotic knowledge was adequate, over 88% of physicians reported prescribing antibiotics for viral infections such as common colds or sore throats.<sup>46</sup> In Sri Lanka, 99.6% of participating physicians in one study stated that antibiotics should not be prescribed to any patient with fever, but 52.3% almost always/often/sometimes felt under pressure to prescribe antibiotics if patients expect these.<sup>160</sup> In the Western Pacific Region, for instance in Lao, 91% of physicians stated that extensive use of antibiotics increases the risk of ABR, but 39% would prescribe

an antibiotic to an adult with a runny nose, cough or fever to hasten recovery.<sup>161</sup>

The need for enhanced training regarding antibiotics, the AWaRe classification, ABR and AMS, was seen as a definite requirement among all prescribers, including physicians and nurses, across the LMICs in the four WHO Regions to improve future prescribing. This also included training surrounding current national STGs, and with the recent publication of the WHO AWaRe book where no current national STGs exist.<sup>27,28</sup> This is important with cited studies (Tables S1 to S4) showing that educational and other interventions surrounding STGs increased the appropriateness of antibiotic prescribing.<sup>68,135,162–165</sup> In addition, prescribers who had access to STGs were less likely to prescribe antibiotics.<sup>166</sup> Multiple initiatives in China in recent years have also enhanced appropriate prescribing of antibiotics with improved training.<sup>128,129</sup>

### Dispensers

A similar picture to that for prescribers was seen among dispensers across the four WHO Regions (Table S18), with again the findings irrespective of World Bank income levels. There were typically high rates of dispensing of antibiotics without a prescription across the four WHO Regions where rates were documented. Rates of self-purchasing included up to 100% of patients visiting community pharmacies in the African and Eastern Mediterranean Regions,<sup>167,168</sup> up to 94% among community pharmacies and drug sellers in the Western Pacific Region,<sup>169</sup> and up to 86.7% among patients visiting community pharmacies in the South-East Asia Region.<sup>166</sup>

Alongside this, similar to the prescribers across LMICs, the dispensing of antibiotics without a prescription was typically for self-limiting conditions including URTIs, acute respiratory infections (ARIs) and acute diarrhoea as well as for aches, pain and inflammation. Identified concerns going forward also included appreciable dispensing of antibiotics from the Watch group in some of the studied LMICs. High rates of dispensing of azithromycin were also seen among patients with COVID-19 (99.1%) in Kenya.<sup>170</sup> Watch antibiotics also accounted for 53.6% of antibiotics dispensed among patients visiting community pharmacists in Bangladesh,<sup>171</sup> and for 49.3% of antibiotics dispensed among patients visiting community pharmacies in Pakistan.<sup>172</sup> Pharmaceutical company promotional activities also enhanced the dispensing of azithromycin without a prescription in Nepal.<sup>173</sup> There were also high rates of dispensing of antibiotics from the Reserve group among community pharmacies in Bangladesh (10%) and Pakistan (19.0%).<sup>171,172</sup>

Overall, the reasons for high rates of dispensing of antibiotics without a prescription where this occurs were similar across the LMICs and WHO Regions irrespective of World Bank income status. Key reasons included high patient copayments, convenience of pharmacies, ease of access to antibiotics in community pharmacies and drugstores as well as pressure from patients or carers to dispense.<sup>52,168,174–176</sup> If anything, rates of dispensing of antibiotics without a prescription were higher in drug outlets versus community pharmacies, and for children versus adults.<sup>176–183</sup> However, this was not always the case.<sup>184</sup>

Not all countries within the four Regions experienced high rates of dispensing of antibiotics without a prescription. There

were stricter regulations and greater access to HCPs in some of the LMICs, for example Botswana and Namibia in Africa and Malaysia in South-East Asia. In these countries, there are strict regulations surrounding the dispensing of antibiotics without a prescription, which is typically enforced, alongside activities of pharmacists to offer alternatives through training.<sup>17,57,185</sup> There was also limited dispensing of antibiotics without a prescription to patients for essentially viral infections in community pharmacies linked to the University of Nairobi in Kenya.<sup>186,187</sup>

Pharmacists with greater experience, and following educational activities, were generally less likely to sell antibiotics without a prescription.<sup>68,188,189</sup> However, this was not always the case.<sup>190</sup> Similar to prescribers, although knowledge of antibiotics, ABR and AMR appeared reasonable to good among dispensers in a number of LMICs, this again did not always translate into actual practice. Dispensers across the four WHO Regions typically recognized their own knowledge and skills gap. They also acknowledged issues of language surrounding the terminology for antibiotics, ABR and AMR as a barrier to reducing inappropriate dispensing of antibiotics for essentially self-limiting conditions.<sup>69–71,191</sup> Real-time feedback to community pharmacists using mobile and other technologies could also facilitate improved dispensing alongside improved adherence to standard guidance including treatment guidance in the recently launched WHO AWaRe book.<sup>27,28</sup>

### Students

There were again similar issues regarding the knowledge of healthcare students concerning antibiotics, ABR, AMR and AMS across the LMICs in the four WHO Regions. There were also similar concerns among the countries and Regions irrespective of country income levels (Table S19).

Good knowledge regarding antibiotics among students was seen in some studies in the African Region. In Zambia, 87.3% of participating medical students had good knowledge regarding antibiotic use and AMR, with 99.2% believing antibiotics can cure bacterial infections and 93.5% that antibiotics cannot cure viral infections.<sup>192</sup> In Malawi, over 88% of surveyed students in one study correctly answered more than half the questions regarding antibiotics and AMR.<sup>193</sup> In addition, in Nigeria 98.4% of surveyed students in one study believed AMR was an important global public health issue,<sup>194</sup> with 94.8% in another study believing that antibiotics are overused in their country.<sup>195</sup> In Rwanda, 98.7% of participating students had heard about AMR, and 78.6% had discussed AMR during their courses, with 95.2% believing that inappropriate use of antimicrobials increases AMR.<sup>196</sup> In Uganda, 87.5% of surveyed students had sufficient knowledge about AMR.<sup>197</sup> In another study among universities in East Africa, 72.9% of participating students had a good attitude and perception regarding antibiotic use. The extent of good attitudes was highest in Uganda versus Kenya and Tanzania.<sup>198</sup> There was also typically greater knowledge regarding antibiotics, ABR and AMR among final-year students versus students with less experience, and among healthcare versus non-healthcare students across studies.<sup>192,193,199,200</sup>

Despite this, there were generally considerable concerns regarding the attitude and practice towards antibiotics and AMR among the surveyed students across Africa.<sup>41,194,201–204</sup>

This was also reflected by high rates of self-purchasing of antibiotics without a prescription, including for self-limiting conditions such as coughs, colds, sore throats and diarrhoea, among students across Africa.<sup>201,203,204</sup> Rates of 55.3% to 93.8% for self-purchasing without a prescription were seen in studies including among students from Ghana, Nigeria, Tanzania and Uganda.<sup>108,199,201,203,205,206</sup> These high rates of self-purchasing were driven by easy access and convenience of pharmacies, perceived only minor illnesses, the costs involved, including cost savings with self-medication, and prior knowledge.<sup>207–209</sup>

A similar situation was seen among students in the Eastern Mediterranean Region. In Palestine in one study, 97.4% of students stated they knew about AMR and 86.7% that the overuse of broad-spectrum antibiotics increases AMR.<sup>210</sup> In Egypt in one study, 95.7% of surveyed students believed indiscriminate use of antibiotics increases ABR and AMR,<sup>211</sup> and in Jordan, 93% of students in one study stated they knew that unnecessary use of antibiotics makes them ineffective, with 85.5% stating that antibiotics are ineffective against viruses.<sup>212</sup> In Pakistan, 85.7% of students in one study stated that they knew about AMR, with 79.4% agreeing that a contributing factor for AMR was poorly designed dosing regimens.<sup>213</sup>

However, despite these encouraging signs, there was again appreciable concerns with the overall knowledge of students regarding antibiotics, ABR and AMS across this Region. This was reflected in high rates of self-purchasing of antibiotics among students in a number of the countries. Rates of self-purchasing of antibiotics ranged from 38.1% to 60.8% among surveyed students in the Region.<sup>214–216</sup> There was also a high percentage of students who believed that antibiotics are effective and/or first line in treating self-limiting viral infections including coughs, colds and influenza.<sup>211,217–219</sup> In the Middle East, including Egypt, Jordan and Lebanon, 51.7% of pharmacy students reported taking antibiotics to manage a fever.<sup>216</sup> In Pakistan in one study, 51.4% of surveyed students believed antibiotics can be taken as preventative measures.<sup>220</sup> Among Arabic-speaking countries, including Jordan, Iraq and Palestine, 44.5% of students believed antibiotics are effective against colds and coughs,<sup>221</sup> and in another study involving students in Palestine, 66.1% were not confident about AMS.<sup>210</sup> Alongside this, 43.3% of students in one study in Egypt believed that ABR is only a problem if antibiotics are taken regularly.<sup>217</sup>

Similar to the situation in the African Region, there was generally greater knowledge regarding antibiotics, ABR and AMR among more mature students in the Eastern Mediterranean Region, and with healthcare compared with non-healthcare students.<sup>213,216</sup>

Similar to the other Regions, appreciable differences were seen between knowledge and practice among countries in the South-East Asia Region. In India, 91.8% of students in one study believed antibiotics were useful for treating bacterial infections,<sup>222</sup> with 84.1% and 82.4% of students, respectively, in another study believing inappropriate use of antibiotics, and not completing the full course, increases ABR.<sup>223</sup> A study in Bangladesh found that 91.7% of students believed that dispensing antibiotics without a prescription increases ABR, with 82%

believing the overuse of antibiotics increases ABR.<sup>224</sup> In one study in Nepal, 83.3% of students believed that ABR means that antibiotics are taken too often and will not work.<sup>225</sup> In three Asian countries, including Indonesia, 80.1% of participating students in one study recognized it was inappropriate to treat diarrhoea with antibiotics, with 75.8% strongly disagreeing or disagreeing that AMR is not a serious problem.<sup>226</sup>

However, again there were appreciable concerns regarding the practical knowledge of students concerning antibiotics, ABR, AMR and AMS. In Nepal in one study, 73.7% of participating students thought that antibiotics can cure infections caused by viruses.<sup>227</sup> In Indonesia, only 26.1% of participating students in one study stated it was inappropriate to treat URIs with antibiotics.<sup>226</sup> In Sri Lanka, between 40.2% and 63% of surveyed students in various studies believed antibiotics could cure colds, sore throats and influenza.<sup>228–231</sup> In Bangladesh, 44.4% of students in one study believed that antibiotics could prevent viral infections.<sup>224</sup> There was also appreciable self-purchasing of antibiotics among students, including 67.78% of students in one study in India,<sup>232</sup> up to 61.7% of students in Nepal,<sup>233,234</sup> and 42.2% to 91.7% in Bangladesh.<sup>224,235,236</sup> However, this was not always the case among the Regions.<sup>237</sup> Self-purchasing of antibiotics among students in this Region was again typically for self-limiting conditions including fevers and ARIs.<sup>234–236,238</sup> Principal reasons for self-medication with antibiotics among students again included convenience, long waiting times to see physicians, previous experiences with antibiotics, the condition is not serious, and costs.<sup>232,236</sup>

Mirroring other Regions, there again appeared greater knowledge regarding antibiotics, ABR and AMR among healthcare and biology students versus other students in the South-East Asia Region as well as among more mature students.<sup>224,227–229</sup>

Lastly, in the Western Pacific Region there were again differences between perceived knowledge and actual practice among students. In one study in China, more than 85.0% of surveyed students were aware that the overuse of antibiotics is a serious problem that could cause AMR as well as lead to difficulties in treating bacterial infections.<sup>239</sup> In other studies, 74.2%–88.5% of students reported being aware of the dangers posed by the overuse of antibiotics.<sup>240,241</sup> In one study in Malaysia, more than 88% of surveyed students totally agreed or agreed that penicillin or amoxicillin are antibiotics, and that antibiotics are useful for treating bacterial infections, with 77.5% also stating they do not take antibiotics to treat a cold or sore throat.<sup>242</sup>

In contrast, only 47.0% of surveyed students in one study in China did not agree that antibiotics could reduce the symptoms of common colds,<sup>239</sup> and in another, 75.2% of students stated they had performed inappropriate antimicrobial practices.<sup>243</sup> Another study in China reported that 54.3% of students were unaware that antibiotics were ineffective against viral infections, with only 41.6% agreeing that every person treated with antibiotics is at an increased risk of antibiotic-resistant infections.<sup>237</sup> There was a similar situation in Malaysia, with only 3.3% of surveyed students in one study having a good level of knowledge about antibiotics,<sup>244</sup> in another study, only 26.1% of students, including those from Malaysia, believed it inappropriate to treat



URTIs with antibiotics.<sup>226</sup> Self-medication with antibiotics was also seen in 48.8%–73.9% of students in China,<sup>240,245</sup> and in up to 33.7% of students for a cough in Malaysia.<sup>244</sup>

Similar to other studies, students with a medical background in the Western Pacific Region were significantly associated with better antibiotic use behaviours and knowledge versus other students.<sup>244,245</sup> In addition, more mature students had better knowledge regarding antibiotics, ABR and AMS than junior students.<sup>242,246</sup>

Overall, there was a recognized need among healthcare students across all four WHO Regions, irrespective of World Bank income levels, for greater training on antibiotics, ABR, AMR and AMS during their undergraduate years, with greater training seen as effective with improving knowledge scores.<sup>244,247</sup> There was recognition that HCPs would need further updated training post-qualification with continuing professional development (CPD). This should include practical tools to help with diagnosis and treatment, as well as tools to provide up-to-date information on local resistance patterns to aid future prescribing and dispensing. To this end, training about robust guidelines such as the recent AWARe classification and guidance is seen as important.<sup>13,27,28</sup>

### Patients/public

There were similar concerns and issues to the other key stakeholder groups among patients and the public across the LMICs in the four WHO Regions with respect to their KAP towards antibiotics, ABR and AMR (Table S20).

In the Democratic Republic of Congo in the WHO African Region, all interviewed patients in one study were familiar with the term ‘antibiotics’, with most describing antibiotics as medicines used to treat infections.<sup>248</sup> Similarly in Ethiopia, 95.9% of participants correctly agreed with the definition of antibiotics, with 59.4% of participants having heard the term AMR, and of these 50.4% were aware of the problems of ABR.<sup>249</sup> In other studies in Ethiopia, 58.2% of participants were aware of the existence of ABR, 48% had moderate knowledge of antibiotics and AMR, with 35% having good knowledge.<sup>249–252</sup> In Cameroon, 88.3% of those surveyed in one study stated that antibiotics treat microbes, with only 11.8% believing that antibiotics can be used to treat viral diseases.<sup>253</sup> In Ghana, 82.4% of respondents in one study were aware that the misuse of antibiotics is a major driver of ABR, and 74.4% knew that ABR is a failure of antibiotics to kill germs.<sup>254</sup> In another study in Ghana, 75.9% of participants had good knowledge regarding bacteria’s ability to become resistant to antibiotics.<sup>255</sup> In Tanzania, 96.33% of participants in one study believed that self-medication with antibiotics increases ABR.<sup>256</sup> A similar situation was seen in Nigeria, Senegal, South Africa and Zambia.<sup>257–262</sup> In Nigeria, 76.6% of patients in one study stated that bacteria would become less resistant to antibiotics after prolonged use, with only 22.1% believing that antibiotics cure colds and sore throats faster than other remedies.<sup>261</sup>

However, a number of concerns were identified across the countries in the African Region. In Tanzania in one study, only 10.9% of participating parents or guardians had good knowledge

about antibiotics,<sup>263</sup> and in the Democratic Republic of Congo, antibiotics were being used to treat a range of conditions including fever, coughs and profuse sweating.<sup>248</sup> In Malawi, 92.4% of participants in one study believed antibiotics could stop a fever,<sup>264</sup> and in Senegal, 72.3% believed antibiotics are effective against a cough,<sup>257</sup> with 68% of participants in Tanzania in one study admitting to using antibiotics to treat coughs.<sup>265</sup> Elsewhere in Tanzania, 61.4%–62.5% of participants in different studies believed sore throats could be treated with antibiotics.<sup>266,267</sup> In South Africa, 66% of participants in one study incorrectly stated that antibiotics are effective for treating viruses,<sup>262</sup> with generally a common misconception that antibiotics can treat colds, influenza and fever.<sup>268</sup> Alongside this in Uganda in one study, 60.2% of children with ARIs were treated with antimicrobials,<sup>269</sup> and in Zambia in different studies, 58.2% of participants had taken an antibiotic for a common cold,<sup>259</sup> and 59.1% had used antibiotics inappropriately.<sup>270</sup> Similar findings were seen in Eritrea, Ethiopia, Ghana, Nigeria, Mozambique and different studies in Zambia.<sup>250,271–277</sup>

Among the countries in the WHO African Region, patients also admitted to appreciable self-purchasing of antibiotics without a prescription. Self-purchasing was typically for self-limiting conditions including coughs, colds and fever, which was driven by ease of access to community pharmacies and drugstores and their availability in these locations, previous experience, condition seen as minor, costs, location (e.g. rural area) as well as a desire for quick relief of their infectious diseases.<sup>262,272,274,278–284</sup> Documented rates of self-purchasing of antibiotics among participating patients in the studies varied across the countries. These were 23.6% in Kenya,<sup>285</sup> 38.9% in Ethiopia,<sup>281</sup> 45.1% among participants in Eritrea including for URTIs,<sup>280</sup> 55.4% in Uganda,<sup>286</sup> 58%–76.3% in Tanzania,<sup>256,284</sup> 36%–76% among participants in Ghana,<sup>100,287,288</sup> 74.0% in Zambia,<sup>259</sup> 47%–81.4% in Cameroon,<sup>253,289</sup> 26%–86.5% in Nigeria,<sup>208,290</sup> and 93.75% in Mozambique.<sup>291</sup>

In the WHO Eastern Mediterranean Region, there were again appreciable differences between stated knowledge and practice. In Jordan, 88.3% of participants agreed that antibiotics are being excessively used to treat URTIs,<sup>292</sup> with up to 92.6% of participants in one study believing antibiotics are ineffective against common colds,<sup>293</sup> and only 27% of participants in another study believing antibiotics can be used to treat both bacteria and viruses.<sup>294</sup> There were similar findings in studies in Morocco, Pakistan and Yemen.<sup>64,295,296</sup> For instance in Pakistan, 61.7% of participants in one study had good knowledge regarding antibiotics and AMR and the fact that antibiotics kill bacteria.<sup>64</sup>

Among participants in a study in Afghanistan there was poor knowledge regarding antibiotics, with surveyed patients typically seeing antibiotics as the ‘desired medication’, with high expectations that antibiotics will be prescribed by physicians for their infections whatever the cause.<sup>297</sup> In Yemen, 88.8% of participants in one study stated they had used antibiotics to treat a fever,<sup>295</sup> and in Morocco, 78% of those surveyed in one study believed that respiratory tract infections (RTIs) should be treated with antibiotics.<sup>296</sup> In Jordan, 72.4% of participants strongly agreed/agreed that once a child develops a fever, they should be given

antibiotics regardless of the cause.<sup>292</sup> In Pakistan in one study, the majority of participants believed antibiotics can treat viral infections and that all types of infection can be cured with the help of antibiotics,<sup>64</sup> with overall knowledge regarding ABR seen as poor.<sup>298</sup> Among Arabic-speaking countries including Egypt, Iraq, Jordan and Palestine, 58.8% of surveyed patients thought antibiotics are effective against sore throats, and 57.6% believed they combat common colds, coughs and nasal congestion.<sup>221</sup> Similar findings were also seen in other studies in Jordan,<sup>292,299</sup> Iraq<sup>300</sup> and Pakistan,<sup>52,301,302</sup> as well as in Lebanon and Tunisia.<sup>303–305</sup>

Self-medication with antibiotics among patients was also common in the WHO Eastern Mediterranean Region. Documented rates among patients ranged from 20.6% of participants in Tunisia<sup>303</sup> to 32.1% in Egypt,<sup>306</sup> 33.4%–59% in Pakistan,<sup>301,307</sup> 44% in Jordan,<sup>308</sup> 45.8% in Iraq,<sup>300</sup> up to 56.1% in Sudan,<sup>309</sup> 61.6% in Iran<sup>310</sup> and 68% in Morocco.<sup>296</sup>

Similarly in the WHO South-East Asia Region, there were again considerable differences between stated knowledge and actual practices. In Nepal, 94.1% of participants in one study answered correctly that antibiotics are useful for killing germs, and 84.1% that antibiotics are often not needed for treating colds and influenza.<sup>311</sup> In India, 78.7%–89.7% of participants in various studies agreed that it is the obligation of everyone to use antimicrobials responsibly to reduce AMR.<sup>312,313</sup> In Indonesia, 67.7% of participants in one study correctly stated that inappropriate use of antibiotics will cause ABR,<sup>314</sup> and only 12.5% believed antibiotics are effective against viruses.<sup>315</sup> Similar findings regarding the appropriate and inappropriate use of antibiotics and ABR were seen in studies in Bangladesh and Myanmar.<sup>316,317</sup>

In contrast in Nepal, 84.6% of participants in one study reported sometimes preferring an antibiotic when they have a cough or sore throat,<sup>311</sup> and in another study participants typically demonstrated a low level of knowledge concerning AMR.<sup>318</sup> In Bangladesh in one study, 90.7% of participants believed antibiotics are effective in treating viral infections, and 84.76% thought that antibiotics speed up recovery from most coughs/colds.<sup>316</sup> In addition, community pharmacists in Bangladesh believed that if they advise patients against taking antibiotics for minor illnesses including colds/coughs and fever, or insisting on dispensing antibiotics only with a prescription, patients may take their business elsewhere.<sup>183</sup> In India, there was minimal understanding of the term antibiotic in one study,<sup>319</sup> and in Sri Lanka, overall knowledge among participants about antibiotics was poor.<sup>320</sup> Similar findings were seen in other studies in Bangladesh and Nepal,<sup>321,322</sup> as well as in India, Indonesia and Myanmar.<sup>313,314,317,323</sup>

Again, self-purchasing of antibiotics was reported by patients across the WHO South-East Asia Region. Reported rates ranged from 11% in Sri Lanka,<sup>320</sup> up to 50.9%–80.9% in Bangladesh, which included high rates of dispensing of antibiotics from the Watch list.<sup>171,324</sup> The top three infectious diseases for self-medication with antibiotics in Bangladesh were coughs, colds and a fever.<sup>325</sup> In one study in India, participants stated that they typically go to informal providers as a result of economic conditions, with coughs and colds typically treated with antibiotics for 3 to 5 days.<sup>326</sup>

Similarly, in the WHO Western Pacific Region there were again appreciable differences between stated knowledge regarding

antibiotics, ABR and AMR and actual practices. In China, 83.5% of participants in one study believed that antibiotics are not necessary for common colds, and 70.2% averred that overuse of antibiotics increases ABR, with similar findings seen in an additional study (62.8% of participants).<sup>327,328</sup> In Cambodia, 74.8% of participants in one study believed that AMR will make infections more difficult to treat,<sup>329</sup> and in Malaysia in one study, the majority of respondents had good knowledge of antibiotic usage and ABR problems.<sup>330</sup> Similar findings regarding good attitudes towards antibiotics were also seen in other studies in China, Malaysia and Vietnam.<sup>331–333</sup>

However, common uses of antibiotics among patients in Lao and Thailand included for coughs (30.5%), fevers (30.5%) and sore throats (28.9%).<sup>70</sup> In Vietnam, only 18.8% of participants in one study knew that ABR has a negative effect on antibiotics in terms of reducing their effectiveness.<sup>334</sup> Frequent and indiscriminate use of antibiotics is common among patients in Vietnam, driven by the powerful appeal that antibiotics have, with 67.9% of participants in one study believing that antibiotics can be used to treat viruses.<sup>335,336</sup> Similarly in China, 91.6% of participants in one study believed that antibiotics can control viruses,<sup>337</sup> with 81.3% of participants in another study admitting to using antibiotics to treat their RTIs.<sup>338</sup> In Malaysia, studies among patients also documented a high percentage believing antibiotics can hasten recovery from viral infections, or cure these, ranging from 41.5% to 83.7% of surveyed participants.<sup>339–342</sup> In other studies in Malaysia, only 20.6% to 23.8% of participants knew that antibiotics are not effective against viral infections.<sup>330,342</sup> Similar findings were also seen in other studies in Cambodia, China, Lao and Vietnam.<sup>327–329,331,343–348</sup>

Self-purchasing of antibiotics was also common in the WHO Western Pacific Region. Drug suppliers in Vietnam stated that patients often request specific antibiotics to treat their infections based on previous experiences.<sup>169</sup> Overall reported rates of self-purchasing among patients ranged from 10.3% to 92.1% in China, depending on the population and location,<sup>123,127,327,332,337,348–353</sup> 15.1% in Malaysia,<sup>341</sup> and up to 81.7% in Vietnam.<sup>354</sup>

Overall, patients' and parents' beliefs regarding the effectiveness of antibiotics across a range of self-limiting infectious diseases are a major challenge across the four WHO Regions as they have resulted in considerable pressures on both prescribers and dispensers to prescribe/issue antibiotics in situations where they are not needed and could cause harm (Table S20). Generally, though, there was a better attitude towards antibiotics among more educated patients.

Encouragingly, education initiatives across the Regions did appear to appreciably improve mean knowledge scores and the practical use of antibiotics in terms of reducing their use for essentially viral infections.<sup>68,139–143</sup> In addition, in the study of Otieku *et al.*,<sup>355</sup> participants exposed to an educational intervention to increase their knowledge base were significantly more likely to recommend restrictive access to antibiotics. However, there can be challenges with undertaking educational campaigns among patients.<sup>356</sup> Issues and concerns with patients' understanding of the terminology regarding antibiotics and ABR were recognized as needing to be addressed, especially by dispensers, when patients or their care givers are educated regarding the

appropriate management of their infectious diseases.<sup>69–71,357</sup> Understanding these issues is crucial when health authorities and others are designing educational initiatives for patients.<sup>358,359</sup> However, it was noted that any intervention or targeted programme must include a range of stakeholders to ensure these are appropriate for the health system, community and cultural context in question to enhance the chances of success.<sup>360</sup>

## Discussion

We believe this to be the most comprehensive narrative synthesis to date of KAPs across multiple stakeholder groups in LMIC primary care settings. The wide range of priorities identified in the reviewed papers have been broken down into different key stakeholder groups: governments/health authorities, prescribers (physicians/nurses), dispensers (principally community pharmacists), universities/students and patients/public. We are aware that dispensers also include the informal sector in many LMICs, e.g. Bangladesh, Malawi and Zimbabwe. However, in this section we will deal just with activities with community pharmacies and discuss implications and activities for the informal sector in future publications. The need for enhanced activities surrounding the AwaRe system and prescribing guidance in the WHO AwaRe book was noted for all key stakeholder groups to meet UNGA targets.<sup>98</sup>

Following the principles of the narrative review, the main training needs identified have been adapted into a range of potential AwaRe-based educational activities and are summarized in Box 1.

As noted above, the main findings with respect to the KAPs of the different stakeholder groups were remarkably similar across the four WHO Regions (Tables S17–S20). As a result, the suggested activities to improve future antibiotic prescribing and dispensing to meet future national targets for the optimal use of Access antibiotics will also be similar across the four WHO Regions. In view of this, just the suggested activities for government/health authorities will be documented here (Table 2). Suggested activities for the other key stakeholders, including prescribers, dispensers, universities/students and patients, are included in Tables S21–S24.

Similar to high-income settings, even when a higher level of theoretical knowledge was identified, attitudes and practice were markedly different to self-reported knowledge about antibiotics and AMR. This was a very notable finding across all the four WHO Regions and countries of varying income groups. There was also a clear lack of any postgraduate education or training on optimal antibiotic use across all relevant stakeholders, with most studies noting the respondents surveyed were typically dissatisfied with their own knowledge and the resultant implications for routine practice.

Overall, across the major stakeholder groups, there was a clear need to improve the level of basic knowledge on optimal antibiotic use. This was particularly notable for the primary healthcare sector, where no detailed WHO guidance on undertaking ASPs in the primary care setting currently exists. However, this will require financial resources and personnel to help instigate, or update, information technology (IT) and surveillance systems, to routinely collect pertinent healthcare data to

improve future antibiotic use in primary care. Such activities can build on systems such as DHIS2, which has been developed for Africa, Asia and the Middle East,<sup>380</sup> as well as GLASS.<sup>362</sup> Activities to enhance IT systems, including electronic patient records, are essential to undertake ASPs on a routine and real-time basis. This is extremely difficult with paper-based systems, with analysis and reporting typically taking several months after the event. As part of this, governments and health authorities also need continually to assess the cost-effectiveness of any ASPs that are being instigated, including the potential role of artificial intelligence, to improve future antibiotic use in a cost-efficient manner. This is particularly important in LMICs given their limited resources.

There also needs to be continuing research regarding effective and efficient ways to educate patients on antibiotics and AMR given their increasing role in LMICs, while remaining mindful of limited resources. Governments and health authorities also need to work closely with key stakeholder groups to refine the WHO AwaRe book guidance for the local situation, building on activities suggested in Box 1. These include developing and agreeing potential indicators to help progress towards UNGA Access targets.

Governments and health authorities also need to work with key pharmacy organizations to reduce inappropriate dispensing of antibiotics, especially for self-limiting conditions. This again includes instigating systems to better monitor antibiotic dispensing patterns alongside any agreed indicators. Universities also need to ensure that on graduation all healthcare students are fully conversant with the WHO AwaRe classification and guidance, as well as ASPs, given ongoing concerns across all four WHO Regions (Table S19). Consequently, universities need critically to re-evaluate their curricula for the different student HCPs to see if they are fit for purpose for the future. Similarly for any CPD activities.

Finally, governments and health authorities need to engage more with patients and their organizations to address current misconceptions regarding the effectiveness of antibiotics for viral infections as well as the need to complete the full course of any antibiotics prescribed or dispensed. This would reduce the pressure on prescribers and dispensers to provide antibiotics for essentially self-limiting conditions. This includes closer working with patient organizations and others to explore new social media channels to more effectively communicate with patients in languages and with concepts that they understand. Such activities can build on the social media experiences and learnings from the COVID-19 pandemic, including vaccinations, where there were appreciable concerns.<sup>369,381–383</sup> These various activities are explored further in Table 2 as well as in Tables S21–S24, for each key stakeholder group.

We are aware of many limitations to this study. It was not practically possible, given the number of studies identified, to report data across all WHO regions and countries. However, all the major stakeholders involved in the day-to-day prescribing and dispensing of antibiotics were included, as well as students across all subject areas alongside patients and the public. Given the wide variety of methodologies used in the included papers, a formal systematic review was not possible. However, the marked similarity of the main findings across all stakeholders and countries included made the wider generalizability of the key findings more likely.

**Table 2.** Potential interventions for government/health authority activities

Short to medium term (1 to 5 y)	<div data-bbox="321 289 568 315"><b>A) National Action Plans</b></div> <div data-bbox="321 319 1513 751"><ul style="list-style-type: none"><li>• Governments and health authorities across LMICs need to focus on reducing inappropriate use of antibiotics in the primary healthcare setting alongside other measures to reduce AMR in a One Health approach.<sup>361</sup> This will necessarily involve resources (technical/personnel and financial) to address current challenges including concerns with the KAP of all key stakeholder groups translating into high levels of inappropriate prescribing and dispensing of antibiotics.</li><li>• This includes a focus on the integration of routine antibiotic surveillance into electronic healthcare systems such as DHIS2 and others as part of the primary care agenda.<sup>362</sup> Routine surveillance would include the monitoring of prescribing and dispensing patterns against agreed AWARe-based prescribing/quality indicators and quantity metrics. As part of this, governments and health authority personnel in each country may need to refine existing indicators to ensure their relevance and enhance their use.<sup>94</sup></li><li>• Continue to encourage the routine use and standardization of primary care ASPs among prescribers and dispensers to improve future antibiotic use.</li><li>• As part of this, continue to assess the cost-effectiveness of policy and ASP interventions with the aid of academic institutions to help refine future strategies given often limited resources in LMICs.<sup>140,363</sup> This includes the future role of artificial intelligence/machine learning to improve future antibiotic use as part of ASPs,<sup>364–366</sup> and possible virtual reality initiatives.<sup>367</sup></li><li>• Seek to pilot/introduce educational campaigns among patients to help attain NAP goals, cognizant of language and potential gender issues, and monitor their outcomes with the help of academic units as part of NAP activities.<sup>70,166,357,368</sup></li></ul></div> <div data-bbox="321 756 698 781"><b>B) Prescribers (physicians and nurses)</b></div> <div data-bbox="321 785 1513 1218"><ul style="list-style-type: none"><li>• Continue to evaluate the KAP of prescribers given current concerns regarding antibiotics, AMR and ASPs among LMICs in practice, and instigate additional educational activities where necessary through the help of universities as part of CPD activities.</li><li>• Encourage prescriber organizations in each country to work with governments/health authorities to refine local guidelines based on the WHO AWARe book and local resistance patterns.</li><li>• As part of this, key organizations need to work closely with physicians and other prescribers in their country to encourage them to regularly consult national/local guidelines concerning optimal treatment for patients, with guideline adherence increasingly seen as a key quality marker.</li><li>• Where necessary, key organizations should work with governments/health authorities to make guidelines easily accessible to prescribers—increasingly through simple, easy-to-use applications and other systems (Box 1).</li><li>• Such activities should be part of local ASP activities to help achieve UNGA goals. This will require key organizations to work with national governments and health authorities to improve electronic systems within countries to enhance real-time monitoring of prescribing practices.</li><li>• Alongside this, key organizations should work with governments and health authorities to refine and prioritize the introduction of pertinent quality indicators based on the AWARe system.</li></ul></div> <div data-bbox="321 1222 714 1247"><b>C) Dispensers (community pharmacists)</b></div> <div data-bbox="321 1251 1513 1684"><ul style="list-style-type: none"><li>• Continue researching the KAP of dispensers given current concerns regarding antibiotics, AMR and ASPs among LMICs in practice across the four WHO Regions.</li><li>• Seek to instigate additional educational activities where necessary through the help of universities as part of CPD activities. This includes activities to help dispel misinformation, which was problematic during the COVID-19 pandemic.<sup>369–372</sup></li><li>• As part of this, governments/health authorities in each country need to work with key pharmacy organizations to help encourage dispensers to regularly consult national/local guidelines concerning optimal treatment for patients presenting with infectious diseases where there are concerns.</li><li>• Where necessary, governments/health authorities need to work with pharmacy organizations to make guidelines easily accessible to dispensers—increasingly through simple, easy-to-use applications and other systems (Box 1).</li><li>• Governments/health authorities also need to ensure that any list of antibiotics that can be dispensed in community pharmacies is in line with WHO AWARe book recommendations where there are current concerns.<sup>373</sup> This includes any antibiotics that are part of potential quality indicators introduced in community pharmacies to improve future antibiotic dispensing and their monitoring.</li><li>• Alongside this, governments/health authorities need to work with pharmacy organizations to improve monitoring systems in community pharmacies, which can include the use of mobile technologies.<sup>17,374</sup></li></ul></div> <div data-bbox="321 1688 470 1713"><b>D) Universities</b></div> <div data-bbox="321 1717 1513 1894"><ul style="list-style-type: none"><li>• Governments/health authorities need to work with universities to ensure that current HCP curricula meet agreed standards. This increasingly includes the fact that no HCP student should leave university without being fully aware of the WHO AWARe classification and guidance.</li><li>• Governments/health authorities also need to work closely with universities to ensure that CPD activities are aimed at improving dispenser knowledge of national/AWARe book guidelines in order to offer appropriate advice to patients when presenting with self-limiting conditions such as URTIs. Alongside this, instigation of communication skills with patients.<sup>375</sup></li></ul></div>
---------------------------------	--



	<ul style="list-style-type: none"><li>• Governments/health authorities also need to work closely with universities to monitor the KAP of all key stakeholders towards antibiotics, AMR and AMS, as well as research the outcomes of any national initiatives, including guideline introduction and possible quality indicators, as part of academic activities. This also includes the future role of artificial intelligence/machine learning activities to improve antibiotic use as part of ASPs.<sup>364–366</sup></li><li>• Alongside this, use academic institutions to help assess the cost-effectiveness of any targeted educational campaign among patients.</li></ul> <p><b>E) Patients/patient organizations</b></p> <ul style="list-style-type: none"><li>• Work with patient organizations across LMICs to improve existing communication regarding the lack of effectiveness of antibiotics for viral infections/high levels of inappropriate use will increase AMR—building on successful campaigns among LMICs.<sup>68,139–143</sup></li><li>• Patient organizations can also help governments/health authorities to understand key issues surrounding the terminology of antibiotics and AMR. These issues are crucial when health authorities and others are designing educational initiatives for patients to enhance their chance of success.<sup>357–359</sup></li><li>• In addition, work with patient organizations to explore new social media channels to effectively communicate with patients as well as address misinformation, which was problematic during the COVID-19 pandemic.<sup>172,363,376–378</sup></li><li>• Alongside this, also explore virtual reality initiatives to increase awareness of antibiotics and AMR among the public.<sup>367</sup></li><li>• Work with patient organizations to encourage people to become antibiotic guardians where such programmes exist.<sup>379</sup></li></ul>
Longer term	<ul style="list-style-type: none"><li>• Regularly monitor antimicrobial use patterns across all sectors as part of agreed NAPs.</li><li>• Instigate where pertinent additional multiple strategies to improve antibiotic utilization based on the WHO AWaRe book guidance. This includes continuing to improve AMR surveillance as well as enhancing electronic monitoring systems where pertinent.</li><li>• Continue to refine international guidance and quality indicators where pertinent to continue to enhance their acceptance. As a result, ensure any indicators remain current as well as avoiding overloading HCPs across locations.</li><li>• In addition, continue to research the cost-effectiveness of ASPs as well as educational activities among patients to help achieve UNGA goals, and refine where necessary.</li><li>• Continue to work with universities to ensure adequate training of HCPs as well as support for CPD and research activities.</li></ul>

**Box 1. Potential cross-stakeholder AWaRe educational initiatives through the WHO and other international organizations and governments.**

- A) *Initial phase*—building AWaRe-ness focused on optimal antibiotic use and the WHO AWaRe book
- Practical updated guidance on the principles of optimal antibiotic use, including recent major research findings that impact on routine clinical care.
  - Simple short guidance based on the WHO AWaRe book. These could include R-U-AWaRe infographics—simple and easy summaries of the AWaRe book guidance for the main infections seen in ambulatory care translated into local languages where necessary.
  - The Quick-AWaRe book—a short version of the most common infections seen in primary care broken down into summary key points for each clinical area locally adapted to reflect normative standards of care.
  - The Your-AWaRe book localization tool—integration of the WHO AWaRe book with local government/national guidelines with an online tool. This would include a localization process to ensure that the guidance has been appropriately adapted to specific settings.
  - AWaRe-ness training—building a series of webinars/talks/training based on the WHO AWaRe book and system relevant to the stakeholder group.
  - Local and nationally adapted AWaRe quality indicators adapted per WHO Region/Country building on ongoing activities.<sup>94</sup>
- B) *Second phase*
- Development and dissemination of a global community of AWaRe champions and trainers including a toolkit to improve AWaRe-ness through local and national projects/audits/implementation projects.

This review was also conducted during the preparations for UNGA-AMR in 2024. The major commitment of UNGA that 70% of all antibiotics used in humans should be Access antibiotics has added urgency to the key findings of the results. There is a clear need for major development in the training across all stakeholders of optimal antibiotic use. This review provides a strong rationale that although there is a need for adaptation and localization, the core components of the practical training required could be addressed by a standard programme with national and regional modules. This is practical and affordable

with a small number of globally relevant current educational training programmes that could be developed and enhanced, with a particular focus on primary care.

**Conclusions**

The development of the AWaRe system by the WHO, including the practical prescribing guidance of the WHO AWaRe book, has created a clear opportunity for the standardization of education regarding optimal antibiotic use, and focusing on practical

training in primary healthcare in LMIC settings. This narrative review has noted that the unmet training needs to enhance KAPs were very similar across all key stakeholder groups, regions and countries.

Consequently, the next step is to identify all current educational provision focusing on improving antibiotic use, developing a network of current providers including universities and educational partners, and building collaboration with the WHO regarding a new, and sustainably funded, AWaRe antibiotic educational training initiative. Alongside this, other key aspects associated with increasing ABR in LMICs, including the role and influence of the informal sector, should be investigated further as part of a co-ordinated holistic approach. Such activities are essential to meet UNGA targets for the use of Access antibiotics and AMR.

## Funding

This work is partly supported by the ADILA project (Antibiotic Data to Inform Local Action). ADILA is funded by the Wellcome Trust [222051/Z/20/Z], who had no role or responsibility in the study design, data collection, analysis, interpretation of the findings and their implication, or writing of the paper.

## Transparency declarations

The authors have no relevant conflicts of interest to declare.

## Supplementary data

Tables S1 to S24 are available as [Supplementary data](#) at JAC-AMR Online.

## Data availability

Additional data is available from the Corresponding Author upon reasonable request. However, all available information has been referenced.

## References

- Global burden of bacterial antimicrobial resistance in 2019: a systematic analysis. *Lancet* 2022; **399**: 629–55. [https://doi.org/10.1016/S0140-6736\(21\)02724-0](https://doi.org/10.1016/S0140-6736(21)02724-0)
- Dadgostar P. Antimicrobial resistance: implications and costs. *Infect Drug Resist* 2019; **12**: 3903–10. <https://doi.org/10.2147/IDR.S234610>
- Global burden of bacterial antimicrobial resistance 1990–2021: a systematic analysis with forecasts to 2050. *Lancet* 2024; **404**: 1199–226. [https://doi.org/10.1016/S0140-6736\(24\)01867-1](https://doi.org/10.1016/S0140-6736(24)01867-1)
- Poudel AN, Zhu S, Cooper N *et al*. The economic burden of antibiotic resistance: a systematic review and meta-analysis. *PLoS One* 2023; **18**: e0285170. <https://doi.org/10.1371/journal.pone.0285170>
- O'Neill J. Tackling drug-resistant infections globally: final report and recommendations. The Review on Antimicrobial Resistance; 2016. [https://amr-review.org/sites/default/files/160525\\_Final%20paper\\_with%20cover.pdf](https://amr-review.org/sites/default/files/160525_Final%20paper_with%20cover.pdf)
- Gautam A. Antimicrobial resistance: the next probable pandemic. *J Nepal Med Assoc* 2022; **60**: 225–8. <https://doi.org/10.31729/jnma.7174>
- Sulis G, Sayood S, Gandra S. Antimicrobial resistance in low- and middle-income countries: current status and future directions. *Expert Rev Anti Infect Ther* 2022; **20**: 147–60. <https://doi.org/10.1080/14787210.2021.1951705>
- Laxminarayan R, Van Boeckel T, Frost I *et al*. The Lancet Infectious Diseases Commission on Antimicrobial Resistance: 6 years later. *Lancet Infect Dis* 2020; **20**: e51–60. [https://doi.org/10.1016/S1473-3099\(20\)30003-7](https://doi.org/10.1016/S1473-3099(20)30003-7)
- Godman B, Egwuenu A, Haque M *et al*. Strategies to improve antimicrobial utilization with a special focus on developing countries. *Life* 2021; **11**: 528. <https://doi.org/10.3390/life11060528>
- Okeke IN, de Kraker MEA, Van Boeckel TP *et al*. The scope of the antimicrobial resistance challenge. *Lancet* 2024; **403**: 2426–38. [https://doi.org/10.1016/S0140-6736\(24\)00876-6](https://doi.org/10.1016/S0140-6736(24)00876-6)
- Klein EY, Milkowska-Shibata M, Tseng KK *et al*. Assessment of WHO antibiotic consumption and Access targets in 76 countries, 2000–15: an analysis of pharmaceutical sales data. *Lancet Infect Dis* 2021; **21**: 107–15. [https://doi.org/10.1016/S1473-3099\(20\)30332-7](https://doi.org/10.1016/S1473-3099(20)30332-7)
- Sulis G, Sayood S, Katukoori S *et al*. Exposure to World Health Organization's AWaRe antibiotics and isolation of multidrug resistant bacteria: a systematic review and meta-analysis. *Clin Microbiol Infect* 2022; **28**: 1193–202. <https://doi.org/10.1016/j.cmi.2022.03.014>
- Sharland M, Gandra S, Huttner B *et al*. Encouraging AWaRe-ness and discouraging inappropriate antibiotic use—the new 2019 essential medicines list becomes a global antibiotic stewardship tool. *Lancet Infect Dis* 2019; **19**: 1278–80. [https://doi.org/10.1016/S1473-3099\(19\)30532-8](https://doi.org/10.1016/S1473-3099(19)30532-8)
- Sharland M, Zanichelli V, Ombajo LA *et al*. The WHO essential medicines list AWaRe book: from a list to a quality improvement system. *Clin Microbiol Infect* 2022; **28**: 1533–5. <https://doi.org/10.1016/j.cmi.2022.08.009>
- Godman B, Haque M, McKimm J *et al*. Ongoing strategies to improve the management of upper respiratory tract infections and reduce inappropriate antibiotic use particularly among lower and middle-income countries: findings and implications for the future. *Curr Med Res Opin* 2020; **36**: 301–27. <https://doi.org/10.1080/03007995.2019.1700947>
- Torres NF, Chibi B, Kuupiel D *et al*. The use of non-prescribed antibiotics; prevalence estimates in low-and-middle-income countries. A systematic review and meta-analysis. *Arch Public Health* 2021; **79**: 2. <https://doi.org/10.1186/s13690-020-00517-9>
- Sono TM, Yeika E, Cook A *et al*. Current rates of purchasing of antibiotics without a prescription across sub-Saharan Africa; rationale and potential programmes to reduce inappropriate dispensing and resistance. *Expert Rev Anti Infect Ther* 2023; **21**: 1025–55. <https://doi.org/10.1080/14787210.2023.2259106>
- Otaigbe II, Elikwu CJ. Drivers of inappropriate antibiotic use in low- and middle-income countries. *JAC Antimicrob Resist* 2023; **5**: dlad062. <https://doi.org/10.1093/jacamr/dlad062>
- Allwell-Brown G, Hussain-Alkhateeb L, Kitutu FE *et al*. Trends in reported antibiotic use among children under 5 years of age with fever, diarrhoea, or cough with fast or difficult breathing across low-income and middle-income countries in 2005–17: a systematic analysis of 132 national surveys from 73 countries. *Lancet Glob Health* 2020; **8**: e799–807. [https://doi.org/10.1016/S2214-109X\(20\)30079-6](https://doi.org/10.1016/S2214-109X(20)30079-6)
- Rony MKK, Sharmi PD, Alamgir HM. Addressing antimicrobial resistance in low and middle-income countries: overcoming challenges and implementing effective strategies. *Environ Sci Pollut Res Int* 2023; **30**: 101896–902. <https://doi.org/10.1007/s11356-023-29434-4>
- Klein EY, Van Boeckel TP, Martinez EM *et al*. Global increase and geographic convergence in antibiotic consumption between 2000 and 2015. *Proc Natl Acad Sci U S A* 2018; **115**: E3463–e70. <https://doi.org/10.1073/pnas.1717295115>
- WHO. Global action plan on antimicrobial resistance. 2016. <https://iris.who.int/bitstream/handle/10665/254352/sea-cd308.pdf?sequence=1>
- World Bank Group. Pulling together to beat superbugs. Knowledge and implementation gaps in addressing antimicrobial resistance. 2019. <https://openknowledge.worldbank.org/bitstream/handle/10986/32552/>

Pulling-Together-to-Beat-Superbugs-Knowledge-and-Implementation-Gaps-in-Addressing-Antimicrobial-Resistance.pdf?sequence=1&isAllowed=y

- 24 Hofer U. The cost of antimicrobial resistance. *Nat Rev Microbiol* 2019; **17**: 3. <https://doi.org/10.1038/s41579-018-0125-x>
- 25 The World Bank. Final report—drug-resistant infections. A threat to our economic future. March 2017. <http://documents1.worldbank.org/curated/en/323311493396993758/pdf/final-report.pdf>
- 26 WHO implementation handbook for national action plans on antimicrobial resistance: guidance for the human health sector. 2022. <https://www.who.int/publications/i/item/9789240041981>
- 27 Moja L, Zanichelli V, Mertz D et al. WHO's essential medicines and AWaRe: recommendations on first- and second-choice antibiotics for empiric treatment of clinical infections. *Clin Microbiol Infect* 2024; **30** Suppl 2: S1–S11. <https://doi.org/10.1016/j.cmi.2024.02.003>
- 28 Zanichelli V, Sharland M, Cappello B et al. The WHO AWaRe (Access, Watch, Reserve) antibiotic book and prevention of antimicrobial resistance. *Bull World Health Organ* 2023; **101**: 290–6. <https://doi.org/10.2471/BLT.22.288614>
- 29 Godman B, Egwuenu A, Wesangula E et al. Tackling antimicrobial resistance across sub-Saharan Africa: current challenges and implications for the future. *Expert Opin Drug Saf* 2022; **21**: 1089–111. <https://doi.org/10.1080/14740338.2022.2106368>
- 30 Charani E, Mendelson M, Pallett SJC et al. An analysis of existing national action plans for antimicrobial resistance—gaps and opportunities in strategies optimising antibiotic use in human populations. *Lancet Glob Health* 2023; **11**: e466–74. [https://doi.org/10.1016/S2214-109X\(23\)00019-0](https://doi.org/10.1016/S2214-109X(23)00019-0)
- 31 Iwu CD, Patrick SM. An insight into the implementation of the global action plan on antimicrobial resistance in the WHO African region: a roadmap for action. *Int J Antimicrob Agents* 2021; **58**: 106411. <https://doi.org/10.1016/j.ijantimicag.2021.106411>
- 32 Chua AQ, Verma M, Hsu LY et al. An analysis of national action plans on antimicrobial resistance in Southeast Asia using a governance framework approach. *Lancet Reg Health West Pac* 2021; **7**: 100084. <https://doi.org/10.1016/j.lanwpc.2020.100084>
- 33 Willemsen A, Reid S, Assefa Y. A review of national action plans on antimicrobial resistance: strengths and weaknesses. *Antimicrob Resist Infect Control* 2022; **11**: 90. <https://doi.org/10.1186/s13756-022-01130-x>
- 34 Vliegenthart-Jongbloed K, Jacobs J. Not recommended fixed-dose antibiotic combinations in low- and middle-income countries—the example of Tanzania. *Antimicrob Resist Infect Control* 2023; **12**: 37. <https://doi.org/10.1186/s13756-023-01238-8>
- 35 Valia D, Ingelbeen B, Nassa GJW et al. Antibiotic use by clinical presentation across all healthcare providers in rural Burkina Faso: a healthcare visit exit survey. *J Antimicrob Chemother* 2024; **79**: 2534–42. <https://doi.org/10.1093/jac/dkac252>
- 36 Mustafa Z U, Batool A, Ibrar H et al. Bacterial co-infections, secondary infections and antimicrobial use among hospitalized COVID-19 patients in the sixth wave in Pakistan: findings and implications. *Expert Rev Anti Infect Ther* 2024; **22**: 229–40. <https://doi.org/10.1080/14787210.2023.2299387>
- 37 Duffy E, Ritchie S, Metcalfe S et al. Antibacterials dispensed in the community comprise 85%–95% of total human antibacterial consumption. *J Clin Pharm Ther* 2018; **43**: 59–64. <https://doi.org/10.1111/jcpt.12610>
- 38 Nepal G, Bhatta S. Self-medication with antibiotics in WHO Southeast Asian region: a systematic review. *Cureus* 2018; **10**: e2428. <https://doi.org/10.7759/cureus.2428>
- 39 Rose J, Crosbie M, Stewart A. A qualitative literature review exploring the drivers influencing antibiotic over-prescribing by GPs in primary care and recommendations to reduce unnecessary prescribing. *Perspect Public Health* 2021; **141**: 19–27. <https://doi.org/10.1177/1757913919879183>
- 40 Guo S, Sun Q, Zhao X et al. Prevalence and risk factors for antibiotic utilization in Chinese children. *BMC Pediatr* 2021; **21**: 255. <https://doi.org/10.1186/s12887-021-02706-z>
- 41 Teague E, Bezuidenhout S, Meyer JC et al. Knowledge and perceptions of final-year nursing students regarding antimicrobials, antimicrobial resistance, and antimicrobial stewardship in South Africa: findings and implications to reduce resistance. *Antibiotics* 2023; **12**: 1742. <https://doi.org/10.3390/antibiotics12121742>
- 42 Lagarde M, Blaauw D. Levels and determinants of overprescribing of antibiotics in the public and private primary care sectors in South Africa. *BMJ Glob Health* 2023; **8**: 1–10. <https://doi.org/10.1136/bmjgh-2023-012374>
- 43 Olaru ID, Ferrand RA, Yeung S et al. Knowledge, attitudes and practices relating to antibiotic use and resistance among prescribers from public primary healthcare facilities in Harare, Zimbabwe. *Wellcome Open Res* 2021; **6**: 72. <https://doi.org/10.12688/wellcomeopenres.16657.1>
- 44 Mabilika RJ, Shirima G, Mpolya E. Prevalence and predictors of antibiotic prescriptions at primary healthcare facilities in the Dodoma region, central Tanzania: a retrospective, cross-sectional study. *Antibiotics* 2022; **11**: 1035. <https://doi.org/10.3390/antibiotics11081035>
- 45 Mathibe LJ, Zwane NP. Unnecessary antimicrobial prescribing for upper respiratory tract infections in children in Pietermaritzburg, South Africa. *Afr Health Sci* 2020; **20**: 1133–42. <https://doi.org/10.4314/ahs.v20i3.15>
- 46 Nair M, Tripathi S, Mazumdar S et al. Knowledge, attitudes, and practices related to antibiotic use in Paschim Bardhaman district: a survey of healthcare providers in West Bengal, India. *PLoS One* 2019; **14**: e0217818. <https://doi.org/10.1371/journal.pone.0217818>
- 47 Adegbite BR, Edoa JR, Schaumburg F et al. Knowledge and perception on antimicrobial resistance and antibiotics prescribing attitude among physicians and nurses in Lambaréné region, Gabon: a call for setting-up an antimicrobial stewardship program. *Antimicrob Resist Infect Control* 2022; **11**: 44. <https://doi.org/10.1186/s13756-022-01079-x>
- 48 Li J, Zhou P, Wang J et al. Worldwide dispensing of non-prescription antibiotics in community pharmacies and associated factors: a mixed-methods systematic review. *Lancet Infect Dis* 2023; **23**: e361–70. [https://doi.org/10.1016/S1473-3099\(23\)00130-5](https://doi.org/10.1016/S1473-3099(23)00130-5)
- 49 Torres NF, Chibi B, Middleton LE et al. Evidence of factors influencing self-medication with antibiotics in low and middle-income countries: a systematic scoping review. *Public Health* 2019; **168**: 92–101. <https://doi.org/10.1016/j.puhe.2018.11.018>
- 50 Abdulrahman Jairoun A, Saleh Al-Hemyari S, Jairoun M. Antibiotics dispensing between legal idealism and implementation: closing the implementation gap between local regulation and practice. *Res Social Adm Pharm* 2020; **16**: 1328–9. <https://doi.org/10.1016/j.sapharm.2020.05.001>
- 51 Makeri D, Dilli PP, Pius T et al. The nature of self-medication in Uganda: a systematic review and meta-analysis. *BMC Public Health* 2025; **25**: 197. <https://doi.org/10.1186/s12889-025-21380-9>
- 52 Waseem H, Ali J, Sarwar F et al. Assessment of knowledge and attitude trends towards antimicrobial resistance (AMR) among the community members, pharmacists/pharmacy owners and physicians in district Sialkot, Pakistan. *Antimicrob Resist Infect Control* 2019; **8**: 67. <https://doi.org/10.1186/s13756-019-0517-3>
- 53 Kumar KS, Saranya S, Rani NV. Community pharmacists' knowledge, attitude, and nonprescription dispensing practices of antibiotics: an explorative study in a selected city of south India. *J Res Pharm Pract* 2022; **11**: 51–8. [https://doi.org/10.4103/jrpp.jrpp\\_48\\_21](https://doi.org/10.4103/jrpp.jrpp_48_21)
- 54 Jairoun AA, Al-Hemyari SS, Shahwan M et al. Top unresolved ethical challenges and dilemmas faced by community pharmacists in providing pharmaceutical care: drawing the line between ethical challenges and the quality of the pharmaceutical care. *Res Social Adm Pharm* 2022; **18**: 3711–3. <https://doi.org/10.1016/j.sapharm.2022.05.009>

- 55 Gebresillassie BM, Howells K, Ashiru-Oredope D. Public health interventions delivered by pharmacy professionals in low- and middle-income countries in Africa: a systematic scoping review. *Pharmacy* 2023; **11**: 24. <https://doi.org/10.3390/pharmacy11010024>.
- 56 Cadogan CA, Hughes CM. On the frontline against COVID-19: community pharmacists' contribution during a public health crisis. *Res Social Adm Pharm* 2021; **17**: 2032–5. <https://doi.org/10.1016/j.sapharm.2020.03.015>
- 57 Kibuule D, Nambahu L, Sefah IA et al. Activities in Namibia to limit the prevalence and mortality from COVID-19 including community pharmacy activities and the implications. *Scholars Academic Journal of Pharmacy* 2021; **10**: 82–92. <https://doi.org/10.36347/sajp.2021.v10i05.001>
- 58 Finnis A. Can pharmacists prescribe antibiotics? The new plans for prescriptions in England, explained. THEIPAPER, 2023. <https://inews.co.uk/news/health/pharmacists-can-prescribe-antibiotics-new-plans-pharmacy-prescriptions-englan-explained-2330773>
- 59 Yevutsey SK, Buabeng KO, Aikins M et al. Situational analysis of antibiotic use and resistance in Ghana: policy and regulation. *BMC Public Health* 2017; **17**: 896. <https://doi.org/10.1186/s12889-017-4910-7>
- 60 Bharti RK, Pathania JS, Sood V et al. Assessing the knowledge, attitude, and practice (KAP) of antimicrobial resistant among MBBS, BDS and BSc nursing students in the northern state of India. An observational-based cross-sectional study. *Adv Biosci Clin Med* 2020; **8**: 10–6. <https://doi.org/10.7575/aic.abcm.v.8n.2p.10>
- 61 Nogueira-Uzal N, Zapata-Cachafeiro M, Vázquez-Cancela O et al. Does the problem begin at the beginning? Medical students' knowledge and beliefs regarding antibiotics and resistance: a systematic review. *Antimicrob Resist Infect Control* 2020; **9**: 172. <https://doi.org/10.1186/s13756-020-00837-z>
- 62 Bulubula ANH, Jenkins A, Mehtar S et al. Education and management of antimicrobials amongst nurses in Africa—a situation analysis: an Infection Control Africa Network (ICAN)/BSAC online survey. *J Antimicrob Chemother* 2018; **73**: 1408–15. <https://doi.org/10.1093/jac/dky023>
- 63 Ashiru-Oredope D, Casale E, Harvey E et al. Knowledge and attitudes about antibiotics and antibiotic resistance of 2404 UK healthcare workers. *Antibiotics* 2022; **11**: 1133. <https://doi.org/10.3390/antibiotics11081133>
- 64 Khan FU, Khan FU, Hayat K et al. Knowledge, attitude and practices among consumers toward antibiotics use and antibiotic resistance in Swat, Khyber-Pakhtunkhwa, Pakistan. *Expert Rev Anti Infect Ther* 2020; **18**: 937–46. <https://doi.org/10.1080/14787210.2020.1769477>
- 65 Antwi AN, Stewart A, Crosbie M. Fighting antibiotic resistance: a narrative review of public knowledge, attitudes, and perceptions of antibiotics use. *Perspect Public Health* 2020; **140**: 338–50. <https://doi.org/10.1177/1757913920921209>
- 66 Mitchell J, Cooke P, Ahorlu C et al. Community engagement: the key to tackling antimicrobial resistance (AMR) across a one health context? *Glob Public Health* 2022; **17**: 2647–64. <https://doi.org/10.1080/17441692.2021.2003839>
- 67 Ramdas N, Meyer JC, Schellack N et al. Knowledge, attitudes, motivations, expectations, and systemic factors regarding antimicrobial use amongst community members seeking care at the primary healthcare level: a scoping review. *Antibiotics* 2025; **14**: 78. <https://doi.org/10.3390/antibiotics14010078>
- 68 Kandeel A, Palms DL, Afifi S et al. An educational intervention to promote appropriate antibiotic use for acute respiratory infections in a district in Egypt—pilot study. *BMC Public Health* 2019; **19** Suppl 3: 498. <https://doi.org/10.1186/s12889-019-6779-0>
- 69 Nayiga S, MacPherson EE, Mankhomwa J et al. “Arming half-baked people with weapons!” information enclaving among professionals and the need for a care-centred model for antibiotic use information in Uganda, Tanzania and Malawi. *Glob Health Action* 2024; **17**: 2322839. <https://doi.org/10.1080/16549716.2024.2322839>
- 70 Haenssge MJ, Charoenboon N, Zanello G et al. Antibiotic knowledge, attitudes and practices: new insights from cross-sectional rural health behaviour surveys in low-income and middle-income South-East Asia. *BMJ Open* 2019; **9**: e028224. <https://doi.org/10.1136/bmjopen-2018-028224>
- 71 Charoenboon N, Haenssge MJ, Warapikuptanun P et al. Translating antimicrobial resistance: a case study of context and consequences of antibiotic-related communication in three northern Thai villages. *Palgrave Commun* 2019; **5**: 23. <https://doi.org/10.1057/s41599-019-0226-9>
- 72 Sono TM, Mboweni V, Jelić AG et al. Pilot study to evaluate patients' understanding of key terms and aspects of antimicrobial use in a rural province in South Africa: findings and implications. *Adv Hum Biol* 2025; **15**: 108–12. [https://doi.org/10.4103/aihb.aihb.119\\_24](https://doi.org/10.4103/aihb.aihb.119_24)
- 73 Nathwani D, Varghese D, Stephens J et al. Value of hospital antimicrobial stewardship programs (ASPs): a systematic review. *Antimicrob Resist Infect Control* 2019; **8**: 35. <https://doi.org/10.1186/s13756-019-0471-0>
- 74 Lee CF, Cowling BJ, Feng S et al. Impact of antibiotic stewardship programmes in Asia: a systematic review and meta-analysis. *J Antimicrob Chemother* 2018; **73**: 844–51. <https://doi.org/10.1093/jac/dkx492>
- 75 Antimicrobial stewardship programmes in health-care facilities in low- and middle-income countries: a WHO practical toolkit. *JAC Antimicrob Resist* 2019; **1**: dlz072. <https://doi.org/10.1093/jacamr/dlz072>
- 76 Gulumbe BH, Haruna UA, Almazan J et al. Combating the menace of antimicrobial resistance in Africa: a review on stewardship, surveillance and diagnostic strategies. *Biol Proced Online* 2022; **24**: 19. <https://doi.org/10.1186/s12575-022-00182-y>
- 77 Harun MGD, Sumon SA, Hasan I et al. Barriers, facilitators, perceptions and impact of interventions in implementing antimicrobial stewardship programs in hospitals of low-middle and middle countries: a scoping review. *Antimicrob Resist Infect Control* 2024; **13**: 8. <https://doi.org/10.1186/s13756-024-01369-6>
- 78 Ashiru-Oredope D, Nabiryo M, Zengeni L et al. Tackling antimicrobial resistance: developing and implementing antimicrobial stewardship interventions in four African commonwealth countries through a health partnership model. *J Public Health Afr* 2023; **14**: 2335. <https://doi.org/10.4081/jphia.2023.2335>
- 79 Sneddon J, Drummond F, Guise T et al. Accreditation of antimicrobial stewardship programmes: addressing a global need to tackle antimicrobial resistance. *JAC Antimicrob Resist* 2024; **6**: dlac007. <https://doi.org/10.1093/jacamr/dlae007>
- 80 The Fleming Fund. Fleming Fund Online AMR Course: new modules! 2021. <https://www.flemingfund.org/publications/fleming-fund-online-amr-course/>
- 81 Chigome A, Ramdas N, Skosana P et al. A narrative review of antibiotic prescribing practices in primary care settings in South Africa and potential ways forward to reduce antimicrobial resistance. *Antibiotics* 2023; **12**: 1540. <https://doi.org/10.3390/antibiotics12101540>
- 82 Masele A, Rogers AM, Gabriel D et al. A narrative review of recent antibiotic prescribing practices in ambulatory care in Tanzania: findings and implications. *Medicina* 2023; **59**: 2195. <https://doi.org/10.3390/medicina59122195>
- 83 Nair MM, Mahajan R, Burza S et al. Behavioural interventions to address rational use of antibiotics in outpatient settings of low-income and lower-middle-income countries. *Trop Med Int Health* 2021; **26**: 504–17. <https://doi.org/10.1111/tmi.13550>
- 84 WHO. Antimicrobial stewardship interventions: a practical guide. <https://iris.who.int/bitstream/handle/10665/340709/9789289054980-eng.pdf>
- 85 Open University. Understanding antibiotic resistance. <https://www.open.edu/openlearn/science-maths-technology/understanding-antibiotic-resistance/content-section-overview?active-tab=description-tab>



- 86** Ogunleye OO, Fadare JO, Eriksen J *et al.* Reported needs of information resources, research tools, connectivity and infrastructure among African pharmacological scientists to improve future patient care and health. *Expert Rev Clin Pharmacol* 2019; **12**: 481–9. <https://doi.org/10.1080/17512433.2019.1605903>
- 87** Infection Control Africa Network. Training—ICAN training courses. <https://icanetwork.co.za/training/>
- 88** Kalungia AC, Kampamba M, Banda D *et al.* Impact of a hub-and-spoke approach to hospital antimicrobial stewardship programmes on antibiotic use in Zambia. *JAC Antimicrob Resist* 2024; **6**: dlae178. <https://doi.org/10.1093/jacamr/dlae178>
- 89** Niaz Q, Godman B, Massele A *et al.* Validity of World Health Organization prescribing indicators in Namibia's primary healthcare: findings and implications. *Int J Qual Health Care* 2019; **31**: 338–45. <https://doi.org/10.1093/intqhc/mzy172>
- 90** Campbell SM, Meyer J, Godman B. Why compliance to national prescribing guidelines is important especially across sub-Saharan Africa and suggestions for the future. *Biomed Pharm Sci* 2021; **4**: 1–7.
- 91** Saleem Z, Godman B, Cook A *et al.* Ongoing efforts to improve antimicrobial utilization in hospitals among African countries and implications for the future. *Antibiotics* 2022; **11**: 1824. <https://doi.org/10.3390/antibiotics11121824>
- 92** Boltana MT, Woldie M, Siraneh Y *et al.* Adherence to evidence-based implementation of antimicrobial treatment guidelines among prescribers in sub-Saharan Africa: a systematic review and meta-analysis. *J Pharm Policy Pract* 2023; **16**: 137. <https://doi.org/10.1186/s40545-023-00634-0>
- 93** Kirchhelle C, Atkinson P, Broom A *et al.* Setting the standard: multidisciplinary hallmarks for structural, equitable and tracked antibiotic policy. *BMJ Glob Health* 2020; **5**: e003091. <https://doi.org/10.1136/bmjgh-2020-003091>
- 94** Funicello E, Lorenzetti G, Cook A *et al.* Identifying AWARe indicators for appropriate antibiotic use: a narrative review. *J Antimicrob Chemother* 2024; **79**: 3063–77. <https://doi.org/10.1093/jac/dkae370>
- 95** Kredt T, Cooper S, Abrams AL *et al.* 'Building on shaky ground'—challenges to and solutions for primary care guideline implementation in four provinces in South Africa: a qualitative study. *BMJ Open* 2020; **10**: e031468. <https://doi.org/10.1136/bmjopen-2019-031468>
- 96** Atalay YA, Abebe Gelaw K. Prevalence of knowledge, attitudes, and practices regarding antimicrobial resistance in Africa: a systematic review and meta-analysis. *Front Microbiol* 2024; **15**: 1345145. <https://doi.org/10.3389/fmicb.2024.1345145>
- 97** Fuller W, Kapona O, Aboderin AO *et al.* Education and awareness on antimicrobial resistance in the WHO African region: a systematic review. *Antibiotics* 2023; **12**: 1613. <https://doi.org/10.3390/antibiotics12111613>
- 98** United Nations. Political declaration of the high-level meeting on antimicrobial resistance. 9 September 2024. <https://www.un.org/pga/wp-content/uploads/sites/108/2024/09/FINAL-Text-AMR-to-PGA.pdf>
- 99** WHO. Antibiotic resistance: multi-country public awareness survey. 2015. [https://iris.who.int/bitstream/handle/10665/194460/9789241509817\\_eng.pdf?sequence=1&isAllowed=y](https://iris.who.int/bitstream/handle/10665/194460/9789241509817_eng.pdf?sequence=1&isAllowed=y)
- 100** Kretchy J-P, Adase SK, Gyansa-Lutterodt M. The prevalence and risks of antibiotic self-medication in residents of a rural community in Accra, Ghana. *Sci Afr* 2021; **14**: e01006. <https://doi.org/10.1016/j.sciaf.2021.e01006>
- 101** van de Maat J, De Santis O, Luwanda L *et al.* Primary care case management of febrile children: insights from the ePOCT routine care cohort in Dar es Salaam, Tanzania. *Front Pediatr* 2021; **9**: 626386. <https://doi.org/10.3389/fped.2021.626386>
- 102** Sefah IA, Essah DO, Kurdi A *et al.* Assessment of adherence to pneumonia guidelines and its determinants in an ambulatory care clinic in Ghana: findings and implications for the future. *JAC Antimicrob Resist* 2021; **3**: dlab080. <https://doi.org/10.1093/jacamr/dlab080>
- 103** Niaz Q, Godman B, Campbell S *et al.* Compliance to prescribing guidelines among public health care facilities in Namibia; findings and implications. *Int J Clin Pharm* 2020; **42**: 1227–36. <https://doi.org/10.1007/s11096-020-01056-7>
- 104** Hadley MB, Beard J. Is 'health for all' synonymous with 'antibiotics for all': changes in antibiotic prescribing in a performance-based financing pilot in Zanzibar. *Health Policy Plan* 2019; **34** Suppl 2: ii28–35. <https://doi.org/10.1093/heapol/czz103>
- 105** The burden of bacterial antimicrobial resistance in the WHO African region in 2019: a cross-country systematic analysis. *Lancet Glob Health* 2024; **12**: e201–16. [https://doi.org/10.1016/S2214-109X\(23\)00539-9](https://doi.org/10.1016/S2214-109X(23)00539-9)
- 106** Chaw PS, Höpner J, Mikolajczyk R. The knowledge, attitude and practice of health practitioners towards antibiotic prescribing and resistance in developing countries—a systematic review. *J Clin Pharm Ther* 2018; **43**: 606–13. <https://doi.org/10.1111/jcpt.12730>
- 107** Pham-Duc P, Sriparamanathan K. Exploring gender differences in knowledge and practices related to antibiotic use in Southeast Asia: a scoping review. *PLoS One* 2021; **16**: e0259069. <https://doi.org/10.1371/journal.pone.0259069>
- 108** Xu R, Mu T, Wang G *et al.* Self-medication with antibiotics among university students in LMIC: a systematic review and meta-analysis. *J Infect Dev Ctries* 2019; **13**: 678–89. <https://doi.org/10.3855/jidc.11359>
- 109** Mwita JC, Ogunleye OO, Olalekan A *et al.* Key issues surrounding appropriate antibiotic use for prevention of surgical site infections in low- and middle-income countries: a narrative review and the implications. *Int J Gen Med* 2021; **14**: 515–30. <https://doi.org/10.2147/IJGM.S253216>
- 110** Haseeb A, Saleem Z, Maqadmi AF *et al.* Ongoing strategies to improve antimicrobial utilization in hospitals across the Middle East and North Africa (MENA): findings and implications. *Antibiotics* 2023; **12**: 827. <https://doi.org/10.3390/antibiotics12050827>
- 111** WHO. Regional offices. <https://www.who.int/about/who-we-are/regional-offices>
- 112** Adekoya I, Maraj D, Steiner L *et al.* Comparison of antibiotics included in national essential medicines lists of 138 countries using the WHO access, watch, reserve (AWARe) classification: a cross-sectional study. *Lancet Infect Dis* 2021; **21**: 1429–40. [https://doi.org/10.1016/S1473-3099\(20\)30854-9](https://doi.org/10.1016/S1473-3099(20)30854-9)
- 113** Joshi MP, Hafner T, Twesigye G *et al.* Strengthening multisectoral coordination on antimicrobial resistance: a landscape analysis of efforts in 11 countries. *J Pharm Policy Pract* 2021; **14**: 27. <https://doi.org/10.1186/s40545-021-00309-8>
- 114** Etando A, Amu AA, Haque M *et al.* Challenges and innovations brought about by the COVID-19 pandemic regarding medical and pharmacy education especially in Africa and implications for the future. *Healthcare* 2021; **9**: 1722. <https://doi.org/10.3390/healthcare9121722>
- 115** Chowdhury K, Haque M, Etando A *et al.* The global impact of the COVID-19 pandemic on the education of healthcare professionals, especially in low- and middle-income countries. *Adv Hum Biol* 2022; **12**: 87–92. [https://doi.org/10.4103/aihb.aihb\\_60\\_22](https://doi.org/10.4103/aihb.aihb_60_22)
- 116** Chinna K, Sundarasan S, Khoshaim HB *et al.* Psychological impact of COVID-19 and lock down measures: an online cross-sectional multi-county study on Asian university students. *PLoS One* 2021; **16**: e0253059. <https://doi.org/10.1371/journal.pone.0253059>
- 117** Bednarčuk N, Golić Jelić A, Stoisavljević Štara S *et al.* Antibiotic utilization during COVID-19: are we over-prescribing? *Antibiotics* 2023; **12**: 308. <https://doi.org/10.3390/antibiotics12020308>
- 118** Sulis G, Batomen B, Kotwani A *et al.* Sales of antibiotics and hydroxy-chloroquine in India during the COVID-19 epidemic: an interrupted time series analysis. *PLoS Med* 2021; **18**: e1003682. <https://doi.org/10.1371/journal.pmed.1003682>

- 119** Gul B, Sana M, Saleem A *et al.* Antimicrobial dispensing practices during COVID-19 and the implications for Pakistan. *Antibiotics* 2023; **12**: 1018. <https://doi.org/10.3390/antibiotics12061018>
- 120** Nkengasong JN, Tessema SK. Africa needs a new public health order to tackle infectious disease threats. *Cell* 2020; **183**: 296–300. <https://doi.org/10.1016/j.cell.2020.09.041>
- 121** Bell D, Schultz Hansen K. Relative burdens of the COVID-19, malaria, tuberculosis, and HIV/AIDS epidemics in sub-Saharan Africa. *Am J Trop Med Hyg* 2021; **105**: 1510–5. <https://doi.org/10.4269/ajtmh.21-0899>
- 122** Williams PCM, Isaacs D, Berkley JA. Antimicrobial resistance among children in sub-Saharan Africa. *Lancet Infect Dis* 2018; **18**: e33–44. [https://doi.org/10.1016/S1473-3099\(17\)30467-X](https://doi.org/10.1016/S1473-3099(17)30467-X)
- 123** Ding G, Vinturache A, Lu M. Addressing inappropriate antibiotic prescribing in China. *Can Med Assoc J* 2019; **191**: E149–e50. <https://doi.org/10.1503/cmaj.181417>
- 124** Luo Y, Tang X, Ding L *et al.* Non-prescription antibiotic use for people aged 15 years or older for cough in China: a community-based survey. *Antimicrob Resist Infect Control* 2021; **10**: 129. <https://doi.org/10.1186/s13756-021-00998-5>
- 125** Xu J, Wang X, Sun KS *et al.* Parental self-medication with antibiotics for children promotes antibiotic over-prescribing in clinical settings in China. *Antimicrob Resist Infect Control* 2020; **9**: 150. <https://doi.org/10.1186/s13756-020-00811-9>
- 126** Zhang T, Lambert H, Zhao L *et al.* Antibiotic stewardship in retail pharmacies and the access-excess challenge in China: a policy review. *Antibiotics* 2022; **11**: 141. <https://doi.org/10.3390/antibiotics11020141>
- 127** Qu W, Wang X, Liu Y *et al.* Self-medication with antibiotics among children in China: a cross-sectional study of parents' knowledge, attitudes, and practices. *Infect Drug Resist* 2023; **16**: 7683–94. <https://doi.org/10.2147/IDR.S431034>
- 128** Li C, Cui Z, Wei D *et al.* Trends and patterns of antibiotic prescriptions in primary care institutions in southwest China, 2017–2022. *Infect Drug Resist* 2023; **16**: 5833–54. <https://doi.org/10.2147/IDR.S425787>
- 129** Xu X, Zhang K, Ma H *et al.* Differences in service and antibiotics use following symptomatic respiratory tract infections between 2016 and 2021 in rural Anhui, China. *Epidemiol Infect* 2022; **150**: e117. <https://doi.org/10.1017/S0950268822000942>
- 130** Zhang T, Lin H, Zhao X *et al.* Influences on treatment-seeking and antibiotic use for common illnesses in eastern China. *BMC Public Health* 2023; **23**: 1849. <https://doi.org/10.1186/s12889-023-16700-w>
- 131** Chan OSK, Lam W, Zhao S *et al.* Why prescribe antibiotics? A systematic review of knowledge, tension, and motivation among clinicians in low-, middle- and high-income countries. *Soc Sci Med* 2024; **345**: 116600. <https://doi.org/10.1016/j.socscimed.2024.116600>
- 132** Sulis G, Adam P, Nafade V *et al.* Antibiotic prescription practices in primary care in low- and middle-income countries: a systematic review and meta-analysis. *PLoS Med* 2020; **17**: e1003139. <https://doi.org/10.1371/journal.pmed.1003139>
- 133** Graells T, Lambraki IA, Cousins M *et al.* Exploring the factors that contribute to the successful implementation of antimicrobial resistance interventions: a comparison of high-income and low-middle-income countries. *Front Public Health* 2023; **11**: 1230848. <https://doi.org/10.3389/fpubh.2023.1230848>
- 134** Villanueva P, Coffin SE, Mekasha A *et al.* Comparison of antimicrobial stewardship and infection prevention and control activities and resources between low-/middle- and high-income countries. *Pediatr Infect Dis J* 2022; **41**: S3–S9. <https://doi.org/10.1097/INF.0000000000003318>
- 135** Naing S, van Wijk M, Vila J *et al.* Understanding antimicrobial resistance from the perspective of public policy: a multinational knowledge, attitude, and perception survey to determine global awareness. *Antibiotics* 2021; **10**: 1486. <https://doi.org/10.3390/antibiotics10121486>
- 136** Taylor W, Whittaker LM, Fletcher T *et al.* Awareness of inappropriate use related to antimicrobial resistance among medical doctors by country economic status: a systematic review. *Int J Risk Saf Med* 2023; **34**: 227–42. <https://doi.org/10.3233/JRS-220044>
- 137** McDonagh MS, Peterson K, Winthrop K *et al.* Interventions to reduce inappropriate prescribing of antibiotics for acute respiratory tract infections: summary and update of a systematic review. *J Int Med Res* 2018; **46**: 3337–57. <https://doi.org/10.1177/0300060518782519>
- 138** Cross EL, Tolfree R, Kipping R. Systematic review of public-targeted communication interventions to improve antibiotic use. *J Antimicrob Chemother* 2017; **72**: 975–87. <https://doi.org/10.1093/jac/dkw520>
- 139** Maarouf L, Amin M, Evans BA *et al.* Knowledge, attitudes and behaviour of Egyptians towards antibiotic use in the community: can we do better? *Antimicrob Resist Infect Control* 2023; **12**: 50. <https://doi.org/10.1186/s13756-023-01249-5>
- 140** Aika IN, Enato E. Bridging the gap in knowledge and use of antibiotics among pediatric caregivers: comparing two educational interventions. *J Pharm Policy Pract* 2023; **16**: 76. <https://doi.org/10.1186/s40545-023-00578-5>
- 141** Tamhankar AJ, Nachimuthu R, Singh R *et al.* Characteristics of a nationwide voluntary antibiotic resistance awareness campaign in India: future paths and pointers for resource limited settings/low and middle income countries. *Int J Environ Res Public Health* 2019; **16**: 5141. <https://doi.org/10.3390/ijerph16245141>
- 142** Herawati F, Yulia R, Arifin B *et al.* Educational video improves knowledge about outpatients' usage of antibiotics in two public hospitals in Indonesia. *Antibiotics* 2021; **10**: 606. <https://doi.org/10.3390/antibiotics10050606>
- 143** Thong KS, Chang CT, Lee M *et al.* Impact of targeted educational intervention towards public knowledge and perception of antibiotic use and resistance in the state of Perak, Malaysia. *Antimicrob Resist Infect Control* 2021; **10**: 29. <https://doi.org/10.1186/s13756-021-00892-0>
- 144** Abdu N, Idrisnur S, Tewelde T *et al.* Antibiotic prescribing practice using WHO access, watch and reserve classification and its determinants among outpatient prescriptions dispensed to elderly population in six community chain pharmacies in Asmara, Eritrea: a cross-sectional study. *BMJ Open* 2024; **14**: e085743. <https://doi.org/10.1136/bmjopen-2024-085743>
- 145** El-Sokkary R, Kishk R, Mohy El-Din S *et al.* Antibiotic use and resistance among prescribers: current status of knowledge, attitude, and practice in Egypt. *Infect Drug Resist* 2021; **14**: 1209–18. <https://doi.org/10.2147/IDR.S299453>
- 146** Amin MT, Abd El Aty MA, Ahmed SM *et al.* Over prescription of antibiotics in children with acute upper respiratory tract infections: a study on the knowledge, attitude and practices of non-specialized physicians in Egypt. *PLoS One* 2022; **17**: e0277308. <https://doi.org/10.1371/journal.pone.0277308>
- 147** Karasneh RA, Al-Azzam SI, Ababneh M *et al.* Prescribers' knowledge, attitudes and behaviors on antibiotics, antibiotic use and antibiotic resistance in Jordan. *Antibiotics* 2021; **10**: 858. <https://doi.org/10.3390/antibiotics10070858>
- 148** Hashmi H, Sasoli NA, Sadiq A *et al.* Prescribing patterns for upper respiratory tract infections: a prescription-review of primary care practice in Quetta, Pakistan and the implications. *Front Public Health* 2021; **9**: 787933. <https://doi.org/10.3389/fpubh.2021.787933>
- 149** Orubu ESF, Al-Dheeb N, Ching C *et al.* Assessing antimicrobial resistance, utilization, and stewardship in Yemen: an exploratory mixed-methods study. *Am J Trop Med Hyg* 2021; **105**: 1404–12. <https://doi.org/10.4269/ajtmh.21-0101>

- 150** Ingelbeen B, Phanhu DM, Phoba MF et al. Antibiotic use from formal and informal healthcare providers in the Democratic Republic of Congo: a population-based study in two health zones. *Clin Microbiol Infect* 2022; **28**: 1272–7. <https://doi.org/10.1016/j.cmi.2022.04.002>
- 151** Kakumba JM, Kindenge JM, Kapepula PM et al. Evaluation of antibiotic prescribing pattern using WHO access, watch and reserve classification in Kinshasa, Democratic Republic of Congo. *Antibiotics* 2023; **12**: 39. <https://doi.org/10.3390/antibiotics12081239>
- 152** Khalfan MA, Sasi P, Mugusi S. Factors influencing receipt of an antibiotic prescription among insured patients in Tanzania: a cross-sectional study. *BMJ Open* 2022; **12**: e062147. <https://doi.org/10.1136/bmjopen-2022-062147>
- 153** Dereje B, Workneh A, Megersa A et al. Prescribing pattern and associated factors in community pharmacies: a cross-sectional study using AWaRe classification and WHO antibiotic prescribing indicators in Dire Dawa, Ethiopia. *Drugs Real World Outcomes* 2023; **10**: 459–69. <https://doi.org/10.1007/s40801-023-00367-1>
- 154** Sulis G, Daniels B, Kwan A et al. Antibiotic overuse in the primary health care setting: a secondary data analysis of standardised patient studies from India, China and Kenya. *BMJ Glob Health* 2020; **5**: e003393. <https://doi.org/10.1136/bmjgh-2020-003393>
- 155** Thi TVL, Canh Pham E, Dang-Nguyen DT. Evaluation of children's antibiotics use for outpatient pneumonia treatment in Vietnam. *Braz J Infect Dis* 2024; **28**: 103839. <https://doi.org/10.1016/j.bjid.2024.103839>
- 156** Kabba JA, Tadesse N, James PB et al. Knowledge, attitude and antibiotic prescribing patterns of medical doctors providing free healthcare in the outpatient departments of public hospitals in Sierra Leone: a national cross-sectional study. *Trans R Soc Trop Med Hyg* 2020; **114**: 448–58. <https://doi.org/10.1093/trstmh/trz137>
- 157** Chukwu EE, Oladele DA, Enwuru CA et al. Antimicrobial resistance awareness and antibiotic prescribing behavior among healthcare workers in Nigeria: a national survey. *BMC Infect Dis* 2021; **21**: 22. <https://doi.org/10.1186/s12879-020-05689-x>
- 158** Farley E, Stewart A, Davies MA et al. Antibiotic use and resistance: knowledge, attitudes and perceptions among primary care prescribers in South Africa. *S Afr Med J* 2018; **108**: 763–71. <https://doi.org/10.7196/SAMJ.2018.v108i9.12933>
- 159** Maraqa B, Nazzal Z, Hamshari S et al. Palestinian physicians' self-reported practice regarding antibiotic use for upper respiratory tract infections in primary healthcare. *Front Med* 2023; **10**: 1139871. <https://doi.org/10.3389/fmed.2023.1139871>
- 160** Shu G, Jayawardena K, Jayaweera Patabandige D et al. Knowledge, perceptions and practices on antibiotic use among Sri Lankan doctors. *PLoS One* 2022; **17**: e0263167. <https://doi.org/10.1371/journal.pone.0263167>
- 161** Sychareun V, Sihavong A, Machowska A et al. Knowledge, attitudes, perception and reported practices of healthcare providers on antibiotic use and resistance in pregnancy, childbirth and children under two in Lao PDR: a mixed methods study. *Antibiotics* 2021; **10**: 1462. <https://doi.org/10.3390/antibiotics10121462>
- 162** Lim AH, Ab Rahman N, Ong SM et al. Impact evaluation of guidelines on antibiotic utilisation and appropriateness in Malaysian public primary care: an interrupted time series analysis. *J Pharm Policy Pract* 2024; **17**: 2355666. <https://doi.org/10.1080/20523211.2024.2355666>
- 163** Igirikwayo ZK, Migisha R, Mukaga H et al. Prescription patterns of antibiotics and associated factors among outpatients diagnosed with respiratory tract infections in Jinja city, Uganda, June 2022–May 2023. *BMC Pulm Med* 2024; **24**: 446. <https://doi.org/10.1186/s12890-024-03246-9>
- 164** Kleczka B, Kumar P, Njeru MK et al. Using rubber stamps and mobile phones to help understand and change antibiotic prescribing behaviour in private sector primary healthcare clinics in Kenya. *BMJ Glob Health* 2019; **4**: e001422. <https://doi.org/10.1136/bmjgh-2019-001422>
- 165** De Vries E, Johnson Y, Willems B et al. Improving primary care antimicrobial stewardship by implementing a peer audit and feedback intervention in Cape Town community healthcare centres. *S Afr Med J* 2022; **112**: 812–8. <https://doi.org/10.7196/SAMJ.2022.v112i10.16397>
- 166** Chandran DS, Manickavasagam PP. Sale of antibiotics without prescription in stand-alone pharmacies in Tamil Nadu. *J Family Med Prim Care* 2022; **11**: 5516–20. [https://doi.org/10.4103/jfmpc.jfmpc\\_2157\\_21](https://doi.org/10.4103/jfmpc.jfmpc_2157_21)
- 167** Ayele AA, Mekuria AB, Tegegn HG et al. Management of minor ailments in a community pharmacy setting: findings from simulated visits and qualitative study in Gondar town, Ethiopia. *PLoS One* 2018; **13**: e0190583. <https://doi.org/10.1371/journal.pone.0190583>
- 168** Alkadhimi A, Dawood OT, Hassali MA. Dispensing of antibiotics in community pharmacy in Iraq: a qualitative study. *Pharm Pract* 2020; **18**: 2095. <https://doi.org/10.18549/PharmPract.2020.4.2095>
- 169** Nguyen HH, Ho DP, Vu TLH et al. “I can make more from selling medicine when breaking the rules”—understanding the antibiotic supply network in a rural community in Viet Nam. *BMC Public Health* 2019; **19**: 1560. <https://doi.org/10.1186/s12889-019-7812-z>
- 170** Gacheri J, Hamilton KA, Munywoki P et al. Antibiotic prescribing practices in community and clinical settings during the COVID-19 pandemic in Nairobi, Kenya. *PLoS Glob Public Health* 2024; **4**: e0003046. <https://doi.org/10.1371/journal.pgph.0003046>
- 171** Islam MA, Akhtar Z, Hassan MZ et al. Pattern of antibiotic dispensing at pharmacies according to the WHO access, watch, reserve (AWaRe) classification in Bangladesh. *Antibiotics* 2022; **11**: 247. <https://doi.org/10.3390/antibiotics11020247>
- 172** Saleem Z, Hassali MA, Godman B et al. Sale of WHO AWaRe groups antibiotics without a prescription in Pakistan: a simulated client study. *J Pharm Policy Pract* 2020; **13**: 26. <https://doi.org/10.1186/s40545-020-00233-3>
- 173** Koju P, Rousseau SP, Van der Putten M et al. Advertisement of antibiotics for upper respiratory infections and equity in access to treatment: a cross-sectional study in Nepal. *J Pharm Policy Pract* 2020; **13**: 4. <https://doi.org/10.1186/s40545-020-0202-1>
- 174** Erku DA, Abera SY. Non-prescribed sale of antibiotics for acute childhood diarrhea and upper respiratory tract infection in community pharmacies: a 2 phase mixed-methods study. *Antimicrob Resist Infect Control* 2018; **7**: 92. <https://doi.org/10.1186/s13756-018-0389-y>
- 175** Edessa D, Kumsa FA, Dinsa G et al. Drug providers' perspectives on antibiotic misuse practices in eastern Ethiopia: a qualitative study. *BMJ Open* 2024; **14**: e085352. <https://doi.org/10.1136/bmjopen-2024-085352>
- 176** Muloi D, Fèvre EM, Bettridge J et al. A cross-sectional survey of practices and knowledge among antibiotic retailers in Nairobi, Kenya. *J Glob Health* 2019; **9**: 010412. <https://doi.org/10.7189/jogh.09.020412>
- 177** Ndaki PM, Mwanga JR, Mushi MF et al. Practices and motives behind antibiotics provision in drug outlets in Tanzania: a qualitative study. *PLoS One* 2023; **18**: e0290638. <https://doi.org/10.1371/journal.pone.0290638>
- 178** Acharya Y, Nepal P, Yang D et al. Economic and social drivers of antibiotic dispensing practices among community pharmacies in Nepal. *Trop Med Int Health* 2021; **26**: 557–71. <https://doi.org/10.1111/tmi.13555>
- 179** Ngyedu EK, Acolatse J, Akafity G et al. Selling antibiotics without prescriptions among community pharmacies and drug outlets: a simulated client study from Ghana. *Expert Rev Anti Infect Ther* 2023; **21**: 1373–82. <https://doi.org/10.1080/14787210.2023.2283037>
- 180** Myemba DT, Maganda BA, Kibwana UO et al. Profiling of antimicrobial dispensing practices in accredited drug dispensing outlets in Tanzania: a mixed-method cross-sectional study focusing on pediatric patients. *BMC Health Serv Res* 2022; **22**: 1575. <https://doi.org/10.1186/s12913-022-08980-6>

- 181** Ndaki PM, Mushi MF, Mwanga JR *et al.* Dispensing antibiotics without prescription at community pharmacies and accredited drug dispensing outlets in Tanzania: a cross-sectional study. *Antibiotics* 2021; **10**: 1025. <https://doi.org/10.3390/antibiotics10081025>
- 182** Matin MA, Khan WA, Karim MM *et al.* What influences antibiotic sales in rural Bangladesh? A drug dispensers' perspective. *J Pharm Policy Pract* 2020; **13**: 20. <https://doi.org/10.1186/s40545-020-00212-8>
- 183** Nizame FA, Shoaib DM, Rousham EK *et al.* Barriers and facilitators to adherence to national drug policies on antibiotic prescribing and dispensing in Bangladesh. *J Pharm Policy Pract* 2021; **14**: 85. <https://doi.org/10.1186/s40545-021-00342-7>
- 184** Rakhshani NS, Kaljee LM, Khan MI *et al.* A formative assessment of antibiotic dispensing/prescribing practices and knowledge and perceptions of antimicrobial resistance (AMR) among healthcare workers in Lahore Pakistan. *Antibiotics* 2022; **11**: 1418. <https://doi.org/10.3390/antibiotics11101418>
- 185** Godman B, Haque M, Islam S *et al.* Rapid assessment of price instability and paucity of medicines and protection for COVID-19 across Asia: findings and public health implications for the future. *Front Public Health* 2020; **8**: 585832. <https://doi.org/10.3389/fpubh.2020.585832>
- 186** Mukokinya MMA, Opanga S, Oluka M *et al.* Dispensing of antimicrobials in Kenya: a cross-sectional pilot study and its implications. *J Res Pharm Pract* 2018; **7**: 77–82. [https://doi.org/10.4103/jrpp.JRPP\\_17\\_88](https://doi.org/10.4103/jrpp.JRPP_17_88)
- 187** Opanga S, Rizvi N, Wamaita A *et al.* Availability of medicines in community pharmacy to manage patients with COVID-19 in Kenya: pilot study and implications. *Sch Acad J Pharm* 2021; **3**: 36–42. <https://doi.org/10.36347/sajp.2021.v10i03.001>
- 188** Abubakar U, Tangiisuran B. Knowledge and practices of community pharmacists towards non-prescription dispensing of antibiotics in northern Nigeria. *Int J Clin Pharm* 2020; **42**: 756–64. <https://doi.org/10.1007/s11096-020-01019-y>
- 189** Bagonza A, Kitutu FE, Peterson S *et al.* Effectiveness of peer-supervision on pediatric fever illness treatment among registered private drug sellers in east-central Uganda: an interrupted time series analysis. *Health Sci Rep* 2021; **4**: e284. <https://doi.org/10.1002/hsr2.284>
- 190** Poyongo BP, Sangeda RZ. Pharmacists' knowledge, attitude and practice regarding the dispensing of antibiotics without prescription in Tanzania: an explorative cross-sectional study. *Pharmacy* 2020; **8**: 238. <https://doi.org/10.3390/pharmacy8040238>
- 191** Greene HC, Makovi K, Abdul-Mumin R *et al.* Challenges in the distribution of antimicrobial medications in community dispensaries in Accra, Ghana. *PLoS One* 2024; **19**: e0281699. <https://doi.org/10.1371/journal.pone.0281699>
- 192** Zulu A, Matafwali SK, Banda M *et al.* Assessment of knowledge, attitude and practices on antibiotic resistance among undergraduate medical students in the school of medicine at the University of Zambia. *Int J Basic Clin Pharmacol* 2020; **9**: 263–2701. <https://doi.org/10.18203/2319-2003.ijbcp20200174>
- 193** Kamoto A, Chapotera G, Suleman F. Knowledge, attitude and perception on antimicrobial use and antimicrobial resistance among final year medical students in the college of medicine, Malawi. *Malawi Med J* 2020; **32**: 120–3. <https://doi.org/10.4314/mmj.v32i3.3>
- 194** Okedo-Alex I, Madubueze UC, Umeokonkwo CD *et al.* Knowledge of antibiotic use and resistance among students of a medical school in Nigeria. *Malawi Med J* 2019; **31**: 133–7. <https://doi.org/10.4314/mmj.v31i2.5>
- 195** Augie BM, van Zyl RL, McInerney PA *et al.* Knowledge and perceptions about antibiotic resistance and prudent antibiotic prescribing among final year medical students in two African countries. *Int J Pharm Pract* 2021; **29**: 508–14. <https://doi.org/10.1093/ijpp/riab044>
- 196** Nisabwe L, Brice H, Umuhire MC *et al.* Knowledge and attitudes towards antibiotic use and resistance among undergraduate healthcare students at university of Rwanda. *J Pharm Policy Pract* 2020; **13**: 7. <https://doi.org/10.1186/s40545-020-00207-5>
- 197** Kanyike AM, Olum R, Kajimu J *et al.* Antimicrobial resistance and rational use of medicine: knowledge, perceptions, and training of clinical health professions students in Uganda. *Antimicrob Resist Infect Control* 2022; **11**: 145. <https://doi.org/10.1186/s13756-022-01186-9>
- 198** Lubwama M, Onyuka J, Ayazika KT *et al.* Knowledge, attitudes, and perceptions about antibiotic use and antimicrobial resistance among final year undergraduate medical and pharmacy students at three universities in east Africa. *PLoS One* 2021; **16**: e0251301. <https://doi.org/10.1371/journal.pone.0251301>
- 199** Chuwa BB, Njau LA, Msigwa KI *et al.* Prevalence and factors associated with self medication with antibiotics among university students in Moshi Kilimanjaro Tanzania. *Afr Health Sci* 2021; **21**: 633–9. <https://doi.org/10.4314/ahs.v21i2.19>
- 200** Sefah IA, Akwaboah E, Sarkodie E *et al.* Evaluation of healthcare students' knowledge on antibiotic use, antimicrobial resistance and antimicrobial stewardship programs and associated factors in a tertiary university in Ghana: findings and implications. *Antibiotics* 2022; **11**: 1679. <https://doi.org/10.3390/antibiotics11121679>
- 201** Owusu-Ofori AK, Darko E, Danquah CA *et al.* Self-medication and antimicrobial resistance: a survey of students studying healthcare programmes at a tertiary institution in Ghana. *Front Public Health* 2021; **9**: 706290. <https://doi.org/10.3389/fpubh.2021.706290>
- 202** Fetensa G, Wakuma B, Tolossa T *et al.* Knowledge and attitude towards antimicrobial resistance of graduating health science students of Wollega University. *Infect Drug Resist* 2020; **13**: 3937–44. <https://doi.org/10.2147/IDR.S264481>
- 203** Ayepola OO, Onile-Ere OA, Shodeko OE *et al.* Dataset on the knowledge, attitudes and practices of university students towards antibiotics. *Data Brief* 2018; **19**: 2084–94. <https://doi.org/10.1016/j.dib.2018.06.090>
- 204** Akande-Sholabi W, Ajamu AT. Antimicrobial stewardship: assessment of knowledge, awareness of antimicrobial resistance and appropriate antibiotic use among healthcare students in a Nigerian university. *BMC Med Educ* 2021; **21**: 488. <https://doi.org/10.1186/s12909-021-02912-4>
- 205** Shitindi L, Issa O, Poyongo BP *et al.* Comparison of knowledge, attitude, practice and predictors of self-medication with antibiotics among medical and non-medical students in Tanzania. *Front Pharmacol* 2023; **14**: 1301561. <https://doi.org/10.3389/fphar.2023.1301561>
- 206** Nakato G, Adongo PR, Iramiot JS *et al.* Practices and drivers of self-medication with antibiotics among undergraduate medical students in eastern Uganda: a cross-sectional study. *PLoS One* 2023; **18**: e0293685. <https://doi.org/10.1371/journal.pone.0293685>
- 207** Amponsah SK, Odamtten G, Adams I *et al.* A comparative analysis of pattern and attitude towards self-medication among pharmacy and non-pharmacy students in university of Ghana. *Pan Afr Med J* 2022; **41**: 254. <https://doi.org/10.11604/pamj.2022.41.254.31013>
- 208** Ajibola O, Omisakin OA, Eze AA *et al.* Self-medication with antibiotics, attitude and knowledge of antibiotic resistance among community residents and undergraduate students in northwest Nigeria. *Diseases* 2018; **6**: 32. <https://doi.org/10.3390/diseases6020032>
- 209** Tuyishimire J, Okoya F, Adebayo AY *et al.* Assessment of self-medication practices with antibiotics among undergraduate university students in Rwanda. *Pan Afr Med J* 2019; **33**: 307. <https://doi.org/10.11604/pamj.2019.33.307.18139>
- 210** Abuawad M, Ziyadeh-Isleem A, Mahamid A *et al.* Knowledge, perception, and attitudes of medical students towards antimicrobial resistance and stewardship: an observational cross-sectional study from Palestine. *BMC Med Educ* 2024; **24**: 302. <https://doi.org/10.1186/s12909-024-05276-7>



- 211** Assar A, Abdelraoof MI, Abdel-Maboud M *et al.* Knowledge, attitudes, and practices of Egypt's future physicians towards antimicrobial resistance (KAP-AMR study): a multicenter cross-sectional study. *Environ Sci Pollut Res Int* 2020; **27**: 21292–8. <https://doi.org/10.1007/s11356-020-08534-5>
- 212** Al-Taani GM, Karasneh RA, Al-Azzam S *et al.* Knowledge, attitude, and behavior about antimicrobial use and resistance among medical, nursing and pharmacy students in Jordan: a cross sectional study. *Antibiotics* 2022; **11**: 1559. <https://doi.org/10.3390/antibiotics11111559>
- 213** Hussain I, Yousaf N, Haider S *et al.* Assessing knowledge and perception regarding antimicrobial stewardship and antimicrobial resistance in university students of Pakistan: findings and implications. *Antibiotics* 2021; **10**: 866. <https://doi.org/10.3390/antibiotics10070866>
- 214** Shah S, Abbas G, Chauhdary Z *et al.* Antibiotic use: a cross-sectional survey assessing the knowledge, attitudes, and practices amongst students of Punjab, Pakistan. *J Am Coll Health* 2022; **70**: 2499–504. <https://doi.org/10.1080/07448481.2020.1865984>
- 215** Elmahi OKO, Musa RAE, Shareef AAH *et al.* Perception and practice of self-medication with antibiotics among medical students in Sudanese universities: a cross-sectional study. *PLoS One* 2022; **17**: e0263067. <https://doi.org/10.1371/journal.pone.0263067>
- 216** Naser AY, Aboutaleb R, Khaleel A *et al.* Knowledge, attitude, and practices of pharmacy students in 7 Middle Eastern countries concerning antibiotic resistance: a cross-sectional study. *Medicine* 2024; **103**: e39378. <https://doi.org/10.1097/MD.00000000000039378>
- 217** Mostafa A, Abdelzaher A, Rashed S *et al.* Is health literacy associated with antibiotic use, knowledge and awareness of antimicrobial resistance among non-medical university students in Egypt? A cross-sectional study. *BMJ Open* 2021; **11**: e046453. <https://doi.org/10.1136/bmjopen-2020-046453>
- 218** Hayat K, Fatima N, Umer MF *et al.* Understanding of future prescribers about antimicrobial resistance and their preparedness towards antimicrobial stewardship activities in Pakistan: findings and implications. *Front Pharmacol* 2022; **13**: 771083. <https://doi.org/10.3389/fphar.2022.771083>
- 219** Abdelkarim OA, Abubakar U, Hussain MA *et al.* Knowledge, perception, and self-confidence of antibiotic resistance, appropriate antibiotic therapy, and antibiotic stewardship among undergraduate pharmacy students in Sudan. *Infect Drug Resist* 2024; **17**: 935–49. <https://doi.org/10.2147/IDR.S435190>
- 220** Hayat K, Jamshed S, Rosenthal M *et al.* Understanding of pharmacy students towards antibiotic use, antibiotic resistance and antibiotic stewardship programs: a cross-sectional study from Punjab, Pakistan. *Antibiotics* 2021; **10**: 66. <https://doi.org/10.3390/antibiotics10010066>
- 221** Alsayed AR, Darwish El Hajji F, Al-Najjar MAA *et al.* Patterns of antibiotic use, knowledge, and perceptions among different population categories: a comprehensive study based in Arabic countries. *Saudi Pharm J* 2022; **30**: 317–28. <https://doi.org/10.1016/j.jsps.2022.01.013>
- 222** Gupta MK, Vohra C, Raghav P. Assessment of knowledge, attitudes, and practices about antibiotic resistance among medical students in India. *J Family Med Prim Care* 2019; **8**: 2864–9. [https://doi.org/10.4103/jfmpc.jfmpc\\_504\\_19](https://doi.org/10.4103/jfmpc.jfmpc_504_19)
- 223** Ritchie O, Shetty V, Prabhu S *et al.* Confidence in antibiotic prescribing intentions among senior medical students in India. *Am J Trop Med Hyg* 2020; **103**: 2561–7. <https://doi.org/10.4269/ajtmh.20-0193>
- 224** Marzan M, Islam DZ, Lugova H *et al.* Knowledge, attitudes, and practices of antimicrobial uses and resistance among public university students in Bangladesh. *Infect Drug Resist* 2021; **14**: 519–33. <https://doi.org/10.2147/IDR.S289964>
- 225** Shrestha R. Knowledge, attitude and practice on antibiotics use and its resistance among medical students in a tertiary care hospital. *J Nepal Med Assoc* 2019; **57**: 74–9. <https://doi.org/10.31729/jnma.4224>
- 226** Abubakar U, Muhammad HT, Sulaiman SAS *et al.* Knowledge and self-confidence of antibiotic resistance, appropriate antibiotic therapy, and antibiotic stewardship among pharmacy undergraduate students in three Asian countries. *Curr Pharm Teach Learn* 2020; **12**: 265–73. <https://doi.org/10.1016/j.cptl.2019.12.002>
- 227** Shah P, Shrestha R, Mao Z *et al.* Knowledge, attitude, and practice associated with antibiotic use among university students: a survey in Nepal. *Int J Environ Res Public Health* 2019; **16**: 3996. <https://doi.org/10.3390/ijerph16203996>
- 228** Sakeena MH, Bennett AA, Mohamed F *et al.* Investigating knowledge regarding antibiotics among pharmacy and allied health sciences students in a Sri Lankan university. *J Infect Dev Ctries* 2018; **12**: 726–32. <https://doi.org/10.3855/jidc.10388>
- 229** Sakeena MHF, Bennett AA, Jamshed S *et al.* Investigating knowledge regarding antibiotics and antimicrobial resistance among pharmacy students in Sri Lankan universities. *BMC Infect Dis* 2018; **18**: 209. <https://doi.org/10.1186/s12879-018-3107-8>
- 230** Jayaweerasingham M, Angulmaduwa S, Liyanapathirana V. Knowledge, beliefs and practices on antibiotic use and resistance among a group of trainee nurses in Sri Lanka. *BMC Res Notes* 2019; **12**: 601. <https://doi.org/10.1186/s13104-019-4640-2>
- 231** Sakeena MHF, Bennett AA, Carter SJ *et al.* A comparative study regarding antibiotic consumption and knowledge of antimicrobial resistance among pharmacy students in Australia and Sri Lanka. *PLoS One* 2019; **14**: e0213520. <https://doi.org/10.1371/journal.pone.0213520>
- 232** Nabi N, Baluja Z, Mukherjee S *et al.* Trends in practices of self-medication with antibiotics among medical undergraduates in India. *J Pharm Bioallied Sci* 2022; **14**: 19–24. [https://doi.org/10.4103/jpbs.jpbs\\_17\\_21](https://doi.org/10.4103/jpbs.jpbs_17_21)
- 233** Mandal NK, Rauniyar GP, Rai DS *et al.* Self-medication practice of antibiotics among medical and dental undergraduate students in a medical college in eastern Nepal: a descriptive cross-sectional study. *J Nepal Med Assoc* 2020; **58**: 328–32. <https://doi.org/10.31729/jnma.4914>
- 234** Shah K, Halder S, Haider SS. Assessment of knowledge, perception, and awareness about self-medication practices among university students in Nepal. *Heliyon* 2021; **7**: e05976. <https://doi.org/10.1016/j.heliyon.2021.e05976>
- 235** Paul GK, Swapon MS, Kaderi Kibria KM. Knowledge, awareness, and attitudes toward antibiotic resistance and practice of self-medication among university students in Bangladesh: a cross-sectional study. *J Educ Health Promot* 2022; **11**: 115. [https://doi.org/10.4103/jehp.jehp\\_602\\_21](https://doi.org/10.4103/jehp.jehp_602_21)
- 236** Wahab A, Alam MM, Hasan S *et al.* Exploring the knowledge, practices and determinants of antibiotic self-medication among Bangladeshi university students in the era of COVID-19: a cross-sectional study. *Heliyon* 2023; **9**: e19923. <https://doi.org/10.1016/j.heliyon.2023.e19923>
- 237** Yang C, Xie J, Chen Q *et al.* Knowledge, attitude, and practice about antibiotic use and antimicrobial resistance among nursing students in China: a cross sectional study. *Infect Drug Resist* 2024; **17**: 1085–98. <https://doi.org/10.2147/IDR.S454489>
- 238** Seam MOR, Bhatta R, Saha BL *et al.* Assessing the perceptions and practice of self-medication among Bangladeshi undergraduate pharmacy students. *Pharmacy* 2018; **6**: 6. <https://doi.org/10.3390/pharmacy6010006>
- 239** Hu Y, Wang X, Tucker JD *et al.* Knowledge, attitude, and practice with respect to antibiotic use among Chinese medical students: a multicentre cross-sectional study. *Int J Environ Res Public Health* 2018; **15**: 1165. <https://doi.org/10.3390/ijerph15061165>
- 240** Lin L, Fearon E, Harbarth S *et al.* Decisions to use antibiotics for upper respiratory tract infections across China: a large-scale cross-sectional survey among university students. *BMJ Open* 2020; **10**: e039332. <https://doi.org/10.1136/bmjopen-2020-039332>

- 241 Min S, Zhou Y, Sun Y et al. Knowledge, attitude, and practice associated with antimicrobial resistance among medical students between 2017 and 2022: a survey in east China. *Front Public Health* 2022; **10**: 1010582. <https://doi.org/10.3389/fpubh.2022.1010582>
- 242 Haque M, Rahman NAA, McKimm J et al. Antibiotic use: a cross-sectional study evaluating the understanding, usage and perspectives of medical students and pathfinders of a public defence university in Malaysia. *Antibiotics* 2019; **8**: 154. <https://doi.org/10.3390/antibiotics8030154>
- 243 Wang Y, Guo F, Wei J et al. Knowledge, attitudes and practices in relation to antimicrobial resistance amongst Chinese public health undergraduates. *J Glob Antimicrob Resist* 2020; **23**: 9–15. <https://doi.org/10.1016/j.jgar.2020.07.023>
- 244 Haque M, Rahman NAA, McKimm J et al. A cross-sectional study evaluating the knowledge and beliefs about, and the use of antibiotics amongst Malaysian university students. *Expert Rev Anti Infect Ther* 2019; **17**: 275–84. <https://doi.org/10.1080/14787210.2019.1581607>
- 245 Peng D, Wang X, Xu Y et al. Antibiotic misuse among university students in developed and less developed regions of China: a cross-sectional survey. *Glob Health Action* 2018; **11**: 1496973. <https://doi.org/10.1080/16549716.2018.1496973>
- 246 Haque M, Ara T, Haq MA et al. Antimicrobial prescribing confidence and knowledge regarding drug resistance: perception of medical students in Malaysia and the implications. *Antibiotics* 2022; **11**: 540. <https://doi.org/10.3390/antibiotics11050540>
- 247 Ahmed S, Tareq AH, Ilyas D. The impact of antimicrobial resistance and stewardship training sessions on knowledge of healthcare students of Wah cantonment, Pakistan. *Inquiry* 2024; **61**: 469580241228443. <https://doi.org/10.1177/00469580241228443>
- 248 Shembo AKP, Musumari PM, Srithanaviboonchai K et al. A qualitative study on community use of antibiotics in Kinshasa, Democratic Republic of Congo. *PLoS One* 2022; **17**: e0267544. <https://doi.org/10.1371/journal.pone.0267544>
- 249 Mengesha Y, Manaye B, Moges G. Assessment of public awareness, attitude, and practice regarding antibiotic resistance in Kemissie town, northeast Ethiopia: community-based cross-sectional study. *Infect Drug Resist* 2020; **13**: 3783–9. <https://doi.org/10.2147/IDR.S280036>
- 250 Dejene H, Birhanu R, Tarekegn ZS. Knowledge, attitude and practices of residents toward antimicrobial usage and resistance in Gondar, Northwest Ethiopia. *One Health Outlook* 2022; **4**: 10. <https://doi.org/10.1186/s42522-022-00066-x>
- 251 Simegn W, Moges G. Awareness and knowledge of antimicrobial resistance and factors associated with knowledge among adults in Dessie city, northeast Ethiopia: community-based cross-sectional study. *PLoS One* 2022; **17**: e0279342. <https://doi.org/10.1371/journal.pone.0279342>
- 252 Muhammed AM, Alemu A, Maidane YO et al. Knowledge, attitudes, and practices of rural communities regarding antimicrobial resistance and climate change in Adadle district, Somali region, Ethiopia: a mixed-methods study. *Antibiotics* 2024; **13**: 292. <https://doi.org/10.3390/antibiotics13040292>
- 253 Ekambi GA E, Ebongue C O, Penda IC et al. Knowledge, practices and attitudes on antibiotics use in Cameroon: self-medication and prescription survey among children, adolescents and adults in private pharmacies. *PLoS One* 2019; **14**: e0212875. <https://doi.org/10.1371/journal.pone.0212875>
- 254 Vicar EK, Walana W, Mbabila A et al. Drivers of household antibiotic use in urban informal settlements in northern Ghana: implications for antimicrobial resistance control. *Health Sci Rep* 2023; **6**: e1388. <https://doi.org/10.1002/hsr2.1388>
- 255 Effah CY, Amoah AN, Liu H et al. A population-base survey on knowledge, attitude and awareness of the general public on antibiotic use and resistance. *Antimicrob Resist Infect Control* 2020; **9**: 105. <https://doi.org/10.1186/s13756-020-00768-9>
- 256 Horumpende PG, Said SH, Mazuguni FS et al. Prevalence, determinants and knowledge of antibacterial self-medication: a cross sectional study in north-eastern Tanzania. *PLoS One* 2018; **13**: e0206623. <https://doi.org/10.1371/journal.pone.0206623>
- 257 Bassoum O, Sougou NM, Diongue M et al. Assessment of general public's knowledge and opinions towards antibiotic use and bacterial resistance: a cross-sectional study in an urban setting, Rufisque, Senegal. *Pharmacy* 2018; **6**: 103. <https://doi.org/10.3390/pharmacy6040103>
- 258 Mudenda S, Simukoko N, Mohamed S. Knowledge, attitude and practices regarding antimicrobial use and resistance among community members of Mtendere township in Lusaka, Zambia: findings and implications on antimicrobial stewardship. *Int J Basic Clin Pharmacol* 2024; **13**: 315–21. <https://doi.org/10.18203/2319-2003.ijbcp20240985>
- 259 Kampamba M, Hamaambo B, Hikaambo CN et al. Evaluation of knowledge and practices on antibiotic use: a cross-sectional study on self-reported adherence to short-term antibiotic utilization among patients visiting level-1 hospitals in Lusaka, Zambia. *JAC Antimicrob Resist* 2024; **6**: dlae120. <https://doi.org/10.1093/jacamr/dlae120>
- 260 Chukwu EE, Oladele DA, Awoderu OB et al. A national survey of public awareness of antimicrobial resistance in Nigeria. *Antimicrob Resist Infect Control* 2020; **9**: 72. <https://doi.org/10.1186/s13756-020-00739-0>
- 261 Isah A, Aina AB, Ben-Umeh KC et al. Assessment of public knowledge and attitude toward antibiotics use and resistance: a community pharmacy-based survey. *J Pharm Policy Pract* 2023; **16**: 107. <https://doi.org/10.1186/s40545-023-00619-z>
- 262 Farley E, van den Bergh D, Coetzee R et al. Knowledge, attitudes and perceptions of antibiotic use and resistance among patients in South Africa: a cross-sectional study. *S Afr J Infect Dis* 2019; **34**: 118. <https://doi.org/10.4102/sajid.v34i1.118>
- 263 Mutagonda RF, Marealle AI, Nkinda L et al. Determinants of misuse of antibiotics among parents of children attending clinics in regional referral hospitals in Tanzania. *Sci Rep* 2022; **12**: 4836. <https://doi.org/10.1038/s41598-022-08895-6>
- 264 Sambakunsi CS, Småbrekke L, Varga CA et al. Knowledge, attitudes and practices related to self-medication with antimicrobials in Lilongwe, Malawi. *Malawi Med J* 2019; **31**: 225–32. <https://doi.org/10.4314/mmj.v31i4.2>
- 265 Sindato C, Mboera LEG, Katala BZ et al. Knowledge, attitudes and practices regarding antimicrobial use and resistance among communities of Ilala, Kilosa and Kibaha districts of Tanzania. *Antimicrob Resist Infect Control* 2020; **9**: 194. <https://doi.org/10.1186/s13756-020-00862-y>
- 266 Mboya EA, Davies ML, Horumpende PG et al. Inadequate knowledge on appropriate antibiotics use among clients in the Moshi municipality northern Tanzania. *PLoS One* 2020; **15**: e0239388. <https://doi.org/10.1371/journal.pone.0239388>
- 267 Gabriel S, Manumbu L, Mkusa O et al. Knowledge of use of antibiotics among consumers in Tanzania. *JAC Antimicrob Resist* 2021; **3**: dlab183. <https://doi.org/10.1093/jacamr/dlab183>
- 268 Mokoena TTW, Schellack N, Brink AJ. Driving antibiotic stewardship awareness through the minibus-taxi community across the Tshwane district, South Africa—a baseline evaluation. *JAC Antimicrob Resist* 2021; **3**: dlab106. <https://doi.org/10.1093/jacamr/dlab106>
- 269 Lanyero H, Eriksen J, Obua C et al. Use of antibacterials in the management of symptoms of acute respiratory tract infections among children under five years in Gulu, northern Uganda: prevalence and determinants. *PLoS One* 2020; **15**: e0235164. <https://doi.org/10.1371/journal.pone.0235164>
- 270 Ngoma MT, Sitali D, Mudenda S et al. Community antibiotic consumption and associated factors in Lusaka district of Zambia: findings and

- implications for antimicrobial resistance and stewardship. *JAC Antimicrob Resist* 2024; **6**: dlac034. <https://doi.org/10.1093/jacamr/dlae034>
- 271** Russom M, Bahta M, Debesai M et al. Knowledge, attitude and practice of antibiotics and their determinants in Eritrea: an urban population-based survey. *BMJ Open* 2021; **11**: e046432. <https://doi.org/10.1136/bmjopen-2020-046432>
- 272** Wildbret S, Stuck L, Luchen CC et al. Drivers of informal sector and non-prescription medication use in pediatric populations in a low- and middle-income setting: a prospective cohort study in Zambia. *PLoS Glob Public Health* 2023; **3**: e0002072. <https://doi.org/10.1371/journal.pgph.0002072>
- 273** Bogale AA, Amhare AF, Chang J et al. Knowledge, attitude, and practice of self-medication with antibiotics among community residents in Addis Ababa, Ethiopia. *Expert Rev Anti Infect Ther* 2019; **17**: 459–66. <https://doi.org/10.1080/14787210.2019.1620105>
- 274** Mate I, Come CE, Gonçalves MP et al. Knowledge, attitudes and practices regarding antibiotic use in Maputo city, Mozambique. *PLoS One* 2019; **14**: e0221452. <https://doi.org/10.1371/journal.pone.0221452>
- 275** Cambaco O, Alonso Menendez Y, Kinsman J et al. Community knowledge and practices regarding antibiotic use in rural Mozambique: where is the starting point for prevention of antibiotic resistance? *BMC Public Health* 2020; **20**: 1183. <https://doi.org/10.1186/s12889-020-09243-x>
- 276** Jimah T, Fenny AP, Ogunseitan OA. Antibiotics stewardship in Ghana: a cross-sectional study of public knowledge, attitudes, and practices among communities. *One Health Outlook* 2020; **2**: 12. <https://doi.org/10.1186/s42522-020-00021-8>
- 277** Akande-Sholabi W, Oyesiji E. Antimicrobial stewardship: knowledge, perceptions, and factors associated with antibiotics misuse among consumer's visiting the community pharmacies in a Nigeria southwestern state. *J Pharm Policy Pract* 2023; **16**: 120. <https://doi.org/10.1186/s40545-023-00629-x>
- 278** Kiragga AN, Najjemba L, Galiwango R et al. Community purchases of antimicrobials during the COVID-19 pandemic in Uganda: an increased risk for antimicrobial resistance. *PLoS Glob Public Health* 2023; **3**: e0001579. <https://doi.org/10.1371/journal.pgph.0001579>
- 279** Eibs T, Koscalova A, Nair M et al. Qualitative study of antibiotic prescription patterns and associated drivers in Sudan, Guinea-Bissau, Central African Republic and Democratic Republic of Congo. *BMJ Open* 2020; **10**: e036530. <https://doi.org/10.1136/bmjopen-2019-036530>
- 280** Ateshim Y, Bereket B, Major F et al. Prevalence of self-medication with antibiotics and associated factors in the community of Asmara, Eritrea: a descriptive cross sectional survey. *BMC Public Health* 2019; **19**: 726. <https://doi.org/10.1186/s12889-019-7020-x>
- 281** Demissie F, Ereso K, Paulos G. Self-medication practice with antibiotics and its associated factors among community of Bule-Hora town, south west Ethiopia. *Drug Healthc Patient Saf* 2022; **14**: 9–18. <https://doi.org/10.2147/DHPS.S325150>
- 282** Machongo RB, Mipando ALN. “I don’t hesitate to use the left-over antibiotics for my child”: practices and experiences with antibiotic use among caregivers of paediatric patients at Zomba central hospital in Malawi. *BMC Pediatr* 2022; **22**: 466. <https://doi.org/10.1186/s12887-022-03528-3>
- 283** Ahiabu MA, Magnussen P, Bygbjerg IC et al. Treatment practices of households and antibiotic dispensing in medicine outlets in developing countries: the case of Ghana. *Res Social Adm Pharm* 2018; **14**: 1180–8. <https://doi.org/10.1016/j.sapharm.2018.01.013>
- 284** Mboya EA, Sanga LA, Ngocho JS. Irrational use of antibiotics in the Moshi municipality northern Tanzania: a cross sectional study. *Pan Afr Med J* 2018; **31**: 165. <https://doi.org/10.11604/pamj.2018.31.165.15991>
- 285** Kimathi G, Kiarie J, Njarambah L et al. A cross-sectional study of antimicrobial use among self-medicating COVID-19 cases in Nyeri county, Kenya. *Antimicrob Resist Infect Control* 2022; **11**: 111. <https://doi.org/10.1186/s13756-022-01150-7>
- 286** Nyeko R, Otim F, Obiya EM et al. Pre-hospital exposures to antibiotics among children presenting with fever in northern Uganda: a facility-based cross-sectional study. *BMC Pediatr* 2022; **22**: 322. <https://doi.org/10.1186/s12887-022-03375-2>
- 287** Afari-Asiedu S, Oppong FB, Tostmann A et al. Determinants of inappropriate antibiotics use in rural central Ghana using a mixed methods approach. *Front Public Health* 2020; **8**: 90. <https://doi.org/10.3389/fpubh.2020.00090>
- 288** Hackman HK, Annison L, Arhin RE et al. Self-medication with antibiotics during the COVID-19 pandemic: a cross-sectional study among adults in Tema, Ghana. *PLoS One* 2024; **19**: e0305602. <https://doi.org/10.1371/journal.pone.0305602>
- 289** Ngu RC, Fetei VF, Kika BT et al. Prevalence and determinants of antibiotic self-medication among adult patients with respiratory tract infections in the Mboppi Baptist Hospital, Douala, Cameroon: a cross-sectional study. *Diseases* 2018; **6**: 49. <https://doi.org/10.3390/diseases6020049>
- 290** Badger-Emeka LI, Emeka PM, Okosi M. Evaluation of the extent and reasons for increased non-prescription antibiotics use in a university town, Nsukka Nigeria. *Int J Health Sci* 2018; **12**: 11–7.
- 291** Torres NF, Solomon VP, Middleton LE. Identifying the commonly used antibiotics for self-medication in urban Mozambique: a qualitative study. *BMJ Open* 2020; **10**: e041323. <https://doi.org/10.1136/bmjopen-2020-041323>
- 292** Hammour KA, Jalil MA, Hammour WA. An exploration of parents’ knowledge, attitudes and practices towards the use of antibiotics in childhood upper respiratory tract infections in a tertiary Jordanian hospital. *Saudi Pharm J* 2018; **26**: 780–5. <https://doi.org/10.1016/j.jsps.2018.04.006>
- 293** Abdel-Qader DH, Albassam A, Ismael NS et al. Awareness of antibiotic use and resistance in Jordanian community. *J Prim Care Community Health* 2020; **11**: 2150132720961255. <https://doi.org/10.1177/2150132720961255>
- 294** Abdelmalek S, AlEjialet R, Rayyan WA et al. Changes in public knowledge and perceptions about antibiotic use and resistance in Jordan: a cross-sectional eight-year comparative study. *BMC Public Health* 2021; **21**: 750. <https://doi.org/10.1186/s12889-021-10723-x>
- 295** Alshakka M, Hatem NA, Al-Abd N et al. Knowledge, attitude, and practice toward antibiotic use among the general public in a resource-poor setting: a case of Aden-Yemen. *J Infect Dev Ctries* 2023; **17**: 345–52. <https://doi.org/10.3855/jidc.17319>
- 296** Elhaddadi H, Hamami A, Sara A et al. Prevalence and determinants of the use of antibiotics by self-medication in the pediatric population in Oujda, Morocco. *Cureus* 2024; **16**: e60126. <https://doi.org/10.7759/cureus.60126>
- 297** Burtcher D, Van den Bergh R et al. ‘They eat it like sweets’: a mixed methods study of antibiotic perceptions and their use among patients, prescribers and pharmacists in a district hospital in Kabul, Afghanistan. *PLoS One* 2021; **16**: e0260096. <https://doi.org/10.1371/journal.pone.0260096>
- 298** Khan FU, Mallhi TH, Khan FU et al. Evaluation of consumers’ perspective on the consumption of antibiotics, antibiotic resistance, and recommendations to improve the rational use of antibiotics: an exploratory qualitative study from post-conflicted region of Pakistan. *Front Pharmacol* 2022; **13**: 881243. <https://doi.org/10.3389/fphar.2022.881243>
- 299** Yusef D, Babaa AI, Bashairah AZ et al. Knowledge, practices and attitude toward antibiotics use and bacterial resistance in Jordan: a cross-sectional study. *Infect Dis Health* 2018; **23**: 33–40. <https://doi.org/10.1016/j.idh.2017.11.001>
- 300** Al-Taie A, Hussein AN, Albasry Z. A cross-sectional study of patients’ practices, knowledge and attitudes of antibiotics among Iraqi population. *J Infect Dev Ctries* 2021; **15**: 1845–53. <https://doi.org/10.3855/jidc.13066>

- 301** Atif M, Asghar S, Mushtaq I et al. What drives inappropriate use of antibiotics? A mixed methods study from Bahawalpur, Pakistan. *Infect Drug Resist* 2019; **12**: 687–99. <https://doi.org/10.2147/IDR.S189114>
- 302** Gillani AH, Chang J, Aslam F et al. Public knowledge, attitude, and practice regarding antibiotics use in Punjab, Pakistan: a cross-sectional study. *Expert Rev Anti Infect Ther* 2021; **19**: 399–411. <https://doi.org/10.1080/14787210.2021.1823216>
- 303** Mabrouk A B, Ammari F L, Werdani A et al. Parental self-medication with antibiotics in a Tunisian pediatric center. *Therapie* 2022; **77**: 477–85. <https://doi.org/10.1016/j.therap.2021.10.007>
- 304** Zahreddine L, Hallit S, Shakaroun S et al. Knowledge of pharmacists and parents towards antibiotic use in pediatrics: a cross-sectional study in Lebanon. *Pharm Pract* 2018; **16**: 1194. <https://doi.org/10.18549/PharmPract.2018.03.1194>
- 305** El Khoury G, Ramia E, Salameh P. Misconceptions and malpractices toward antibiotic use in childhood upper respiratory tract infections among a cohort of Lebanese parents. *Eval Health Prof* 2018; **41**: 493–511. <https://doi.org/10.1177/0163278716686809>
- 306** Ali HT, Barakat M, Abdelhalim AR et al. Unravelling the dilemma of self-medication in Egypt: a cross-sectional survey on knowledge, attitude, and practice of the general Egyptian population. *BMC Public Health* 2024; **24**: 652. <https://doi.org/10.1186/s12889-024-17913-3>
- 307** Akhund R, Jamshed F, Jaffry HA et al. Knowledge and attitude of general Pakistani population towards antibiotic resistance. *Cureus* 2019; **11**: e4266. <https://doi.org/10.7759/cureus.4266>
- 308** Muflih SM, Al-Azzam S, Karasneh RA et al. Public knowledge of antibiotics, self-medication, and household disposal practices in Jordan. *Expert Rev Anti Infect Ther* 2023; **21**: 477–87. <https://doi.org/10.1080/14787210.2023.2182770>
- 309** Hussain MA, Mohamed AO, Abdelkarim OA et al. Prevalence and predictors of antibiotic self-medication in Sudan: a descriptive cross-sectional study. *Antibiotics* 2023; **12**: 612. <https://doi.org/10.3390/antibiotics12030612>
- 310** Nazari J, Chezani-Sharahi N, Eshtrati B et al. Prevalence and determinants of self-medication consumption of antibiotics in children in Iran: a population-based cross-sectional study, 2018–19. *PLoS One* 2022; **17**: e0278843. <https://doi.org/10.1371/journal.pone.0278843>
- 311** Nepal A, Hendrie D, Robinson S et al. Knowledge, attitudes and practices relating to antibiotic use among community members of the Rupandehi district in Nepal. *BMC Public Health* 2019; **19**: 1558. <https://doi.org/10.1186/s12889-019-7924-5>
- 312** Turankar TB, Gaidhane SA, Gaidhane AM et al. Comparison of health care professionals' and laypeople's knowledge, attitudes, and practices on the use of antimicrobial and antimicrobial resistance. *J Educ Health Promot* 2023; **12**: 313. [https://doi.org/10.4103/jehp.jehp\\_39\\_23](https://doi.org/10.4103/jehp.jehp_39_23)
- 313** Chatterjee S, Hazra A, Chakraverty R et al. A multicentric knowledge-attitude-practice survey in the community about antimicrobial use and resistance in India. *Trans R Soc Trop Med Hyg* 2021; **115**: 785–91. <https://doi.org/10.1093/trstmh/traa124>
- 314** Karuniawati H, Hassali MAA, Suryawati S et al. Assessment of knowledge, attitude, and practice of antibiotic use among the population of Boyolali, Indonesia: a cross-sectional study. *Int J Environ Res Public Health* 2021; **18**: 8258. <https://doi.org/10.3390/ijerph18168258>
- 315** Yunita SL, Yang HW, Chen YC et al. Knowledge and practices related to antibiotic use among women in Malang, Indonesia. *Front Pharmacol* 2022; **13**: 1019303. <https://doi.org/10.3389/fphar.2022.1019303>
- 316** Azim MR, Iftakhar KMN, Rahman MM et al. Public knowledge, attitudes, and practices (KAP) regarding antibiotics use and antimicrobial resistance (AMR) in Bangladesh. *Heliyon* 2023; **9**: e21166. <https://doi.org/10.1016/j.heliyon.2023.e21166>
- 317** Miyano S, Htoon TT, Nozaki I et al. Public knowledge, practices, and awareness of antibiotics and antibiotic resistance in Myanmar: the first national mobile phone panel survey. *PLoS One* 2022; **17**: e0273380. <https://doi.org/10.1371/journal.pone.0273380>
- 318** Marasini S, Sharma S, Joshi A et al. Exploring knowledge, perceptions, and practices of antimicrobials, and their resistance among medicine dispensers and community members in Kavrepalanchok district of Nepal. *PLoS One* 2024; **19**: e0297282. <https://doi.org/10.1371/journal.pone.0297282>
- 319** Kotwani A, Joshi J, Lamkang AS et al. Knowledge and behavior of consumers towards the non-prescription purchase of antibiotics: an insight from a qualitative study from New Delhi, India. *Pharm Pract* 2021; **19**: 2206. <https://doi.org/10.18549/PharmPract.2021.1.2206>
- 320** Zawahir S, Lekamwasam S, Halvorsen KH et al. Self-medication behavior with antibiotics: a national cross-sectional survey in Sri Lanka. *Expert Rev Anti Infect Ther* 2021; **19**: 1341–52. <https://doi.org/10.1080/14787210.2021.1911647>
- 321** Rijal KR, Banjara MR, Dhungel B et al. Use of antimicrobials and antimicrobial resistance in Nepal: a nationwide survey. *Sci Rep* 2021; **11**: 11554. <https://doi.org/10.1038/s41598-021-90812-4>
- 322** Islam MW, Shahjahan M, Azad AK et al. Factors contributing to antibiotic misuse among parents of school-going children in Dhaka city, Bangladesh. *Sci Rep* 2024; **14**: 2318. <https://doi.org/10.1038/s41598-024-52313-y>
- 323** Sinuraya RK, Wulandari C, Amalia R et al. Understanding public knowledge and behavior regarding antibiotic use in Indonesia. *Infect Drug Resist* 2023; **16**: 6833–42. <https://doi.org/10.2147/IDR.S427337>
- 324** Saha A, Marma KKS, Rashid A et al. Risk factors associated with self-medication among the indigenous communities of Chittagong hill tracts, Bangladesh. *PLoS One* 2022; **17**: e0269622. <https://doi.org/10.1371/journal.pone.0269622>
- 325** Mannan A, Chakma K, Dewan G et al. Prevalence and determinants of antibiotics self-medication among indigenous people of Bangladesh: a cross-sectional study. *BMJ Open* 2024; **14**: e071504. <https://doi.org/10.1136/bmjopen-2022-071504>
- 326** Gautham M, Bhattacharyya S, Maity S et al. “Just as curry is needed to eat rice, antibiotics are needed to cure fever”—a qualitative study of individual, community and health system-level influences on community antibiotic practices in rural West Bengal, India. *BMJ Open* 2024; **14**: e076616. <https://doi.org/10.1136/bmjopen-2023-076616>
- 327** Wang J, Sheng Y, Ni J et al. Shanghai parents' perception and attitude towards the use of antibiotics on children: a cross-sectional study. *Infect Drug Resist* 2019; **12**: 3259–67. <https://doi.org/10.2147/IDR.S219287>
- 328** Wang Q, Wu Y, Wang D et al. The impacts of knowledge and attitude on behavior of antibiotic use for the common cold among the public and identifying the critical behavioral stage: based on an expanding KAP model. *BMC Public Health* 2023; **23**: 1683. <https://doi.org/10.1186/s12889-023-16595-7>
- 329** Lim JM, Chhoun P, Tuot S et al. Public knowledge, attitudes and practices surrounding antibiotic use and resistance in Cambodia. *JAC Antimicrob Resist* 2021; **3**: dlaa115. <https://doi.org/10.1093/jacamr/dlaa115>
- 330** Wong LP, Alias H, Husin SA et al. Factors influencing inappropriate use of antibiotics: findings from a nationwide survey of the general public in Malaysia. *PLoS One* 2021; **16**: e0258698. <https://doi.org/10.1371/journal.pone.0258698>
- 331** Ulaya G, Nguyen TCT, Vu BNT et al. Awareness of antibiotics and antibiotic resistance in a rural district of Ha Nam province, Vietnam: a cross-sectional survey. *Antibiotics* 2022; **11**: 1751. <https://doi.org/10.3390/antibiotics11121751>
- 332** Lin R, Duan L, Liu C et al. The public's antibiotic use behavioural patterns and their determinants for upper respiratory tract infections: a latent

- class analysis based on consumer behaviour model in China. *Front Public Health* 2023; **11**: 1231370. <https://doi.org/10.3389/fpubh.2023.1231370>
- 333** Chang CT, Lee M, Lee JCY et al. Public KAP towards COVID-19 and antibiotics resistance: a Malaysian survey of knowledge and awareness. *Int J Environ Res Public Health* 2021; **18**: 3964. <https://doi.org/10.3390/ijerph18083964>
- 334** Ha TV, Nguyen AMT, Nguyen HST. Public awareness about antibiotic use and resistance among residents in highland areas of Vietnam. *Biomed Res Int* 2019; **2019**: 9398536. <https://doi.org/10.1155/2019/9398536>
- 335** McKinn S, Trinh DH, Drabarek D et al. Drivers of antibiotic use in Vietnam: implications for designing community interventions. *BMJ Glob Health* 2021; **6**: e005875. <https://doi.org/10.1136/bmjgh-2021-005875>
- 336** Di KN, Tay ST, Ponnampalavanar SSS et al. Socio-demographic factors associated with antibiotics and antibiotic resistance knowledge and practices in Vietnam: a cross-sectional survey. *Antibiotics* 2022; **11**: 471. <https://doi.org/10.3390/antibiotics11040471>
- 337** Cheng J, Coope C, Chai J et al. Knowledge and behaviors in relation to antibiotic use among rural residents in Anhui, China. *Pharmacoepidemiol Drug Saf* 2018; **27**: 652–9. <https://doi.org/10.1002/pds.4429>
- 338** Diao M, Shen X, Cheng J et al. How patients' experiences of respiratory tract infections affect healthcare-seeking and antibiotic use: insights from a cross-sectional survey in rural Anhui, China. *BMJ Open* 2018; **8**: e019492. <https://doi.org/10.1136/bmjopen-2017-019492>
- 339** Choo SJ, Chang CT, Lee JCY et al. A cross-sectional study on public belief, knowledge and practice towards antibiotic use in the state of Perak, Malaysia. *J Infect Dev Ctries* 2018; **12**: 960–9. <https://doi.org/10.3855/jidc.10723>
- 340** Irawati L, Alrasheedy AA, Hassali MA et al. Low-income community knowledge, attitudes and perceptions regarding antibiotics and antibiotic resistance in Jelutong district, Penang, Malaysia: a qualitative study. *BMC Public Health* 2019; **19**: 1292. <https://doi.org/10.1186/s12889-019-7718-9>
- 341** Aslam A, Zin CS, Ab Rahman NS et al. Self-medication practices with antibiotics and associated factors among the public of Malaysia: a cross-sectional study. *Drug Healthc Patient Saf* 2021; **13**: 171–81. <https://doi.org/10.2147/DHPS.S331427>
- 342** Kong LS, Islahudin F, Muthupalaniappen L et al. Knowledge and expectations on antibiotic use among the general public in Malaysia: a nationwide cross-sectional survey. *Patient Prefer Adherence* 2021; **15**: 2405–16. <https://doi.org/10.2147/PPA.S328890>
- 343** Kounnavong S, Yan W, Sihavong A et al. Antibiotic knowledge, attitudes and reported practice during pregnancy and six months after birth: a follow-up study in Lao PDR. *BMC Pregnancy Childbirth* 2022; **22**: 701. <https://doi.org/10.1186/s12884-022-05018-x>
- 344** Sychareun V, Phounsavath P, Sihavong A et al. Perceptions and reported practices of pregnant women and mothers of children under two years of age regarding antibiotic use and resistance in Vientiane province, Lao PDR: a qualitative study. *BMC Pregnancy Childbirth* 2022; **22**: 569. <https://doi.org/10.1186/s12884-022-04894-7>
- 345** Thi Quynh Nhi L, de Alwis R, Khanh Lam P et al. Quantifying antimicrobial access and usage for paediatric diarrhoeal disease in an urban community setting in Asia. *J Antimicrob Chemother* 2018; **73**: 2546–54. <https://doi.org/10.1093/jac/dky231>
- 346** Cai HTN, Tran HT, Nguyen YHT et al. Challenges and lessons learned in the development of a participatory learning and action intervention to tackle antibiotic resistance: experiences from northern Vietnam. *Front Public Health* 2022; **10**: 822873. <https://doi.org/10.3389/fpubh.2022.822873>
- 347** Li P, Hayat K, Shi L et al. Knowledge, attitude, and practices of antibiotics and antibiotic resistance among Chinese pharmacy customers: a multicenter survey study. *Antibiotics* 2020; **9**: 184. <https://doi.org/10.3390/antibiotics9040184>
- 348** Zhu Y, Tang X, Yan R et al. Non-prescription antibiotic use for cough among Chinese children under 5 years of age: a community-based cross-sectional study. *BMJ Open* 2021; **11**: e051372. <https://doi.org/10.1136/bmjopen-2021-051372>
- 349** Duan L, Liu C, Wang D. The general population's inappropriate behaviors and misunderstanding of antibiotic use in China: a systematic review and meta-analysis. *Antibiotics* 2021; **10**: 497. <https://doi.org/10.3390/antibiotics10050497>
- 350** Lin L, Harbarth S, Hargreaves JR et al. Large-scale survey of parental antibiotic use for paediatric upper respiratory tract infections in China: implications for stewardship programmes and national policy. *Int J Antimicrob Agents* 2021; **57**: 106302. <https://doi.org/10.1016/j.ijantimicag.2021.106302>
- 351** Yin X, Mu K, Yang H et al. Prevalence of self-medication with antibiotics and its related factors among Chinese residents: a cross-sectional study. *Antimicrob Resist Infect Control* 2021; **10**: 89. <https://doi.org/10.1186/s13756-021-00954-3>
- 352** Zhou Z, Zhao D, Zhang H et al. Understanding parental self-medication with antibiotics among parents of different nationalities: a cross-sectional study. *Glob Health Res Policy* 2021; **6**: 42. <https://doi.org/10.1186/s41256-021-00226-y>
- 353** Yuan J, Du W, Li Z et al. Prevalence and risk factors of self-medication among the pediatric population in China: a national survey. *Front Public Health* 2021; **9**: 770709. <https://doi.org/10.3389/fpubh.2021.770709>
- 354** Nguyen TTP, Do TX, Nguyen HA et al. A national survey of dispensing practice and customer knowledge on antibiotic use in Vietnam and the implications. *Antibiotics* 2022; **11**: 1091. <https://doi.org/10.3390/antibiotics11081091>
- 355** Otioku E, Fenny AP, Labi AK et al. Knowledge, attitudes and practices regarding antimicrobial use and resistance among healthcare seekers in two tertiary hospitals in Ghana: a quasi-experimental study. *BMJ Open* 2023; **13**: e065233. <https://doi.org/10.1136/bmjopen-2022-065233>
- 356** Haenssge MJ, Xayavong T, Charoenboon N et al. The consequences of AMR education and awareness raising: outputs, outcomes, and behavioural impacts of an antibiotic-related educational activity in Lao PDR. *Antibiotics* 2018; **7**: 95. <https://doi.org/10.3390/antibiotics7040095>
- 357** Anstey Watkins J, Wagner F, Xavier Gómez-Olivé F et al. Rural South African community perceptions of antibiotic access and use: qualitative evidence from a health and demographic surveillance system site. *Am J Trop Med Hyg* 2019; **100**: 1378–90. <https://doi.org/10.4269/ajtmh.18-0171>
- 358** Davis M, Whittaker A, Lindgren M et al. Understanding media publics and the antimicrobial resistance crisis. *Glob Public Health* 2018; **13**: 1158–68. <https://doi.org/10.1080/17441692.2017.1336248>
- 359** Karvanen M, Cars O. The language of antimicrobial and antibiotic resistance is blocking global collective action. *Infect Dis (Lond)* 2024; **56**: 487–95. <https://doi.org/10.1080/23744235.2024.2332455>
- 360** King R, Hicks J, Rassi C et al. A process for developing a sustainable and scalable approach to community engagement: community dialogue approach for addressing the drivers of antibiotic resistance in Bangladesh. *BMC Public Health* 2020; **20**: 950. <https://doi.org/10.1186/s12889-020-09033-5>
- 361** Endale H, Mathewos M, Abdeta D. Potential causes of spread of antimicrobial resistance and preventive measures in one health perspective—a review. *Infect Drug Resist* 2023; **16**: 7515–45. <https://doi.org/10.2147/IDR.S428837>
- 362** Tornimbene B, Eremin S, Abednego R et al. Global antimicrobial resistance and use surveillance system on the African continent: early implementation 2017–2019. *Afr J Lab Med* 2022; **11**: 1594. <https://doi.org/10.4102/ajlm.v11i1.1594>
- 363** Batura N, Cuevas C, Khan M et al. How effective and cost-effective are behaviour change interventions in improving the prescription and use of antibiotics in low-income and middle-income countries? A protocol



- for a systematic review. *BMJ Open* 2018; **8**: e021517. <https://doi.org/10.1136/bmjopen-2018-021517>
- 364** Amin D, Garzón-Orjuela N, Garcia Pereira A *et al*. Artificial intelligence to improve antibiotic prescribing: a systematic review. *Antibiotics* 2023; **12**: 1293. <https://doi.org/10.3390/antibiotics12081293>
- 365** Pinto-de-Sá R, Sousa-Pinto B, Costa-de-Oliveira S. Brave new world of artificial intelligence: its use in antimicrobial stewardship—a systematic review. *Antibiotics* 2024; **13**: 307. <https://doi.org/10.3390/antibiotics13040307>
- 366** Yang J, Eyre DW, Lu L *et al*. Interpretable machine learning-based decision support for prediction of antibiotic resistance for complicated urinary tract infections. *NPJ Antimicrob Resist* 2023; **1**: 14. <https://doi.org/10.1038/s44259-023-00015-2>
- 367** Plechatá A, Makransky G, Böhm R. A randomized controlled trial investigating experiential virtual reality communication on prudent antibiotic use. *NPJ Digit Med* 2024; **7**: 244. <https://doi.org/10.1038/s41746-024-01240-3>
- 368** Rousham EK, Nahar P, Uddin MR *et al*. Gender and urban-rural influences on antibiotic purchasing and prescription use in retail drug shops: a one health study. *BMC Public Health* 2023; **23**: 229. <https://doi.org/10.1186/s12889-023-15155-3>
- 369** Schellack N, Strydom M, Pepper MS *et al*. Social media and COVID-19—perceptions and public deceptions of ivermectin, colchicine and hydroxychloroquine: lessons for future pandemics. *Antibiotics* 2022; **11**: 445. <https://doi.org/10.3390/antibiotics11040445>
- 370** Erku DA, Belachew SA, Abrha S *et al*. When fear and misinformation go viral: pharmacists' role in deterring medication misinformation during the 'infodemic' surrounding COVID-19. *Res Social Adm Pharm* 2021; **17**: 1954–63. <https://doi.org/10.1016/j.sapharm.2020.04.032>
- 371** Ghibu S, Juncan AM, Rus LL *et al*. The particularities of pharmaceutical care in improving public health service during the COVID-19 pandemic. *Int J Environ Res Public Health* 2021; **18**: 9776. <https://doi.org/10.3390/ijerph18189776>
- 372** Pantasri T. Expanded roles of community pharmacists in COVID-19: a scoping literature review. *J Am Pharm Assoc* 2022; **62**: 649–57. <https://doi.org/10.1016/j.japh.2021.12.013>
- 373** Saleem Z, Sono TM, Godman B. Concerns with current drug laws regarding the purchasing antibiotics without a prescription in Pakistan; ways forward to assist the national action plan. *Expert Rev Anti Infect Ther* 2023; **21**: 1163–5. <https://doi.org/10.1080/14787210.2023.2260096>
- 374** Kalungia A, Godman B. Implications of non-prescription antibiotic sales in China. *Lancet Infect Dis* 2019; **19**: 1272–3. [https://doi.org/10.1016/S1473-3099\(19\)30408-6](https://doi.org/10.1016/S1473-3099(19)30408-6)
- 375** Lorencatto F, Charani E, Sevdalis N *et al*. Driving sustainable change in antimicrobial prescribing practice: how can social and behavioural sciences help? *J Antimicrob Chemother* 2018; **73**: 2613–24. <https://doi.org/10.1093/jac/dky222>
- 376** Ogunleye OO, Basu D, Mueller D *et al*. Response to the novel corona virus (COVID-19) pandemic across Africa: successes, challenges, and implications for the future. *Front Pharmacol* 2020; **11**: 1205. <https://doi.org/10.3389/fphar.2020.01205>
- 377** Calac AJ, Haupt MR, Li Z *et al*. Spread of COVID-19 vaccine misinformation in the ninth inning: retrospective observational infodemic study. *JMIR Infodemiology* 2022; **2**: e33587. <https://doi.org/10.2196/33587>
- 378** Tsao SF, Chen H, Tisseverasinghe T *et al*. What social media told us in the time of COVID-19: a scoping review. *Lancet Digit Health* 2021; **3**: e175–94. [https://doi.org/10.1016/S2589-7500\(20\)30315-0](https://doi.org/10.1016/S2589-7500(20)30315-0)
- 379** Kesten JM, Bhattacharya A, Ashiru-Oredope D *et al*. The antibiotic guardian campaign: a qualitative evaluation of an online pledge-based system focused on making better use of antibiotics. *BMC Public Health* 2017; **18**: 5. <https://doi.org/10.1186/s12889-017-4552-9>
- 380** DHIS2. Homepage. <https://dhis2.org/>
- 381** Joseph AM, Fernandez V, Kritzman S *et al*. COVID-19 misinformation on social media: a scoping review. *Cureus* 2022; **14**: e24601. <https://doi.org/10.7759/cureus.24601>
- 382** Hernandez RG, Hagen L, Walker K *et al*. The COVID-19 vaccine social media infodemic: healthcare providers' missed dose in addressing misinformation and vaccine hesitancy. *Hum Vaccin Immunother* 2021; **17**: 2962–4. <https://doi.org/10.1080/21645515.2021.1912551>
- 383** Stewart R, Madonsela A, Tshabalala N *et al*. The importance of social media users' responses in tackling digital COVID-19 misinformation in Africa. *Digit Health* 2022; **8**: 20552076221085070. <https://doi.org/10.1177/20552076221085070>