



ORIGINAL ARTICLE

WILEY

External validation of a prediction model for estimating fat mass in Arab children and adolescents

Tareq Al-Ati PhD¹ | Khalid El Kari PhD² | Lara Nasreddine PhD³ |
Hessa Al-Kandari MRCPCH⁴ | Richard D. Riley PhD⁵ | Peter H. Whincup FRCP⁶ |
Christopher G. Owen PhD⁶ | Mohammed T. Hudda PhD⁴  

¹Food and Nutrition Program, Environment and Life Sciences Research Institute Center, Kuwait Institute for Scientific Research, Safat, Kuwait

²RDC-Nutrition, Centre National de l'Energie Des Sciences et Techniques Nucleaires, Ibn Tofail University, Kenitra, Morocco

³Department of Nutrition and Food Sciences, American University of Beirut, Beirut, Lebanon

⁴Department of Population Health, Dasman Diabetes Institute, Kuwait City, Kuwait

⁵School of Health Sciences, University of Birmingham, Birmingham, UK

⁶Population Health Research Institute, St George's School of Health & Medical Sciences, City St. George's University of London, London, UK

Correspondence

Mohammed T. Hudda, Department of Population Health, Dasman Diabetes Institute, Kuwait City, Kuwait.

Email: mohammed.hudda@dasmaninstitute.org

Funding information

Pfizer Inc. Global Obesity ASPIRE Research Grant, Grant/Award Numbers: Ref, 91629725; National Institute for Health and Care Research (NIHR) Birmingham Biomedical Research Centre

Abstract

Background/Aims: Current childhood fat mass (FM) assessment techniques are not suitable for clinical and population-level adiposity assessment. A prediction model, which accurately estimates childhood FM using predictor variables of weight, height, age, sex and ethnicity, requires validation in Arab populations. We evaluate the model's performance in Kuwaiti, Lebanese and Moroccan children/adolescents.

Methods: Data from three cross-sectional studies on 471 individuals, aged 6–15 years, were obtained with complete information on predictors and the outcome of log transformed fat-free mass assessed by reference standard deuterium dilution (lnFFM). Country-specific predictive performance statistics of R^2 , calibration slope and calibration-in-the-large (measures the calibration/agreement between observed and predicted lnFFM with ideal values of 1 and 0, respectively) and root mean square error (RMSE) were quantified and pooled across countries via random-effects meta-analysis. FM estimates from bioimpedance were also available for Lebanese children and were compared to the reference standard.

Results: The model showed strong predictive ability in all populations. Pooled R^2 calibration slope and calibration-in-the-large values on the original lnFFM scale were 87.73% (95% CI: 77.20, 98.26%), 0.95 (95% CI: 0.83, 1.08) and -0.03 (95% CI: -0.16 , 0.11), respectively. Model intercepts were recalibrated in each country to improve accuracy; updated country-specific equations are provided. After recalibration, RMSEs on the FM scale were 1.3, 1.6 and 2.8 kg in Kuwait, Lebanon and Morocco, respectively. The RMSE from the model was lower than bioimpedance (2.4 kg) amongst Lebanese children.

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial](https://creativecommons.org/licenses/by-nc/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

© 2025 The Author(s). *Diabetes, Obesity and Metabolism* published by John Wiley & Sons Ltd.

Interpretation: The model explained a large proportion of the variance in FM, produced well-calibrated predictions and relatively low RMSEs in Arab settings. It predicted FM more accurately than bioimpedance, indicating its potential for implementation in clinical- and population-level settings, particularly in low- and middle-income countries.

KEYWORDS

body composition, meta-analysis, observational study, real-world evidence

1 | INTRODUCTION

The global prevalence of paediatric overweight and obesity is increasing rapidly, becoming a major risk factor for numerous chronic diseases of public health importance.¹ Overweight and obesity are defined by the World Health Organization as abnormal or excessive fat accumulation that presents a risk to health.² Historically, in the absence of more direct techniques to assess (or estimate) body fat, overweight and obesity, proxy indirect markers of body fat were relied upon. The most widely used proxy marker of body fat is body mass index (BMI) which is calculated by dividing an individual's weight by the square of their height (weight/height²), and thresholded to define overweight and obesity.^{3,4} However, as BMI is a weight-based proxy marker of body fat, it is unable to discriminate between fat mass (FM) and fat-free mass (FFM), which can vary markedly in children and adolescents with a given BMI.⁵ Several more direct in vivo techniques now exist to assess body fat, such as magnetic resonance imaging (MRI) scanning, dual energy X-ray absorptiometry (DXA), bioelectric impedance (BIA), the four-component model and the deuterium dilution technique.³ However, these methods can be invasive (e.g. MRI), costly (e.g. MRI and deuterium dilution) and some have limited accuracy when used for individual-level assessment (e.g. DXA and BIA), which inhibits use for routine clinical and/or public health body fat assessment.^{3,6–18} Thus, BMI has remained the preferred marker of body fat in children and adolescents.

An alternative approach for assessing FM in children and adolescents, an equation that relies solely on weight, height, age, sex and ethnicity to accurately estimate FFM (and thus FM) amongst individuals aged 4–15 years, has recently been proposed. The equation was initially developed and validated within UK children and adolescents,^{12,19} and was found to produce highly accurate estimates of childhood FM. Crucially, when using reference-standard deuterium dilution as a benchmark, this model assessed FM more accurately than established techniques of BIA and DXA in UK children.¹² The UK-based model equation also underwent extensive external model validation in children and adolescents in 20 other settings outside the United Kingdom and demonstrated strong predictive performance in estimating FM across all settings.^{20,21}

Recent evidence has suggested that childhood overweight and obesity rates are exceptionally high across the Middle East and North Africa (MENA) region,^{22,23} highlighting the need for improved childhood adiposity assessment in order to aid efforts to tackle this public health crisis. Therefore, we performed an external validation of the

UK-based FM prediction model to assess its predictive performance amongst children and adolescents from three countries in the MENA region: Kuwait, Lebanon and Morocco, and (where required) aimed to produce country-specific recalibrated model equations.

2 | METHODS

2.1 | Data sources and study population

To externally validate and potentially recalibrate the UK-based prediction equation for the MENA region, data were obtained from three observational studies conducted in Kuwait,²⁴ Lebanon²⁵ and Morocco,²⁶ which each utilised the deuterium dilution method to assess body composition in children. Each study contributed data on deuterium assessed fat-free mass and fat mass, weight, height, sex and age for children aged between 4 and 15 years and in total included 476 children. The Lebanese study also provided data on body composition ascertained by the BIA technique (assessed using the Imp DF50 tetrapolar single frequency [200 mA at 50 kHz] electrical bioimpedance analyser).^{25,27} After removing implausible (negative) values of deuterium dilution assessed fat mass ($n = 5$), the analysis dataset was made up of 471 children (193 from Kuwait, 162 from Lebanon and 116 from Morocco).

2.2 | Outcome and predictor assessment

The outcome of the model being validated was natural log transformed FFM (lnFFM), ascertained using reference standard deuterium dilution.^{8,28} This technique was used as the outcome for the initial development of the UK-based prediction model and the previous external validations of the model.^{12,19,20} There were no missing data on the outcome (lnFFM) or the predictors (weight, height, sex, age or ethnicity).

2.3 | Statistical analysis

Statistical analyses were conducted in Stata (version 18.0). The Transparent Reporting of a multivariable model for Individual Prognosis Or Diagnosis (TRIPOD)²⁹ guideline for the reporting of studies validating a multivariable prediction model was followed.

2.4 | Evaluation of the performance of UK-based prediction model equation

The UK-based model equation¹⁹ was applied to all individuals within this external validation dataset, and predictions of InFFM were obtained using predictors of weight, height, sex, age and ethnicity. The country-specific predictive performance of the model equation was assessed by comparing InFFM values from the deuterium dilution technique with the predicted values of InFFM (obtained using the UK-based model). As recommended by the TRIPOD guidelines,²⁹ predictive performance was assessed using the following established predictive performance statistics:

1. R^2 —The proportion of the variance in InFFM that is explained by the model predicted InFFM within these childhood settings. The variance and 95% confidence interval (95% CI) for R^2 was estimated using the Wald-type method by Tan.³⁰
2. Calibration
 - a. Calibration Slope—Provides a quantification of the accuracy of the predictions of InFFM from the model across the range of InFFM values from deuterium dilution. The ideal slope value is 1.
 - b. Calibration-in-the-large (CITL)—Measurement of the overall agreement between average model predicted values of InFFM and average deuterium dilution assessed values of InFFM. The ideal value of CITL is 0.
 - c. Calibration plot with flexible calibration curve—Graph of deuterium dilution assessed InFFM plotted against model predicted values of InFFM using a smoothed curve.
3. Root mean square error (RMSE)—The average difference between model predicted InFFM values and the deuterium dilution assessed InFFM.

While the primary results on the model performance are presented in terms of InFFM, the outcome of the original UK-based model, we also present the performance in terms of FFM and FM.

To summarise the performance of the UK-based model across the MENA childhood settings and obtain estimates of the average performance and between-country heterogeneity, we pooled the country-specific predictive performance measures of R^2 Calibration Slope and CITL using a random effects meta-analysis fitted using restricted maximum likelihood estimation (REML)³¹ with the Hartung–Knapp³² approach used to derive confidence intervals. τ^2 , the estimate of between-study variance, was used to summarise heterogeneity for the three pooled performance measures.³³ Model performance was also assessed by sex.

2.5 | Country-specific recalibration of UK-based prediction model equation

Where systematic error in the prediction of InFFM from the UK-based model was observed across countries, we recalibrated the model's intercept term and provided updated country-specific model equations.

2.6 | Comparisons of FM from the prediction equation and bioelectrical impedance with estimates from deuterium dilution from Lebanon

We also compared the accuracy of FM predictions from the Hudda et al. UK-based equation with FM predictions from BIA, using the deuterium dilution method as a reference standard in Lebanese children for whom FM estimates were available from all three techniques. To do so, FM estimates from BIA were compared with FM from deuterium dilution in terms of the R^2 calibration (slope and CITL) and RMSE values. We then compared these predictive performance statistics to those obtained when comparing the UK-based equation to deuterium dilution (before and after recalibration).

3 | RESULTS

The key characteristics of included participants are summarised by country in Table 1. Almost half of the participants from Kuwait and Lebanon were girls, while most of the Moroccan participants (73%) were girls. The median age of individuals from Kuwait (8.2 years) and Lebanon (9.2 years) was lower than those from Morocco (13.9 years). These age differences were reflected in the differences across countries in average levels of height and deuterium dilution assessed fat-free mass (Table 1). However, while the average age of Kuwaiti children was lower than that of their counterparts from Lebanon, their average weight and deuterium dilution assessed fat mass were higher. Median FM amongst children from the original UK development population (8.4 kg)¹⁹ was similar to those from Lebanon (7.8 kg) but lower than that from Kuwait and Morocco in this external validation study.

3.1 | Assessment of UK-based model performance in MENA region countries

When applied to these external validation data, the UK-based model equation produced (in terms of InFFM) high R^2 values of 91.44% (95% CI: 89.13%–93.75%), 87.35% (95% CI: 83.71%–90.99%) and 82.76% (95% CI: 77.05%–88.47%) from Kuwait, Lebanon and Morocco, respectively (Table 2). When the country-specific R^2 values were pooled via random-effects meta-analysis (Figure 1), the overall R^2 value was 87.73% (95% CI: 77.20%–98.26%) with some evidence of between-country heterogeneity in the R^2 values ($\tau^2 = 13.59$). RMSE values were 0.10, 0.07 and 0.08 in terms of InFFM (Table 2). For inter-prebility, in terms of FM, the RMSE values from the original UK-based model were 2.2, 1.7 and 2.9 kg from Kuwait, Lebanon and Morocco, respectively. These average errors represent approximately 16.5%, 21.7% and 18.4% of the average deuterium dilution assessed FM in each of the three countries.

The UK-based model demonstrated high levels of calibration between observed and predicted InFFM in all three countries with calibration slopes of 1.01 (95% CI: 0.96–1.05), 0.91 (95% CI: 0.86–0.97) and 0.93 (95% CI: 0.85–1.01) from Kuwait, Lebanon and Morocco,

TABLE 1 Basic summary statistics of the analysis population, by country.

Variable	Kuwait (N = 193)	Lebanon (N = 162)	Morocco (N = 116)
Age (years)	8.2 (7.6, 8.7)	9.2 (8.6, 9.8)	13.9 (13.3, 14.4)
Height (m)	1.29 (1.25, 1.35)	1.34 (1.28, 1.40)	1.58 (1.51, 1.62)
Weight (kg)	32.83 (26.20, 41.60)	30.90 (26.05, 37.60)	54.00 (44.00, 63.25)
DD assessed fat mass (kg)	13.37 (8.17, 18.30)	7.82 (5.22, 11.87)	15.72 (9.91, 21.80)
DD assessed fat-free mass (kg)	20.49 (17.71, 23.43)	23.20 (20.52, 26.38)	36.11 (32.58, 41.58)
Boys (n, %)	92 (47.7)	77 (47.5)	32 (27.6)
Ethnic Group (n, %)			
White	0 (0.0)	0 (0.0)	0 (0.0)
Black	0 (0.0)	0 (0.0)	0 (0.0)
South Asian	0 (0.0)	0 (0.0)	0 (0.0)
Other Asian	0 (0.0)	0 (0.0)	0 (0.0)
Other	193 (100.0)	162 (100.0)	116 (100.0)

Note: Values are median (25th to 75th centile) unless stated otherwise.

Abbreviation: DD, deuterium dilution.

TABLE 2 External validation predictive performance statistics based on natural log transformed fat-free mass before and after recalibration of the intercept term, by country.

Variable	Kuwait (N = 193)	Lebanon (N = 162)	Morocco (N = 116)
Before recalibration			
R^2 (95% CI)	91.44 (89.13 to 93.75)	87.35 (83.71 to 90.99)	82.76 (77.05 to 88.47)
Calibration Slope (95% CI)	1.01 (0.96 to 1.05)	0.91 (0.86 to 0.97)	0.93 (0.85 to 1.01)
Calibration-in-the-large (95% CI)	−0.08 (−0.09 to −0.07)	0.02 (0.01 to 0.03)	−0.02 (−0.03 to −0.01)
RMSE	0.10	0.07	0.08
After recalibration			
R^2 (95% CI)	91.44 (89.13 to 93.75)	87.35 (83.71 to 90.99)	82.76 (77.05 to 88.47)
Calibration Slope (95% CI)	1.01 (0.96 to 1.05)	0.91 (0.86 to 0.97)	0.93 (0.85 to 1.01)
Calibration-in-the-large (95% CI)	0.00 (−0.01 to 0.01)	0.00 (−0.01 to 0.01)	0.00 (−0.01 to 0.01)
RMSE	0.06	0.07	0.07

Note: Hudda et al. is the fat-free mass prediction equation proposed in Hudda et al. BMJ 2019 doi: [10.1136/bmj.l4293](https://doi.org/10.1136/bmj.l4293), recalibration refers to recalibration of the intercept term.

Abbreviations: BIA, bioelectrical impedance analysis; CI, confidence interval; DD, deuterium dilution; RMSE, root mean square error of lnFFM.

respectively (Table 2). The pooled calibration slope (Figure 1) was 0.95 (95% CI: 0.83–1.08) with no evidence of between-country heterogeneity in the calibration slopes ($\tau^2 = 0$). While country-specific CITL values were close to the ideal value of 0, the associated 95% confidence intervals failed to contain the ideal value of 0 in any of the settings (Table 2). When pooled, the overall pooled CITL value was close to the ideal value of zero (pooled CITL = −0.03) (Figure 1) and there was no evidence of heterogeneity in the CITL values ($\tau^2 = 0$). The calibration plots demonstrated the good levels of calibration across the range of lnFFM (Figure 2) and FM values (Figure S1) within each of the three countries, including at the lower- and upper-ends of the distribution, with the flexible calibration curve close to the ideal 45 degree line of perfect calibration. However, as observed from the CITL results, the graphs demonstrated some systematic error in the prediction of lnFFM and FM within each of the settings, particularly amongst Kuwaiti children (Figure 2, Figure S1). As a result of this

observed systematic error in the UK-based model's prediction of lnFFM, the model intercept was recalibrated to provide updated country-specific equations (Box 1).

After recalibration, the CITL and RMSE values on the lnFFM scale were closer to the ideal values of 0 in all three settings, with all of the country-specific 95% CIs for the CITL now containing the ideal value (Table 2). Updated calibration plots after recalibration on the lnFFM scale are provided in Figure S2. We also provide the country-specific predictive performance statistics after recalibration of the intercept on the FM scale overall (Table 3, Figure S3) and by sex (Figure S4). The RMSE values from the recalibrated model equation in terms of FM were 1.28, 1.60 and 2.76 kg from Kuwait, Lebanon and Morocco, respectively (Table 3). On the FM scale, these average errors from the recalibrated prediction model represent 9.6%, 20.5% and 17.6% of average deuterium dilution assessed FM in Kuwait, Lebanon and Morocco, respectively.




FIGURE 1 Forest plot of R^2 values, calibration slope and calibration-in-the-large based on natural log transformed fat-free mass before recalibration of the intercept term, by country and overall. Study labels: 1 = Kuwait, 2 = Lebanon, 3 = Morocco. Overall estimates are from the random effect restricted maximum likelihood model with Hartung–Knapp standard errors.

3.2 | Comparisons of FM from the UK-based equation and bioelectrical impedance with estimates from deuterium dilution from Lebanon

Table 4 contains a comparison of the predictive performance statistics from BIA and the UK-based equation with estimates from the deuterium dilution reference standard for children from Lebanon. Figure S5 contains a calibration plot comparing FM estimates from BIA with those of deuterium dilution observed FM. When using deuterium

dilution assessed FM as the reference, FM predictions obtained from BIA were not as accurate as the predictions from the UK-based equation even before local recalibration. The R^2 value from BIA was lower than that of the UK-based equation (87.7% vs. 91.9%), and BIA produced higher values of the CITL (1.29 kg vs. -0.52 kg) and a higher RMSE (2.36 kg vs. 1.66 kg) (Table 4). This RMSE from BIA of 2.36 kg represents 30.2% of the average deuterium dilution assessed FM (7.82 kg) in Lebanese children. These differences in accuracy were more marked after the recalibration of the UK-based equation.




FIGURE 2 Calibration assessment of the model based on natural log transformed fat-free mass (lnFFM) before recalibration of the intercept term, by country. Each plot shows the agreement between individuals' predicted values of lnFFM from the model (x-axis) against the individuals' observed lnFFM values (y-axis). The dashed line represents the line of equality. The solid line is the calibration curve, which is a smooth fit through the individual data points, showing the average calibration across individuals (with lighter shading the 95% CI of the curve). Slope = calibration slope (ideal value 1); CITL = calibration-in-the-large (ideal value 0).

BOX 1 Recalibrated country-specific constant terms for the prediction of natural log transformed fat-free mass in children and adolescents.

Country	Constant term
Kuwait	2.7227667
Lebanon	2.8283926
Morocco	2.7863799

$$\begin{aligned} \text{Fat Mass} = & \text{Weight} - \exp(0.3073 * \text{height}^2 - 10.0155 * \text{weight}^{-1} \\ & + 0.004571 * \text{weight} + 0.01408 * \text{BA} - 0.06509 * \text{SA} \\ & - 0.02624 * \text{AO} - 0.01745 * \text{Other} - 0.9180 * \ln(\text{age}) \\ & + 0.6488 * \text{age}^{0.5} + 0.04723 * \text{male} \\ & + \text{Country specific Constant Term from above table}). \end{aligned}$$

Ln = natural logarithmic transformation, FFM = fat-free mass. BA, SA, AO and Other = 1 if child is of Black, South Asian, Other Asian or Other ethnic origins respectively and = 0 if not. If child is of unknown ethnic group, treat as of White ethnic origins. Height is measured in metres, weight in kilograms and age in years.

The UK-based model equation showed very strong predictive ability in all the three new settings, with large percentage of variation explained, and good overall calibration of observed (deuterium dilution) and model predicted FM values. The RMSE values (in terms of FM for interpretability) were 2.19 kg in Kuwaiti children, 1.66 kg in Lebanese children and 2.86 kg in Moroccan children. This demonstrates the generalisability of the model within the MENA region, where it had not previously been extensively externally validated. While the calibration slopes were close to the ideal of 1 for all three countries, there was evidence of a small systematic error in the predictions from the UK-based model. Therefore, the model intercept was recalibrated and updated country-specific prediction equations were provided to further improve country-specific accuracy. Following recalibration, and as expected, the model performance showed improvement in the CITL and RMSE values in all three settings. Finally, amongst Lebanese children for whom BIA-assessed FM was also available, the UK-based prediction model validated in this study, both before and after local recalibration of the intercept term, produced more accurate predictions of FM than BIA, from which the CITL was 1.29 kg and the RMSE was 2.36 kg.

4.2 | Comparison with other studies

The UK-based prediction model¹⁹ has previously been extensively externally validated in 19 settings outside the United Kingdom,^{20,21} and has shown to produce accurate predictions of FM in children aged 4–15 years across all settings. However, it was noted that further validation within settings from the MENA region was required, as Tunisia was the only country included in the external validation.²⁰ Amongst Tunisian children, the predictive performance of the UK-based equation prior to recalibration of the intercept was observed to be strong with good calibration of observed and model predicted values of

4 | DISCUSSION

4.1 | Principal findings

We externally validated, within three MENA countries, the predictive performance of an earlier proposed UK-based model which uses weight, height, sex, age and ethnicity to estimate childhood FM levels.

TABLE 3 External validation predictive performance statistics based on fat-free mass and fat mass after recalibration of the intercept term, by country.

Variable	Kuwait (N = 193)	Lebanon (N = 162)	Morocco (N = 116)
Fat-free Mass			
R ² (95% CI)	91.16 (88.78 to 93.54)	86.50 (82.64 to 90.37)	82.96 (77.31 to 88.61)
Calibration Slope (95% CI)	1.03 (0.99 to 1.08)	0.91 (0.85 to 0.96)	0.94 (0.87 to 1.02)
Calibration-in-the-large (95% CI)	0.04 (−0.14 to 0.23)	−0.02 (−0.27 to 0.23)	0.03 (−0.48 to 0.54)
RMSE	1.28	1.60	2.76
Fat Mass			
R ² (95% CI)	96.83 (95.96 to 97.71)	91.94 (89.56 to 94.32)	89.99 (86.53 to 93.44)
Calibration Slope (95% CI)	0.98 (0.95 to 1.00)	1.10 (1.05 to 1.15)	1.08 (1.01 to 1.15)
Calibration-in-the-large (95% CI)	−0.04 (−0.23 to 0.14)	0.02 (−0.23 to 0.27)	−0.03 (−0.54 to 0.48)
RMSE	1.28	1.60	2.76

Note: Recalibration refers to recalibration of the intercept term.

Abbreviation: RMSE, root mean square error.

TABLE 4 Comparison of the performance of fat mass estimates from bioelectrical impedance and the Hudda et al. equation with estimates from the deuterium dilution reference standard for children from Lebanon.

Variable	BIA versus DD	Hudda et al. versus DD (before recalibration)	Hudda et al. versus DD (after recalibration)
Fat Mass			
R ² (95% CI)	87.66 (84.09 to 91.23)	91.92 (89.54 to 94.31)	91.94 (89.56 to 94.32)
Calibration Slope (95% CI)	0.90 (0.85 to 0.95)	1.08 (1.03 to 1.13)	1.10 (1.05 to 1.15)
Calibration-in-the-large (95% CI)	1.29 (0.99 to 1.60)	−0.52 (−0.76 to −0.27)	0.02 (−0.23 to 0.27)
RMSE	2.36	1.66	1.60

Note: Hudda et al. is the fat-free mass prediction equation proposed in Hudda et al. BMJ 2019 doi: [10.1136/bmj.l4293](https://doi.org/10.1136/bmj.l4293), recalibration refers to recalibration of the intercept term.

Abbreviations: BIA, bioelectrical impedance analysis; CI, confidence interval; DD, deuterium dilution; RMSE, root mean square error of FM.

InFFM, a high R² value of 81.0% and an RMSE value of 0.08.²⁰ These performance statistics from Tunisia are very similar to those observed when assessing the performance of the UK-based equation in Lebanese and Moroccan children in the current study. Moreover, the updated constant term for Tunisian children of 2.7858²⁰ is remarkably similar to that of Moroccan children found in this study of 2.7864.

While several other studies have developed prediction models to estimate FM in childhood populations outside of the United Kingdom,^{21,26,34–39} as discussed previously,²⁰ direct comparisons of the performance of those models with the model being validated in this study are difficult for several reasons such as models adopting different outcome variables (e.g., FM%), most models contain additional measurements (e.g., skinfold thickness, waist circumference, bioimpedance) as predictors rather than being based on readily available anthropometric and demographic predictors, studies not reporting the calibration and/or the RMSE statistics of their models. To our knowledge, just one published equation exists from the MENA region which uses only readily available information on weight, height, sex and age.²⁶ This study developed a model equation in Moroccan children and adults to estimate FM (via total body water estimation) and validated the developed equation in adults from Tunisia. This previous study did not report the

performance of the developed equation in children separately from adults and therefore is not suitable for comparison with the UK-based childhood equation being validated in this study.

Furthermore, results presented in this study demonstrate the superior performance of the UK-based equation compared with BIA at predicting FM amongst Lebanese children (using deuterium dilution as the reference standard) were consistent with those of earlier studies which made the same comparisons amongst UK children.^{7,12} Just as we found in this study, we previously also observed that when using BIA to assess FM, while R² values were high, there was evidence of miscalibration between BIA-produced FM predictions and deuterium assessed FM. Crucially, we demonstrated in our earlier study that RMSE values were lower from the UK-based model (2.6 kg) than from BIA (3.1 kg) and DXA (3.4 kg) when deuterium dilution was used as the reference standard.¹²

4.3 | Strengths and limitations

The current study has several strengths in validating the previously developed UK-based model equation.¹⁹ Most importantly, the same reference standard technique which was used to assess body fat in

the UK model development study, deuterium dilution, was used as the reference standard in each of the three countries being used in this validation study. This technique is known to provide accurate, safe and minimally invasive measurements of TBW (and FFM) with very low error.^{8,28} Furthermore, the sample size of each of the country-specific databases was reasonable to assess the generalisability of the UK-based model in each of the settings. There were also a few limitations. Firstly, while there were a reasonable number of individuals from each of the three settings, there was a fairly narrow age range available, with the youngest children being 8 years of age, and thus, the generalisability of the model to younger children from these settings could not be ascertained. However, as the model performed very well across the range of FM values included, it is likely the model will also provide accurate predictions of FM for children aged 5–7 years of age for whom FM values will on average be lower than their older counterparts included here. Finally, BIA assessments were only available from the Lebanese study, and therefore we could not make comparisons between the UK-based model equation and BIA across all three settings.

4.4 | Implications for clinicians and policymakers

The World Health Organization defines obesity by 'excessive fat deposits that can impair health', which can influence an individual's quality of life,⁴⁰ and therefore the availability of simple, accurate methods for FM assessment would represent an important advance in the field of adiposity. All available techniques to assess fat mass (and thus excessive or insufficient levels) *in vivo* are based upon a predictive model equation which in most cases is embedded within apparatus (such as DXA, MRI or BIA). While some of these techniques are being utilised in research settings, they are not being utilised at the clinical- or population-level largely due to either their accuracy, the time and/or cost associated with implementing these procedures in practice. However, a new predictive model was recently developed and extensively validated which, as it only requires information on childhood age, sex, weight, height and ethnicity, can easily be integrated into clinical- and population-level practice with no additional monetary or time burdens. In MENA region settings where electronic medical systems are in routine use at the clinical- and population-level, the validated model equation can be programmed into existing electronic software to automatically generate the estimated FM values for children where information on weight, height, sex and age are available. In settings where electronic medical systems are still not routinely utilised, then a Microsoft Excel based calculator has been developed (Data S2) to quickly and easily estimate FM based on the inserted values of weight, height, sex and age. The findings of this external validation study emphasise the strong predictive ability of the developed UK-based model in three new childhood settings from the MENA region across a range of FM values and ages. While the country-specific RMSE values from the model, particularly when presented as a proportion of the average deuterium dilution assessed FM

levels, appear to be considerable, this magnitude of error is considerable lower than that produced from other widely accepted and utilised body composition assessment methods of BIA (as shown amongst Lebanese children in this study and UK children in our earlier study¹²) and DXA (in our earlier study¹²). Therefore, the consistently high performance of the model at predicting FM across more than 20 settings further strengthens conclusions that, particularly after local-level recalibration, the model can be used to accurately assess body composition at the individual-level, where it outperforms established current methods of BIA and DXA. At the population-level, where given the extremely high reported prevalence of obesity, its related complications and consequences within the MENA region, this model can play a valuable role in providing improved FM assessment for epidemiological and public health obesity research and policy. For example, the model can provide accurate childhood FM assessment in order to (i) quantify associations between childhood body composition and short- and long-term disease risk, (ii) assess geographical, ethnic, socio-economic and temporal variations in FM within surveillance and monitoring initiatives from MENA region settings and (iii) assess the efficacy and effectiveness of novel interventions for the prevention of childhood obesity using FM (predicted from the model) as the outcome measure of interest as opposed to weight-based proxy measures.

4.5 | Further research

Further external validation in other MENA region settings would add to the model's generalisability, particularly including Arab children under the age of 8 years where deuterium dilution reference data are currently lacking and settings within the region of different socio-economic groups and/or those with different dietary patterns. Furthermore, to reduce RMSE further and thus improve the precision of individual-level predictions, additional predictors could be identified that add predictive value over those included in our models; ideally, these should also be readily available or easily measured variables to ensure the model retains easy implementation. Comparisons between DD, BIA and the prediction model in other MENA settings in the future would also be of value to further demonstrate the superiority of the model over BIA for total body FM assessment. Additionally, the development of sex- and age-specific FM reference values, based upon prospectively associated disease risks, would allow individuals to be classified into groups based on future disease risk attributable to their current FM levels, as opposed to current centile-based approaches to classify FM.⁴¹

AUTHOR CONTRIBUTIONS

Study design—Hudda MT, Al-Ati T. Data contributors—Al-Ati T, El Kari K, Nasreddine L. Data analysis—Hudda MT. Data interpretation—All authors. Drafting manuscript—Hudda MT. Critical evaluation and revision of manuscript—All authors.

FUNDING INFORMATION

MTH is supported by a Research Grant from Pfizer Inc. (Global Obesity ASPIRE Research Grant Ref: 91629725). RDR is supported by the National Institute for Health and Care Research (NIHR) Birmingham Biomedical Research Centre at the University Hospitals Birmingham NHS Foundation Trust and the University of Birmingham. RDR is an NIHR Senior Investigator. The views expressed are those of the author(s) and not necessarily those of the funders. This publication is the work of the authors who will serve as guarantors for the contents of this paper. The views expressed in this paper are those of the authors and not necessarily those of the funding agencies. Ethical approval was not required for this study based on secondary data analysis.

CONFLICT OF INTEREST STATEMENT

The authors confirm we have no competing interests.

PEER REVIEW

The peer review history for this article is available at <https://www.webofscience.com/api/gateway/wos/peer-review/10.1111/dom.16281>.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are from three independent research studies, each with their own ethical approvals and data sharing legislations. The PIs of each of the three studies [Kuwait: T Al-Ati, Morocco: K El Kari, Lebanon: L Nasreddine] should be contacted for data requests via the corresponding author.

ORCID

Mohammed T. Hudda  <https://orcid.org/0000-0001-7894-1159>

TWITTER

Mohammed T. Hudda  [mohammedhudda](https://twitter.com/mohammedhudda)

REFERENCES

- Gao L, Peng W, Xue H, et al. Spatial-temporal trends in global childhood overweight and obesity from 1975 to 2030: a weight mean center and projection analysis of 191 countries. *Global Health*. 2023; 19(1):53.
- World Health Organisation. Obesity. <https://www.who.int/health-topics/obesity>. Accessed 18/04/2024.
- Wells JC, Fewtrell MS. Measuring body composition. *Arch Dis Child*. 2006;91(7):612-617.
- Hall DM, Cole TJ. What use is the BMI? *Arch Dis Child*. 2006;91(4):283-286.
- Wells JC. A Hattori chart analysis of body mass index in infants and children. *Int J Obes Relat Metab Disord*. 2000;24(3):325-329.
- Prentice AM, Jebb SA. Beyond body mass index. *Obes Rev*. 2001;2(3):141-147.
- Reilly JJ, Gerasimidis K, Paparacleous N, et al. Validation of dual-energy x-ray absorptiometry and foot-foot impedance against deuterium dilution measures of fatness in children. *Int J Pediatr Obes*. 2010; 5(1):111-115.
- Wells JC, Fuller NJ, Dewit O, Fewtrell MS, Elia M, Cole TJ. Four-component model of body composition in children: density and hydration of fat-free mass and comparison with simpler models. *Am J Clin Nutr*. 1999;69(5):904-912.
- Williams JE, Wells JC, Wilson CM, Haroun D, Lucas A, Fewtrell MS. Evaluation of lunar prodigy dual-energy X-ray absorptiometry for assessing body composition in healthy persons and patients by comparison with the criterion 4-component model. *Am J Clin Nutr*. 2006; 83(5):1047-1054.
- Sinaiko AR, Donahue RP, Jacobs DR, Prineas RJ. Relation of weight and rate of increase in weight during childhood and adolescence to body size, blood pressure, fasting insulin, and lipids in young adults. *Circulation*. 1999;99(11):1471-1476.
- Wong WW, Hergenroeder AC, Stuff JE, Butte NF, Smith EO, Ellis KJ. Evaluating body fat in girls and female adolescents: advantages and disadvantages of dual-energy X-ray absorptiometry. *Am J Clin Nutr*. 2002;76(2):384-389.
- Hudda MT, Owen CG, Rudnicka AR, Cook DG, Whincup PH, Nightingale CM. Quantifying childhood fat mass: comparison of a novel height-and-weight-based prediction approach with DXA and bioelectrical impedance. *Int J Obes (Lond)*. 2020;45:99-103.
- Shypailo RJ, Butte NF, Ellis KJ. DXA: can it be used as a criterion reference for body fat measurements in children? *Obesity (Silver Spring)*. 2008;16(2):457-462.
- González-Ruiz K, Medrano M, Correa-Bautista JE, et al. Comparison of bioelectrical impedance analysis, slaughter skinfold-thickness equations, and dual-energy X-ray absorptiometry for estimating body fat percentage in Colombian children and adolescents with excess of adiposity. *Nutrients*. 2018;10(8):1086.
- Martin-Matillas M, Mora-Gonzalez J, Migueles JH, Ubago-Guisado E, Gracia-Marco L, Ortega FB. Validity of slaughter equations and bioelectrical impedance against dual-energy X-ray absorptiometry in children. *Obesity (Silver Spring)*. 2020;28(4):803-812.
- Nightingale CM, Rudnicka AR, Owen CG, et al. Are ethnic and gender specific equations needed to derive fat free mass from bioelectrical impedance in children of south asian, black african-Caribbean and white European origin? Results of the assessment of body composition in children study. *PLoS One*. 2013;8(10):e76426.
- Tinsley GM. Five-component model validation of reference, laboratory and field methods of body composition assessment. *Br J Nutr*. 2020;125(11):1246-1259.
- Khan S, Xanthakos SA, Hornung L, Arce-Clachar C, Siegel R, Kalkwarf HJ. Relative accuracy of bioelectrical impedance analysis for assessing body composition in children with severe obesity. *J Pediatr Gastroenterol Nutr*. 2020;70(6):e129-e135.
- Hudda MT, Fewtrell MS, Haroun D, et al. Development and validation of a prediction model for fat mass in children and adolescents: meta-analysis using individual participant data. *BMJ*. 2019; 366:l4293.
- Hudda MT, Wells JCK, Adair LS, et al. External validation of a prediction model for estimating fat mass in children and adolescents in 19 countries: individual participant data meta-analysis. *BMJ*. 2022; 378:e071185.
- Choy CC, Johnson W, Duckham RL, et al. Prediction of fat mass from anthropometry at ages 7 to 9 years in Samoans: a cross-sectional study in the Ola Tuputupua'e cohort. *Eur J Clin Nutr*. 2023;77(4):495-502.
- Farrag NS, Cheskin LJ, Farag MK. A systematic review of childhood obesity in the Middle East and North Africa (MENA) region: prevalence and risk factors meta-analysis. *Adv Pediatr Res*. 2017;4:8.
- Ministry of Health. Kuwait nutrition surveillance system - annual report. 2018.
- Al-Ati T, Preston T, Al-Hooti S, et al. Total body water measurement using the 2H dilution technique for the assessment of body composition of Kuwaiti children. *Public Health Nutr*. 2015;18(2):259-263.
- Nasreddine L, Naja F, Hills AP, Youssef SK, Chahine J, Hwalla N. Validity of predictive equations developed to estimate body fat from anthropometry and bioelectrical impedance analysis in 8-10 year-old children. *Clin Nutr*. 2012;31(3):364-371.

26. El Kari K, Mankai A, El Kouki D, et al. Anthropometry-based prediction equation of body composition in a population aged 12-88 years. *J Nutr*. 2023;153(3):657-664.
27. Liu A, Byrne NM, Ma G, et al. Validation of bioelectrical impedance analysis for total body water assessment against the deuterium dilution technique in Asian children. *Eur J Clin Nutr*. 2011;65(12):1321-1327.
28. Deurenberg P, Yap M. The assessment of obesity: methods for measuring body fat and global prevalence of obesity. *Baillieres Best Pract Res Clin Endocrinol Metab*. 1999;13(1):1-11.
29. Moons KM, Altman DG, Reitsma JB, et al. Transparent reporting of a multivariable prediction model for individual prognosis or diagnosis (tripod): explanation and elaboration. *Ann Intern Med*. 2015;162(1):W1-W73.
30. Tan LJ. Confidence Intervals for Comparison of the Squared Multiple Correlation Coefficients of Non-nested Models. 2012. Electronic Thesis and Dissertation Repository., The University of Western Ontario.
31. Viechtbauer W. Bias and efficiency of meta-analytic variance estimators in the random-effects model. *J Educ Behav Stat*. 2005;30(3):261-293.
32. Hartung J, Knapp G. On tests of the overall treatment effect in meta-analysis with normally distributed responses. *Stat Med*. 2001;20(12):1771-1782.
33. Riley RD, Higgins JPT, Deeks JJ. Interpretation of random effects meta-analyses. *BMJ*. 2011;342:d549.
34. Dugas LR, Cao G, Luke AH, Durazo-Arvizu RA. Adiposity is not equal in a multi-race/ethnic adolescent population: NHANES 1999-2004. *Obesity (Silver Spring)*. 2011;19(10):2099-2101.
35. Stevens J, Cai J, Truesdale KP, Cuttler L, Robinson TN, Roberts AL. Percent body fat prediction equations for 8- to 17-year-old American children. *Pediatr Obes*. 2014;9(4):260-271.
36. Stevens J, Ou FS, Cai J, Heymsfield SB, Truesdale KP. Prediction of percent body fat measurements in Americans 8 years and older. *Int J Obes (Lond)*. 2015;40:587.
37. Stevens J, Truesdale KP, Cai J, Ou FS, Reynolds KR, Heymsfield SB. Nationally representative equations that include resistance and reactance for the prediction of percent body fat in Americans. *Int J Obes (Lond)*. 2017;41(11):1669-1675.
38. Truesdale KP, Roberts A, Cai J, Berge JM, Stevens J. Comparison of eight equations that predict percent body fat using skinfolds in American youth. *Childhood Obes (Print)*. 2016;12(4):314-323.
39. Lohman TG, Caballero B, Himes JH, et al. Estimation of body fat from anthropometry and bioelectrical impedance in native American children. *Int J Obes Relat Metab Disord*. 2000;24(8):982-988.
40. World Health Organisation. Obesity and overweight. Fact sheets. 2022. <https://www.who.int/en/news-room/fact-sheets/detail/obesity-and-overweight>. Accessed 29/08/2024, 2024.
41. McCarthy HD, Cole TJ, Fry T, Jebb SA, Prentice AM. Body fat reference curves for children. *Int J Obes (Lond)*. 2006;30(4):598-602.

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Al-Ati T, El Kari K, Nasreddine L, et al. External validation of a prediction model for estimating fat mass in Arab children and adolescents. *Diabetes Obes Metab*. 2025;1-10. doi:[10.1111/dom.16281](https://doi.org/10.1111/dom.16281)