



Suppl. Figure 1: Nucleoprotein serum IgG and saliva IgA



Suppl. Figure 2: Spike protein serum IgG before and after first and second immunisation



Suppl. Table 1:

No COVID		n	Median	lower quartile - upper quartile		Difference in medians (p)
Spike serum IgG	T0 (a)	25	2.00	0.10 - 41.90		
	T1 (b)	40	258.51	61.46 - 440.67	a/b	0.000
	T2 (c)	34	150.92	72.58 - 349.60	b/c	0.265
	T3 (d)	61	2473.03	1197.02 - 5054.24	c/d	0.000
	T4 (e)	60	1171.74	328.2 - 2445.34	d/e	0.000
	T5 (f)	59	399.38	207.8 - 1207.45	e/f	0.000
					b/d	0.000

COVID		n	Median	lower quartile - upper quartile		Difference in medians (p)
Spike serum IgG	T0 (a)	2	omit	omit		
	T1 (b)	19	4186.62	1207.1 - 9585.57	a/b	
	T2 (c)	21	2807.47	1211.51 - 4952.68	b/c	0.109
	T3 (d)	34	7094.07	3802.29 - 14858.93	c/d	0.022
	T4 (e)	21	2105.83	1275.15 - 3532.52	d/e	0.001
	T5 (f)	21	1077.50	711.40 - 1510.21	e/f	0.001
					b/d	0.109

Suppl. Figure 3: Spike protein serum IgA before and after first and second immunisation

