



HYPERTENSION
 • ↓ prevalence in women than men (24% vs. 27%)
 • Systolic BP ↓ in women than men until 60 y

DASH
 49%

EnligHTN III 38%
 RADIANCE-HTN
 SOLO RADIANCE II 29%
 TARGET BP I 26%

SURGICAL REVASC
 Women referred for CABG at an older age than men

ROMA
 Women (ongoing)
 100%

CV IMAGING
 ↑ non-obstructive CAD in women
 ↑ women refuse to undergo invasive assessment

PROMISE 53%
 CONFIRM 47%
 EVAREST 45%
 SCOT-HEART 44%
 OPT-PACE 40%

ISCHEMIA
 22%

Pregnant women usually excluded from RCTs

HVD
 Females 43% of TAVI patients (2022/2023)

TAVI RCTs
 almost 50%

HF
 ↑ HFpEF prevalence in women than men
 Prevalence ↑ with age

PARAGON-HF
 52%

HF RCTs (2001-2016)
 27%

Rehabilitation studies
 30% (15% in CAD RCTs)

ANOCA/INOCA
 SCAD causes most MI among pregnant women
 ↑ non-obstructive CAD among women (50-70%) compared to men

WISE
 100%

DISCO registry
 88.9%

CorMicA
 73.5%

SENIOR-RITA
 45%

ACS RCTs ('90s-2000)
 25%

OBSTRUCTIVE CAD
 Women older and ↑ comorbidities than men
 ↓ likely to be referred for diagnostic assessment