


RESEARCH ARTICLE OPEN ACCESS

Disparities in Stillbirths in England: Analysis of A Population-Based Study of 1.3 Million Births

Ggenga Kayode¹ | Baskaran Thilaganathan^{2,3} | Christy Burden¹ | Amy Howell¹ | Vincent Cheng¹ | Jane Sandall⁴ | Maria Viner⁵ | Lia Brigante⁶ | Dilly Anumba⁷ | Cathy Winter⁸ | Birte Harlev-Lam⁶ | Timothy Draycott^{1,2,8} | Andrew Judge¹ | Erik Lenguerrand¹  | Tommy's National Centre for Maternity Improvement

¹Translational Health Science, Bristol Medical School, Southmead Hospital, University of Bristol, Bristol, UK | ²Royal College of Obstetricians and Gynaecologists, London, UK | ³St. George's University Hospitals, London, UK | ⁴Department of Women and Children's Health, Faculty of Life Sciences & Medicine, King's College London, London, UK | ⁵Mothers for Mothers, Bristol, UK | ⁶Royal College of Midwives, London, UK | ⁷Academic Unit of Reproductive and Developmental Medicine-Obstetrics and Gynaecology, Faculty of Medicine Dentistry and Health, The University of Sheffield, Sheffield, UK | ⁸The PROMPT Maternity Foundation, Department of Women's Health, Southmead Hospital, Bristol, UK

Correspondence: E. Lenguerrand (erik.lenguerrand@bristol.ac.uk)

Received: 17 April 2024 | **Revised:** 27 February 2025 | **Accepted:** 9 March 2025

Funding: This work was supported by Tommy's Baby Charity.

Keywords: ethnicity | maternity care provision | social inequality | stillbirth

ABSTRACT

Objective: To examine the variation in stillbirth rates between different ethnic and socioeconomic groups within each organisational hospital group (health trust).

Design: National registry study.

Setting: All health trusts (HT) in National Health Service England.

Population: All mothers and babies born between April 2015 and March 2017.

Methods: This observational study examined ethnic and socioeconomic disparities in stillbirth rates for 1 268 367 births in 133 HTs compared to the national average.

Outcome: Stillbirth at or after 24 gestational weeks.

Results: The average stillbirth rates ranged from 3.4/1000 births for White women up to 7.1/1000 births for Black women. The rates ranged from 2.9/1000 births for women living in the least deprived areas to 4.7/1000 births for those in the most deprived. The proportions of HTs with stillbirth rates well above the national average (more than 2 standard deviations) for White, Asian and Black women were 0.8%, 21.8% and 38.6%, respectively. When HTs were ranked by stillbirth rate, there were notable variations, with some trusts demonstrating lower than average stillbirth rates for White women while concurrently having higher than average stillbirth rates for Asian and/or Black women. There were no units exhibiting lower than national average stillbirth rates for Asian/Black women while concurrently having higher than average stillbirth rates for White women.

Conclusions: These findings suggest that access to and delivery of maternity care vary depending on the mother's ethnicity and level of socioeconomic deprivation. Social factors are likely determinants of inequality in stillbirth rather than maternity care alone.

Abbreviations: IMD, index of multiple deprivation; LSOA, lower layer super output area; MIS, maternity information systems; NHS, National Health Service; NMPA, National Maternity and Perinatal Audit; SD, standard deviation; UK, United Kingdom.

G Kayode, and B Thilaganathan contributed equally to this work as first author; A Judge and E Lenguerrand contributed equally to this work as last author.

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2025 The Author(s). BJOG: An International Journal of Obstetrics and Gynaecology published by John Wiley & Sons Ltd.

1 | Introduction

Stillbirth has a devastating as well as protracted psychosocial and economic impact on families and society at large, making prevention of stillbirth a major global ambition [1–3]. Ethnicity, migration and socio-economic status of women are strong determinants of adverse pregnancy outcomes and drivers of health inequity [4–10]. Despite this, there is a paucity of published data on disparities in stillbirth rates across health trusts based on maternal ethnicity and socio-economic status.

Recent, national surveillance datasets within the United Kingdom (UK) show substantial variation in stillbirth rates, with stillbirth rates twice as high in Black women compared to White women, as well as variation in maternity units across the country [9, 11–13]. However, research into ethnic inequalities in stillbirth is limited in the UK. Current reports and published studies have focused mainly on descriptive analysis of geographical disparities in stillbirth and perinatal mortality without investigating the relationship to markers of health inequity and the possible impact of access to perinatal care [14, 15]. In particular, the extent of institutional bias on the associations between ethnicity and stillbirth has been suggested, but not systematically evaluated.

The aim of this study is to examine variation in stillbirth rates across different ethnic groups within organisational hospital groups (health trusts) and estimate the extent to which maternal ethnicity and socio-economic status influence disparities. Such findings may provide vital information to help stakeholders offer more tailored and targeted services to reduce the impact of health inequity on stillbirth rates.

2 | Methods

2.1 | Study Population

This observational study used data from mothers and babies born in England from April 1, 2015, to March 31, 2017, captured on hospital maternity information systems (MIS)—a subset of the National Maternity and Perinatal Audit (NMPA) [16] data related to National Health Service (NHS) England maternity units with approval from the Healthcare Quality Improvement Partnership (DARS-NIC-430380-F7L4Z-v0.4 HQIP348). During the study period, 234 maternity units from 133 health trusts submitted specific maternity information in England. The MIS datasets cover about 97% of all total births in England, and the data are of high quality [17, 18]. Publicly available information describing the lower layer super output area (LSOA) were linked to the anonymised MIS dataset to provide information on the index of multiple deprivation (IMD) of maternal residential areas [19].

2.2 | Outcome and Exposure

Stillbirth was defined as a baby born at or after 24 weeks gestational age with no sign of life. Maternal ethnicity was reported as recorded by healthcare providers: The NHS uses a standardised list of 16 categories to determine a patient's ethnicity. The NHS recommends that organisations ask patients about

their ethnicity using self-reporting, along with questions about national identity and religion. We classified ethnic groups as Asian, Black, mixed ethnicity, other and White [20, 21].

Women included in this analysis were categorised into 5 IMD groups; an aggregated index of socio-economic deprivation of the maternal residential area was used as a proxy for socio-economic status [19]. In England, deprivation is measured in small geographical areas known as LSOA. LSOAs are defined as geographical areas of a similar population size, with an average of 1500 residents. As a measure of socioeconomic deprivation, we used the IMD score, a relative measure of deprivation based on LSOAs. Publicly available information describing the LSOA, produced by the Office of National Statistics, was linked to the anonymised MIS dataset to provide information on the IMD of maternal residential areas [19]. The IMD is the most used measure of deprivation within small areas in England. The seven domains used to generate deprivation scores include income, employment, education, health, crime, barriers to housing and services, and living environment. We categorised IMD into five groups (quintiles), with 1 being the most deprived and 5 denoting the least deprived group.

2.3 | Statistical Analysis

Participants' characteristics were reported as frequencies and percentages (%). Disparities in stillbirth rates were calculated for both individual maternity units and their amalgamated health trusts. Standard deviations [SD] were used to visualise stillbirth rate variation between health trusts as they are commonly in national audits [22]. The national average of stillbirth and the SD were determined across all Trusts included in the analysis. Using the national average of stillbirth and corresponding SD, Trusts were classified into five categories based on their stillbirth rates (Figure S1) Well below average ($< -2SD$ below the national average, Green), below average ($-2SD$ to $-1SD$, Dark blue), average ($-1SD$ to $+1SD$, Sky blue), above average ($+1SD$ to $+2SD$, Orange) and well above average ($> +2SD$, Red). Stillbirth rates estimated by maternal ethnicity and IMD were compared to the overall national average or national average by ethnic/socioeconomic group. We determined the average rate of stillbirth per each IMD and ethnic group, as well as the national stillbirth rate across all ethnic and IMD groups. Average stillbirth rates estimated by maternal ethnicity and IMD were compared to the national average. We then performed a one-sample *t*-test to compare the average stillbirth rate for each group to the national average. All statistical analyses were performed in RStudio statistical software package version 4.0.2 [23].

We determined the mean stillbirth rate in each health trust, with both the 95% and 99.8% confidence intervals around the mean for each individual health trust. We then plotted the national average of stillbirths to establish if the confidence intervals of each individual health trust are above or below the national average [24].

3 | Results

The maternal characteristics for the 1 260 567 births (Figure S2) and 4890 stillbirths (3.4 stillbirths/1000 births) are shown in Table 1 Most stillbirths occurred in nulliparous women (41.1%,

TABLE 1 | Participant characteristics.

Characteristics	Overall (<i>n</i> = 1260567)	Livebirths (<i>n</i> = 1255677)	Stillbirths (<i>n</i> = 4890)	Stillbirth rate (Per 1000 births)
Maternal age (years)				
< 20	39 515 (3.2%)	39 340 (3.2%)	175 (3.6%)	4.4
20–24	184 745 (14.8%)	183 966 (14.8%)	779 (16.0%)	4.2
25–29	351 010 (28.1%)	349 706 (28.1%)	1304 (26.8%)	3.7
30–34	394 907 (31.6%)	393 524 (31.6%)	1383 (28.4%)	3.5
≥ 35	278 879 (22.3%)	277 652 (22.3%)	1227 (25.2%)	4.4
Unknown	11 511	11 489	22	
Parity				
0 (Nulliparous)	457 517 (40.6%)	455 709 (40.6%)	1808 (41.1%)	4.0
1	395 590 (35.2%)	395 305 (35.2%)	1285 (29.2%)	3.2
2	164 371 (14.6%)	163 689 (14.6%)	682 (15.5%)	4.1
3	64 262 (5.7%)	63 930 (5.7%)	332 (7.5%)	5.2
4	24 754 (2.2%)	24 607 (2.2%)	151 (3.4%)	6.1
≥ 5 (Grand multiparous)	19 127 (1.7%)	18 984 (1.7%)	143 (3.3%)	7.5
Unknown	133 942	133 453	489	
Body mass index (kg/m²)				
< 18.5	28 769 (2.9%)	28 667 (2.9%)	102 (2.8%)	3.5
18.5 to < 25	472 278 (48.2%)	470 775 (48.3%)	1503 (41.4%)	3.2
25 to < 30	272 786 (27.9%)	271 721 (27.9%)	1075 (29.6%)	3.9
30 to < 35	126 463 (12.9%)	125 931 (12.9%)	532 (14.7%)	4.2
≥ 35	78 685 (8.1%)	78 269 (8.0%)	416 (11.5%)	5.3
Unknown	281 576	280 314	1262	
Ethnicity				
Asian	136 383 (11.9%)	135 644 (11.9%)	741 (16.7%)	5.4
Black	57 774 (5.1%)	57 365 (5.1%)	409 (9.2%)	7.1
Mixed	21 734 (1.9%)	21 637 (1.9%)	97 (2.2%)	3.9
Others	48 994 (4.3%)	48 802 (4.3%)	192 (4.3%)	4.5
White	875 217 (76.8%)	872 217 (76.8%)	3000 (67.6%)	3.4
Unknown	120 463	120 012	451	
Index multiple deprivation				
1 (most deprived)	317 294 (26.7%)	315 799 (26.8%)	1495 (33.5%)	4.7
2	267 810 (22.6%)	266 747 (22.6%)	1063 (23.8%)	4.0
3	222 781 (18.9%)	221 984 (18.8%)	797 (17.9%)	3.6
4	198 009 (16.8%)	197 414 (16.8%)	595 (13.4%)	3.0
5 (least deprived)	177 334 (15.0%)	176 826 (15.0%)	508 (11.4%)	2.9
Unknown	77 339	76 907	432	

$n=1808$), those aged 30–34 years (28.4%, $n=1383$) and with a BMI between 18.5 and 25 kg/m² (41.4%, $n=1503$). However, the highest rates of stillbirth (4.4/1000 births) were observed at the extremes of maternal age (<20 years and ≥35 years), respectively, in women with five or more births (7.5/1000 births) and those with a BMI ≥35 kg/m² (5.3/1000 births). The variation in stillbirth rate across health trusts in NHS England is shown in Figure 1 and Figure S3. The proportion of health trusts classified as having stillbirth rates well below average was 5.3% (Green, $n=7$), whereas the proportion classified as well above average was 1.5% (Red, $n=2$).

3.1 | Stillbirth Rate Variation

The stillbirth rate in average health trusts (blue) was similar for all ethnicities, ranging from 3.7/1000 births in White women up to 4.1/1000 births for Asian women (Table 1). The stillbirth rates in the well above average (red) health trusts varied from 9.8/1000 births in White women to 15.6/1000 births in Black women. The proportions of health trusts with well above average (red) stillbirth rates for White, Asian and Black women were 0.8%, 21.8%, and 38.6%, respectively (Figure 2). When health trusts were ranked in the same order according to the overall stillbirth rate in White women, there were notable variations in rates of stillbirth within the same health trust for White, Asian and Black women (Figure S4). Some health trusts demonstrating lower than average stillbirth rates for White women concurrently demonstrated higher than average stillbirth rates for Asian and/or Black women. There were no units exhibiting lower than average stillbirth rates for Asian/Black women while concurrently having higher than average stillbirth rates for White women.

The proportion of health trusts with well above average stillbirth rates was 4.5% for women living in the least deprived areas, while the corresponding figure was 17.3% for those living in the most deprived areas, respectively (Figure 3). When health trusts were ranked according to overall stillbirth rate in White women, there were notable variations in rates of stillbirth within the same health trust for the least and most deprived women (Figures S5 and S6). The proportions of health trusts with well above average stillbirth rates for White, Asian and Black women living in the most deprived areas were 15%, 27.8%, and 31.2%, respectively; The stillbirth rates in White, Asian and Black women from the most deprived areas were 4.3/1000 births, 6.7/1000 births and 5.7/1000 births, respectively (Figure S7). The rates from the least deprived areas were 2.6/1000 births, 5.9/1000 births, and 4.6/1000 births, respectively, with fewer trusts well above average stillbirth rates for White women (Figure S8). The rates of stillbirth by ethnicity or socioeconomic level are presented in Table S1.

4 | Discussion

This study investigated stillbirths across 133 NHS health trusts in England and confirmed the health disparity conferred by both ethnicity and socioeconomic deprivation. Most notably, some health trusts reported below average stillbirth rates for White women and concurrently reported above average stillbirth rates for Black and Asian women delivering in the same trust. Similar findings were evident for women living in areas with the highest and lowest socioeconomic deprivation. Stillbirth rates are concordant with socioeconomic deprivation, with women living in the poorest areas having the highest stillbirth rates and accounting for the majority (IMD 1 and





FIGURE 1 | Variation in stillbirth rate across health trusts in NHS England compared to the national rate (3.4 stillbirths/1000 births).



Ethnic differences in stillbirth rates across NHS Trusts

FIGURE 2 | Variation in stillbirth rate across health trusts in NHS England by ethnicity compared to the national rate (3.4 stillbirths/1000 births).

2, $n = 2802$, 57.3%) of all stillbirths. Similarly, Black and Asian women had the highest stillbirth rates, accounting for 25.9% ($n = 1150$) of stillbirths.

A substantial portion of the variation in stillbirth rates occurring in England can be attributed to socioeconomic and ethnic inequality [10]. Ethnicity is a social construct with strong intersectionality with socioeconomic deprivation [25]. Our findings are consistent with previous studies conducted in the UK, Europe and North America, showing that ethnic minority mothers residing in the most socioeconomically deprived neighbourhoods were more likely to have stillbirths [4, 6, 9]. These could be attributed to the inverse relationship between socioeconomic deprivation and access to quality perinatal care [26, 27]. Considering the wide disparity in stillbirths among mothers of the same ethnicity or neighbourhood and the existing evidence of differences in the quality of perinatal care received by women, patient-level characteristics cannot fully explain the observed disparities in stillbirth without considering the context and content of healthcare received by these women. Women of low socioeconomic status are more likely to face obstacles such as being disadvantaged and vulnerable [28]; therefore, poverty could be the leading factor preventing equal access to maternity care. To address inequalities in maternity care, enhancement in living standards for disadvantaged women is required to provide access to education and increase employment opportunities [29]. Addressing the complex association between stillbirth and socioeconomic deprivation will depend upon understanding these underlying patient-level factors influencing stillbirth [30]. Inequity in access to quality perinatal care due to mistrust of health services, language/communication difficulties, racial discrimination, poor nutrition, tobacco


use, alcohol consumption and substance use could be central to these inequalities [31–35]. Several medical audits, including the Perinatal Confidential Enquiries, have been carried out in the UK, all of which highlight the importance of targeted perinatal care in reducing avoidable stillbirths [36–39].

4.1 | Clinical Implications

Ethnic minority mothers living in the most deprived areas had the highest risk of stillbirth. Though previous studies conducted in the UK [7, 40, 41] and other European countries [42–44] have identified ethnicity as an independent risk factor for stillbirth, this is the first study to report ethnic and socioeconomic disparities in stillbirth rates at the level of individual health trusts. Both ethnicity and socioeconomic deprivation predispose individuals to adverse pregnancy outcomes through complex mechanisms—biological, financial, social and cultural. However, it is apparent that some trusts with below average stillbirth rates for White women demonstrated above average stillbirth rates for Black and Asian women. This observed disparity in stillbirth rates within the same health trust suggests that inequalities in access and/or delivery of quality antenatal care may be an important target for intervention and improvement [45, 46].

4.2 | Research and Health Policy Implications

The complex mechanisms responsible for the increase in stillbirth rates with ethnicity and socioeconomic deprivation do not necessarily mean that solutions should be complicated or



Index of multiple deprivation (IMD) and differences in stillbirth rates across NHS Trusts

FIGURE 3 | Variation in stillbirth rate across health trusts in NHS England by Index of Multiple Deprivation (IMD) compared to the national rate (3.4 stillbirths/1000 births).

impossible. Even though stillbirth rates are highest in Black women, 9% of pregnancy losses are formed by this ethnic group; the proportion of stillbirths was highest in White women (67.6%, $n=3000$) and from the most deprived neighbourhoods (33.5%, $n=1495$). This distribution of stillbirth suggests that targeting interventions based on ethnicity alone is unlikely to be effective in reducing stillbirth rates and could even perpetuate the flawed societal concept that ethnic and racial categories are biological determinants of health [47, 48]. The latter approach also runs the risk of stigmatisation of women on the basis of their ethnicity and also worsening the very health inequalities that need addressing [49]. A recent study reported a threefold reduction in perinatal death in Black and Asian women after early pregnancy risk personalised assessment using a model that included demographic, biophysical and biochemical characteristics [50, 51]. Targeting interventions based on risk prediction models that also include ethnicity and socioeconomic deprivation may represent the most effective approach towards stillbirth reduction. These findings should guide care and policymakers in addressing this challenge [29, 52].

4.3 | Strengths and Limitations

Some trusts were smaller than others, and the number of mothers from specific ethnic and/or IMD groups also varied between trusts. However, this study is using data from all trusts in NHS England over several years and is therefore providing robust, representative evidence. This is not an aetiological study aiming to further our understanding of the causal factors of stillbirth but a report of the variability of stillbirth among mothers of different

ethnicities and IMD groups. This study utilised the most reliable official routine maternity service data in England. Some studies have examined the geographical differences in stillbirths to quantify the neighbourhood effect on stillbirths [14, 53]. This study explores disparities across NHS Trusts and further considers inequality-related factors. A limitation is that no causal relationship can be established as this is a descriptive study. Other inequality-related factors, such as unemployment, cultural/religious beliefs, non-English language and migration status, were not available in the data source used. Therefore, these factors could explain the inequalities observed. Ethnicity and IMD are likely to have a direct but also indirect relationship on stillbirth, and factors such as cultural beliefs, language barriers and access to care are likely to play an important role in the occurrence of this adverse outcome. However, this is a descriptive study that highlights inequalities that currently exist. Thus, disparities in stillbirth could be reduced by targeting populations that have higher than average rates of stillbirth as early as possible in the antenatal care pathway, as well as health trusts with demonstrable inequalities in care delivery. Some mitigation is provided using IMD metrics, which capture employment deprivation among other factors related to deprivation in the area where a woman lives. However, IMD is a broad measure and cannot give information specifically about the individual social class of women living in a particular area. Furthermore, we were unable to subdivide the presented ethnic groups and, therefore, could not examine internal variation within each ethnic group, thus potentially masking inequalities. In addition, the availability of individual-level data for each woman, rather than the use of aggregated national data, allowed comparisons in stillbirth rates based upon maternal ethnicity and socioeconomic background.

However, the number of livebirths and stillbirths was very small for some trusts, especially when examining the intersection between ethnicity and socioeconomic background; thus, there is an increased chance of false positives [54]. In this descriptive study, we solely focused on stillbirth rather than perinatal mortality, as neonatal deaths are typically less frequent and therefore prone to substantial variability. In addition, neonatal deaths are subjected to confounding by acute care in labour rather than antenatal care provision, which was the focus of this analysis.

5 | Conclusion

Risk of stillbirth varied substantially by ethnic group and/or socioeconomic deprivation within individual health trusts. This is a descriptive study that highlights inequalities that currently exist, but as not all social and cultural confounders were available, inference cannot be established. This study emphasises the importance of considering factors that result in variability in delivery and/or access to healthcare at the level of the health trust. We also demonstrate that strategies to reduce stillbirth have to target both ethnic minority women and those who are socioeconomically deprived, if existing disparities are to be reduced. Findings from this study should guide care and policy stakeholders in prioritising interventions addressing the important public health challenge of reducing stillbirth.

Author Contributions

All authors were involved in the conception, design and data collection of the study. G.K., E.L., and A.J. performed data analysis, and G.K., A.H., E.L., A.J. and B.T. drafted the first version of the manuscript. All authors critically reviewed the article and contributed to interpreting the results. The final version of the manuscript was submitted for publication after obtaining approval from all authors.

Acknowledgements

The authors are grateful to Tommy's Charity, United Kingdom, for the financial support of this study towards alleviating the stillbirth impact in the United Kingdom. We also thank the Royal College of Midwives and the Royal College of Obstetrics and Gynaecology, for their joint leadership of the Tommy's National Centre for Maternity Improvement. We extend our gratitude to all members of the Tommy's National Centre for Maternity Improvement team and to the patient and public involvement group members for their consultation and input, which forms a key component in our publication procedure. J.S. is an NIHR Senior Investigator and is supported by the National Institute for Health Research (NIHR) Applied Research Collaboration South London (NIHR ARC South London) at King's College Hospital NHS Foundation Trust. E.L. and A.J. are supported by the NIHR Biomedical Research Centre at University Hospitals Bristol and Weston NHS Foundation Trust and the University of Bristol. The views expressed are those of the authors and not necessarily those of the NIHR or the Department of Health and Social Care.

Conflicts of Interest

The authors declare no conflicts of interest.

Data Availability Statement

The data that support the findings of this study are available from the National Maternity and Perinatal Audit] but restrictions apply to the

availability of these data, which were used under licence for the current study, and so are not publicly available. Data are, however, available from the authors upon reasonable request and with the permission of [National Maternity and Perinatal Audit].

References

1. A. E. P. Heazell, D. Siassakos, H. Blencowe, et al., "Stillbirths: Economic and Psychosocial Consequences," *Lancet* 387, no. 10018 (2016): 604–616.
2. H. Mistry, A. E. Heazell, O. Vincent, and T. Roberts, "A Structured Review and Exploration of the Healthcare Costs Associated With Stillbirth and a Subsequent Pregnancy in England and Wales," *BMC Pregnancy and Childbirth* 13 (2013): 236.
3. United Nations Inter-agency Group for Child Mortality Estimation, "Never Forgotten the Situation of Stillbirth Around the Globe Report," 2022 United Nations Children's Fund; 2023, <https://data.unicef.org/resources/never-forgotten-stillbirth-estimates-report/>.
4. S. E. Seaton, D. J. Field, E. S. Draper, et al., "Socioeconomic Inequalities in the Rate of Stillbirths by Cause: A Population-Based Study," *BMJ Open* 2, no. 3 (2012): e001100, <https://doi.org/10.1136/bmjopen-2012-001100>.
5. L. K. Smith, B. N. Manktelow, E. S. Draper, E. M. Boyle, S. J. Johnson, and D. J. Field, "Trends in the Incidence and Mortality of Multiple Births by Socioeconomic Deprivation and Maternal Age in England: Population-Based Cohort Study," *BMJ Open* 4, no. 4 (2014): e004514.
6. K. E. Best, S. E. Seaton, E. S. Draper, et al., "Assessing the Deprivation Gap in Stillbirths and Neonatal Deaths by Cause of Death: A National Population-Based Study," *Archives of Disease in Childhood. Fetal and Neonatal Edition* 104, no. 6 (2019): F624–F630, <https://doi.org/10.1136/archdischild-2018-316124>.
7. N. Penn, E. Oteng-Ntim, L. L. Oakley, and P. Doyle, "Ethnic Variation in Stillbirth Risk and the Role of Maternal Obesity: Analysis of Routine Data From a London Maternity Unit," *BMC Pregnancy and Childbirth* 14 (2014): 404.
8. S. F. Villadsen, L. H. Mortensen, and A. M. Andersen, "Ethnic Disparity in Stillbirth and Infant Mortality in Denmark 1981–2003," *Journal of Epidemiology and Community Health* 63, no. 2 (2009): 106–112.
9. E. S. Draper, I. D. Gallimore, L. K. Smith, et al., "MBRRACE-UK Perinatal Mortality Surveillance Report. UK Perinatal Deaths for Births from January to December 2018," 2021 Leicester: The Infant Mortality and Morbidity Studies, Department of Health Sciences, University of Leicester, 2020 March 28.
10. J. Jardine, K. Walker, I. Gurol-Urganci, et al., "Adverse Pregnancy Outcomes Attributable to Socioeconomic and Ethnic Inequalities in England: A National Cohort Study," *Lancet* 398, no. 10314 (2021): 1905–1912.
11. E. S. Draper, I. D. Gallimore, L. K. Smith, et al., "MBRRACE-UK Perinatal Mortality Surveillance Report. UK Perinatal Deaths for Births from January to December 2013," 2015 The Infant Mortality and Morbidity Studies, Department of Health Sciences, University of Leicester.
12. E. S. Draper, I. D. Gallimore, L. K. Smith, et al., "MBRRACE-UK Perinatal Mortality Surveillance Report. UK Perinatal Deaths for Births from January to December 2016," 2021 Leicester: The Infant Mortality and Morbidity Studies, Department of Health Sciences, University of Leicester; 2018 March 28.
13. E. S. Draper, I. D. Gallimore, L. K. Smith, et al., "MBRRACE-UK Perinatal Mortality Surveillance Report-UK Perinatal Deaths for Births from January to December 2020," 2022, <https://www.npeu.ox.ac.uk/mbrrace-uk/reports>.
14. S. Pattenden, K. Casson, S. Cook, and H. Dolk, "Geographical Variation in Infant Mortality, Stillbirth and Low Birth Weight in Northern

- Ireland, 1992-2002," *Journal of Epidemiology and Community Health* 65, no. 12 (2011): 1159-1165.
15. MBRRACE-UK, "Perinatal Mortality Rates by Trust/Health Board," <https://timms.le.ac.uk/mbrrace-uk-perinatal-mortality/2022>.
 16. "National Maternity and Perinatal Audit," 2023, <https://maternityaudit.org.uk/pages/home>.
 17. S. J. Stock, M. Horne, M. Bruijn, et al., "Development and Validation of a Risk Prediction Model of Preterm Birth for Women With Preterm Labour Symptoms (The QUIDS Study): A Prospective Cohort Study and Individual Participant Data Meta-Analysis," *PLoS Medicine* 18, no. 7 (2021): e1003686.
 18. H. Aughey, A. Blotkamp, F. Carroll, et al., "National Maternity and Perinatal Audit Clinical Report 2019."
 19. *National Statistics: English Indices of Deprivation 2015* (Office of National statistics, 2015), <https://www.gov.uk/government/statistics/english-indices-of-deprivation-2015>.
 20. Gov.UK, "List of Ethnic Groups 2021," <https://www.ethnicity-facts-figures.service.gov.uk/style-guide/ethnic-groups/>.
 21. J. E. Jardine, A. Frémeaux, M. Coe, I. Gurol Urganç, D. Pasupathy, and K. Walker, "Validation of Ethnicity in Administrative Hospital Data in Women Giving Birth in England: Cohort Study," *BMJ Open* 11, no. 8 (2021): e051977.
 22. Y. Ben-Shlomo, A. Blom, C. Boulton, et al., "The National Joint Registry 17th Annual Report 2020," London: National Joint Registry; 2020 Sep. Unit-level activity and outcomes 2020, <https://www.ncbi.nlm.nih.gov/books/NBK566661/>.
 23. Team RC, "A Language and Environment for Statistical Computing: R Foundation for Statistical Computing," <https://www.R-project.org/>.
 24. S. Hawley, D. Inman, C. L. Gregson, M. Whitehouse, A. Johansen, and A. Judge, "Predictors of Returning Home After Hip Fracture: A Prospective Cohort Study Using the UK National Hip Fracture Database (NHFD)," *Age and Ageing* 51, no. 8 (2022): afac131.
 25. M. A. Bohren, A. Iyer, A. J. D. Barros, et al., "Towards a Better Tomorrow: Addressing Intersectional Gender Power Relations to Eradicate Inequities in Maternal Health," *EclinicalMedicine* 67 (2024): 102180.
 26. P. Sauvegrain, M. Carayol, A. Piedvache, et al., "Understanding High Rates of Stillbirth and Neonatal Death in a Disadvantaged, High-Migrant District in France: A Perinatal Audit," *Acta Obstetrica et Gynecologica Scandinavica* 99, no. 9 (2020): 1163-1173.
 27. S. Hesselman, A. K. Wikström, A. Skalkidou, I. Sundström-Poromaa, and A. Wikman, "Neighborhood Deprivation and Adverse Perinatal Outcomes in Sweden: A Population-Based Register Study," *Acta Obstetrica et Gynecologica Scandinavica* 98, no. 8 (2019): 1004-1013.
 28. S. M. Harvey and K. S. Faber, "Obstacles to Prenatal Care Following Implementation of a Community-Based Program to Reduce Financial Barriers," *Family Planning Perspectives* 25, no. 1 (1993): 32-36.
 29. V. Flenady, P. Middleton, G. C. Smith, et al., "Stillbirths: The Way Forward in High-Income Countries," *Lancet* 377, no. 9778 (2011): 1703-1717.
 30. C. Kingdon, D. Roberts, M. A. Turner, et al., "Inequalities and Stillbirth in the UK: A Meta-Narrative Review," *BMJ Open* 9, no. 9 (2019): e029672.
 31. Y. Li, M. A. Quigley, A. Macfarlane, H. Jayaweera, J. J. Kurinczuk, and J. Hollowell, "Ethnic Differences in Singleton Preterm Birth in England and Wales, 2006-12: Analysis of National Routinely Collected Data," *Paediatric and Perinatal Epidemiology* 33, no. 6 (2019): 449-458.
 32. C. Opondo, R. Gray, J. Hollowell, Y. Li, J. J. Kurinczuk, and M. A. Quigley, "Joint Contribution of Socioeconomic Circumstances and Ethnic Group to Variations in Preterm Birth, Neonatal Mortality and Infant Mortality in England and Wales: A Population-Based Retrospective Cohort Study Using Routine Data From 2006 to 2012," *BMJ Open* 9, no. 7 (2019): e028227.
 33. W. M. Abel and J. T. Efid, "The Association Between Trust in Health Care Providers and Medication Adherence Among Black Women With Hypertension," *Frontiers in Public Health* 1 (2013): 66.
 34. K. Armstrong, M. Putt, C. H. Halbert, et al., "Prior Experiences of Racial Discrimination and Racial Differences in Health Care System Distrust," *Medical Care* 51, no. 2 (2013): 144-150.
 35. S. Yeo, "Language Barriers and Access to Care," *Annual Review of Nursing Research* 22 (2004): 59-73.
 36. R. Rowe, E. S. Draper, S. Kenyon, et al., "Intrapartum-Related Perinatal Deaths in Births Planned in Midwifery-Led Settings in Great Britain: Findings and Recommendations From the ESMiE Confidential Enquiry," *BJOG: An International Journal of Obstetrics and Gynaecology* 127, no. 13 (2020): 1665-1675.
 37. T. Norris, B. N. Manktelow, L. K. Smith, and E. S. Draper, "Causes and Temporal Changes in Nationally Collected Stillbirth Audit Data in High-Resource Settings," *Seminars in Fetal & Neonatal Medicine* 22, no. 3 (2017): 118-128.
 38. E. S. Draper, J. J. Kurinczuk, S. Kenyon, on behalf of MBRRACE-UK, "MBRRACE-UK 2015 Perinatal Confidential Enquiry: Term, Singleton, Normally-Formed, Antepartum Stillbirth 2015," 2015 The Infant Mortality and Morbidity Studies, Department of Health Sciences, University of Leicester: Leicester.
 39. E. S. Draper, J. J. Kurinczuk, S. Kenyon, and on behalf of MBRRACE-UK, "MBRRACE-UK Perinatal Confidential Enquiry: Term, Singleton, Intrapartum Stillbirth and Intrapartum-Related Neonatal death," 2017 The Infant Mortality and Morbidity Studies, Department of Health Sciences, University of Leicester: Leicester.
 40. L. M. Webster, K. Bramham, P. T. Seed, et al., "Impact of Ethnicity on Adverse Perinatal Outcome in Women With Chronic Hypertension: A Cohort Study," *Ultrasound in Obstetrics & Gynecology* 54, no. 1 (2019): 72-78.
 41. J. Gardosi, V. Madurasinghe, M. Williams, A. Malik, and A. Francis, "Maternal and Fetal Risk Factors for Stillbirth: Population Based Study," *BMJ* 346 (2013): f108.
 42. J. Zeitlin, E. Combier, F. De Caunes, and E. Papiernik, "Socio-Demographic Risk Factors for Perinatal Mortality. A Study of Perinatal Mortality in the French District of Seine-Saint-Denis," *Acta Obstetrica et Gynecologica Scandinavica* 77, no. 8 (1998): 826-835.
 43. B. Essén, B. S. Hanson, P. O. Ostergren, P. G. Lindquist, and S. Gudmundsson, "Increased Perinatal Mortality Among Sub-Saharan Immigrants in a City-Population in Sweden," *Acta Obstetrica et Gynecologica Scandinavica* 79, no. 9 (2000): 737-743.
 44. S. Vangen, C. Stoltenberg, R. E. Johansen, J. Sundby, and B. Stray-Pedersen, "Perinatal Complications Among Ethnic Somalis in Norway," *Acta Obstetrica et Gynecologica Scandinavica* 81, no. 4 (2002): 317-322.
 45. L. Goodwin, A. Jones, and B. Hunter, "Addressing Social Inequity Through Improving Relational Care: A Social-Ecological Model Based on the Experiences of Migrant Women and Midwives in South Wales," *Health Expectations* 25, no. 5 (2022): 2124-2133.
 46. R. Wood, D. Stockton, and H. Brown, "Moving From a Universal to Targeted Child Health Programme: Which Children Receive Enhanced Care? A Population-Based Study Using Routinely Available Data," *Child: Care, Health and Development* 39, no. 6 (2013): 772-781.
 47. A. Heazell, J. Budd, L. K. Smith, et al., "Associations Between Social and Behavioural Factors and the Risk of Late Stillbirth - Findings From the Midland and North of England Stillbirth Case-Control Study," *BJOG: An International Journal of Obstetrics and Gynaecology* 128, no. 4 (2021): 704-713.
 48. A. C. Ravelli, M. Tromp, M. Eskes, et al., "Ethnic Differences in Stillbirth and Early Neonatal Mortality in The Netherlands," *Journal of*

Epidemiology and Community Health 65, no. 8 (2011): 696–701, <https://doi.org/10.1136/jech.2009.095406>.

49. B. Liu, M. Alakaloko, A. Frick, A. Bhide, and B. Thilaganathan, “Authors’ Reply Re: The Dangers of Biological Essentialism in Addressing Birth Equity,” *BJOG: An International Journal of Obstetrics and Gynaecology* 129, no. 11 (2022): 1945–1946.

50. B. Liu, U. Nadeem, A. Frick, M. Alakaloko, A. Bhide, and B. Thilaganathan, “Reducing Health Inequality in Black, Asian and Other Minority Ethnic Pregnant Women: Impact of First Trimester Combined Screening for Placental Dysfunction on Perinatal Mortality,” *BJOG: An International Journal of Obstetrics and Gynaecology* 129, no. 10 (2022): 1750–1756.

51. J. Carter, D. Anumba, L. Brigante, et al., “The Tommy’s Clinical Decision Tool, a Device for Reducing the Clinical Impact of Placental Dysfunction and Preterm Birth: Protocol for a Mixed-Methods Early Implementation Evaluation Study,” *BMC Pregnancy and Childbirth* 22, no. 1 (2022): 639.

52. O. A. Esan, N. K. Adjei, S. Saberian, et al., “Mapping Existing Policy Interventions to Tackle Ethnic Health Inequalities in Maternal and Neonatal Health in England: A Systematic Scoping Review With Stakeholder Engagement,” www.nhs.uk/publications/new-research-identifies-gaps-in-ethnicity-research-in-maternal-care-2 2022.

53. H. O. Dickinson, J. L. Hutton, L. H. Greaves, T. J. Dummer, and L. Parker, “Deprivation and Stillbirth Risk in Rural and Urban Areas,” *Paediatric and Perinatal Epidemiology* 16, no. 3 (2002): 249–254.

54. A. D. Althouse, “Adjust for Multiple Comparisons? It’s Not That Simple,” *Annals of Thoracic Surgery* 101, no. 5 (2016): 1644–1645.

Supporting Information

Additional supporting information can be found online in the Supporting Information section.