

RAPID REPORT

Inhibition of native TRPC6 channel activity by phosphatidylinositol 4,5-bisphosphate in mesenteric artery myocytes

Anthony P. Albert, Sohag N. Saleh and William A. Large

Ion Channels and Cell Signalling Research Centre, Division of Basic Medical Sciences, St George's, University of London, Cranmer Terrace, London SW17 0RE, UK

The present work investigates the effect of phosphatidylinositol-4,5-bisphosphate (PIP₂) on native TRPC6 channel activity in freshly dispersed rabbit mesenteric artery myocytes using patch clamp recording and co-immunoprecipitation methods. Inclusion of 100 μM diC8-PIP₂ in the patch pipette and bathing solutions, respectively, inhibited angiotensin II (Ang II)-evoked whole-cell cation currents and TRPC6 channel activity by over 90%. In inside-out patches diC8-PIP₂ also inhibited TRPC6 activity induced by the diacylglycerol analogue 1-oleoyl-2-acetyl-*sn*-glycerol (OAG) with an IC₅₀ of 7.6 μM. Anti-PIP₂ antibodies potentiated Ang II- and OAG-evoked TRPC6 activity by about 2-fold. Depleters of tissue PIP₂ wortmannin and LY294002 stimulated TRPC6 activity, as did the polycation PIP₂ scavenger poly-L-lysine. Wortmannin reduced Ang II-evoked TRPC6 activity by over 75% but increased OAG-induced TRPC6 activity by over 50-fold. Co-immunoprecipitation studies demonstrated association between PIP₂ and TRPC6 proteins in tissue lysates. Pre-treatment with Ang II, OAG and wortmannin reduced TRPC6 association with PIP₂. These results provide for the first time compelling evidence that constitutively produced PIP₂ exerts a powerful inhibitory action on native TRPC6 channels.

(Received 10 March 2008; accepted after revision 1 May 2008; first published online 8 May 2008)

Corresponding author A. Albert: Ion Channels & Cell Signalling Research Centre, Division of Basic Medical Sciences, St George's, University of London, Cranmer Terrace, London SW17 0RE, UK. Email: aalbert@sgul.ac.uk

Members of the canonical class of transient receptor potential (TRPC) non-selective Ca²⁺-permeable cation channels are present in many types of smooth muscle (see Large, 2002; Beech *et al.* 2004; Albert & Large, 2006; Albert *et al.* 2007). These channels have been implicated in many physiological responses in vascular tissue such as depolarization, contraction, gene expression, cell growth and proliferation. Classically TRPC channels are often described as receptor-operated (ROCs) or store-operated channels (SOCs) with ROCs being stimulated by G-protein-coupled receptors linked to phospholipase C (PLC) and SOCs stimulated by depletion of internal Ca²⁺ stores. However, little is known about the precise mechanisms by which TRPC-mediated ROCs and SOCs are gated.

In rabbit vascular myocytes there are several TRPC ROC isoforms where one product of phosphatidylinositol-4,5-bisphosphate hydrolysis (PIP₂) by PLC, diacylglycerol (DAG), initiates channel opening by a protein kinase C (PKC)-independent mechanism. DAG

stimulates TRPC6 activity in portal vein and mesenteric artery, TRPC3 in ear artery and TRPC3/TRPC7 in coronary artery myocytes in this manner although it is not known how DAG produces channel gating (Helliwell & Large, 1997; Inoue *et al.* 2001; Albert *et al.* 2005, 2006; Saleh *et al.* 2006; Peppiatt-Wildman *et al.* 2007). Moreover the other product of PIP₂ hydrolysis, inositol 1,4,5-trisphosphate (IP₃), markedly potentiates TRPC6-like and TRPC3/TRPC7 channel opening in, respectively, portal vein and coronary artery myocytes (Albert & Large, 2003; Peppiatt-Wildman *et al.* 2007). In the present work we have investigated the role of PIP₂ in regulating native TRPC6 activity since PIP₂ is the precursor of both DAG and IP₃ and has been shown to independently regulate the function of many ion channel proteins including members of the TRP superfamily (see Suh & Hille, 2005; Rohacs, 2007; Voets & Nilius, 2007). Recently it was demonstrated that PIP₂ increased expressed TRPC3, -C6 and -C7 activity in HEK293 cells (Lemonnier *et al.* 2008) and it was suggested that

phosphoinositides, including PIP₂, mediate increases in TRPC6 activity due to disruption of calmodulin (CaM) binding to fusion proteins containing the C-termini of TRPC6 (Kwon *et al.* 2007). In contrast, PIP₂ inhibited receptor-operated TRPC4 α activity in HEK293 cells (Otsuguro *et al.* 2008). These results illustrate complex effects of PIP₂ on expressed TRPC channels but to date there have been no studies on the effect of PIP₂ on native TRPC channels. Therefore, in the present study, we investigated the effect of PIP₂ on TRPC6 channels in freshly dispersed rabbit mesenteric artery myocytes. These novel results show that PIP₂ exerts a powerful inhibitory brake on agonist-evoked TRPC6 activity. Moreover, simultaneous depletion of PIP₂ and production of DAG are necessary for optimal channel activation.

Methods

Cell isolation

New Zealand White rabbits (2–3 kg) were killed using i.v. sodium pentobarbitone (120 mg kg⁻¹, in accordance with the UK Animals (Scientific Procedures) Act, 1986). 1st to 5th order mesenteric arteries were dissected free from fat and connective tissue and enzymatically digested into single myocytes using methods previously described (Saleh *et al.* 2006).

Electrophysiology

Whole-cell and single cation currents were recorded with an Axopatch 200B patch-clamp amplifier (Axon Instruments, USA) at room temperature (20–23°C) using whole-cell recording, cell-attached, inside-out and outside-out patch configurations and data acquisition and analysis protocols as previously described (Saleh *et al.* 2006). Briefly, single channel current amplitudes were calculated from idealized traces of at least 60 s in duration using the 50% threshold method with events lasting for less than 0.664 ms (2 \times rise time for a 1 kHz, -3 db, low-pass filter) being excluded from analysis. Figure preparation was carried out using MicroCal Origin software 6.0 (MicroCal Software Inc., MA, USA) where inward single channel currents are shown as downward deflections. Open probability (NP_o) was calculated using the equation: $NP_o = \frac{\sum(O_n)}{T}$ where n is the number of channels in the patch, O_n is the time spent at each open level and T is the total recording time.

Immunoprecipitation and Western blotting

Dissected tissues were either flash frozen and stored at -80°C for subsequent use or immediately placed into 10 mg ml⁻¹ RIPA lysis buffer (Santa Cruz Biotechnology)

supplemented with protease inhibitors and then homogenized on ice by sonication for at least 3 h. The total cell lysate (TCL) was collected by centrifugation at 10 g for 10 min at 4°C and then protein content was quantified using the Bio-Rad protein dye reagent (Bradford method). TCL was pre-cleared using A/G agarose beads (Santa Cruz Biotechnology) and immunoprecipitated overnight at 4°C using the appropriate antibody and A/G agarose bead conjugate. Alternatively, the immunoprecipitation protocol was carried out using the Upstate Catch and Release kit (Millipore), where spin columns were loaded with 500 μ g of cell lysates, 4 μ g of antibody and immunoprecipitated for 2 h at room temperature.

Protein samples were eluted with Laemmli sample buffer and incubated at 95°C for 2 min. One-dimensional protein gel electrophoresis was performed in 4–12% Bis-Tris gels in a Novex mini-gel system (Invitrogen) with 20 μ g of total protein loaded in each lane. Separated proteins were electrophoretically transferred onto polyvinylidene difluoride membranes in a Biorad trans-blot SD semi-dry transfer cell or using the iBlot apparatus (Invitrogen). Blots were incubated for 1–4 h with 5% (weight/volume) non-fat milk in phosphate-buffered saline (PBS) containing 0.1% Tween 20 (PBST) to block non-specific protein binding. Membranes were then incubated with appropriate primary antibody overnight at 4°C (where possible alternative antibodies raised against different epitopes were used for immunoprecipitation and Western blot analysis). Following antibody removal blots were washed for 2 h with milk-PBST. Blots were subsequently incubated with horseradish peroxidase-conjugated goat anti-rabbit (Sigma) or sheep anti-goat (Santa Cruz Biotechnology) IgG secondary antibody diluted 1 : 1000–1 : 5000 in milk-PBST, washed 3 times in PBST, and treated with ECL chemiluminescence reagents (Pierce) for 1 min and exposed to photographic films. In loading control experiment anti- β -actin antibodies (mouse monoclonal, Sigma, UK) were also added to the immunoprecipitate and immunoblot procedures to show that expression levels of β -actin did not change during the experimental conditions. Moreover, immunoblot control data showed that anti-PIP₂ and anti-TRPC6 antibodies did not recognize β -actin following immunoprecipitation with only anti- β -actin antibodies. Data shown represent n values of at least three separate experiments.

Anti-TRPC6 and anti-PIP₂ antibodies

Polyclonal TRPC6 antibody generated in rabbits against an intracellular epitope was purchased from Alomone Laboratories (Jerusalem, Israel) and the selectivity and negligible cross-reactivity of this antibody for its target

protein has been previously confirmed (Liu *et al.* 2005; Sours *et al.* 2006). An alternative polyclonal antibody for TRPC6 generated in goats against a different intracellular epitope was purchased from Santa Cruz Biotechnology and the specificity of this antibody has also been previously confirmed (Liu *et al.* 2005). Mouse monoclonal PIP₂ antibody generated against liposomes of human origin constituted with the phospholipid was purchased from Assay Designs (USA). This anti-PIP₂ antibody has previously been used in electrophysiological (Liou *et al.* 1999; Bian *et al.* 2001; Ma *et al.* 2002; Pian *et al.* 2006; Xie *et al.* 2008) and immunoprecipitation studies (Asteggiano *et al.* 2001; Beauge *et al.* 2002; Yue *et al.* 2002) to investigate the role of PIP₂ in regulating ion channels and exchangers.

Solutions and drugs

The bathing solution used to measure whole-cell cation currents was K⁺ free and contained (mM): NaCl (126), CaCl₂ (1.5), Hepes (10), glucose (11), DIDS (0.1), niflumic acid (0.1) and nicardipine (0.005), pH to 7.2 with NaOH. In cell-attached patch experiments the membrane potential was set to approximately 0 mV by perfusing cells in a KCl external solution containing (mM): KCl (126), CaCl₂ (1.5), Hepes (10) and glucose (11), pH to 7.2 with 10 M KOH. Nicardipine (5 μM) was also included to prevent smooth muscle cell contraction by blocking Ca²⁺ entry through voltage-dependent Ca²⁺ channels.

The patch pipette solution used to measure whole-cell cation currents (intracellular solution) was also K⁺ free and contained (mM): CsCl (18), caesium aspartate (108), MgCl₂ (1.2), Hepes (10), glucose (11), BAPTA (10), CaCl₂ (4.8, free internal Ca²⁺ concentration approximately 100 nM as calculated using EQCAL software), Na₂ATP (1) and NaGTP (0.2), pH 7.2 with Tris. The patch pipette solution used for both cell-attached and inside-out patch recording (extracellular solution) was K⁺ free and contained (mmol l⁻¹): NaCl (126), CaCl₂ (1.5), Hepes (10), glucose (11), TEA (10), 4-AP (5), iberiotoxin (0.0002), DIDS (0.1), niflumic acid (0.1) and nicardipine (0.005), pH to 7.2 with NaOH. The composition of the bathing solution used in inside-out experiments (intracellular solution) was the same as the patch pipette solution used for whole-cell recording except that 1 mM BAPTA and 0.48 mM CaCl₂ were included (free internal Ca²⁺ concentration approximately 100 nM). Under these conditions VDCCs, K⁺ currents, swell-activated Cl⁻ currents and Ca²⁺-activated conductances are abolished and non-selective cation currents could be recorded in isolation.

DiC8-PIP₂ was purchased from Cayman chemicals (UK). All other drugs were purchased from Calbiochem (UK), Sigma (UK) or Tocris (UK) and agents were dissolved in distilled H₂O or DMSO (0.1%). DMSO alone had no effect on TRPC6 channel activity. The values are

the mean of *n* cells ± s.e.m. Statistical analysis was carried out using paired (comparing effects of agents on the same cell) or unpaired (comparing effects of agents between cells) Students' *t* test with the level of significance set at *P* < 0.05.

Results

DiC8-PIP₂ inhibits native TRPC6 channel activity in rabbit mesenteric artery myocytes

In initial experiments we investigated the effect of including 100 μM diC8-PIP₂, a water-soluble form of PIP₂, in the patch pipette solution on angiotensin II (Ang II)-induced whole-cell cation currents in mesenteric artery myocytes. In these experiments we used 1 nM Ang II since previous work showed that 1 nM Ang II activated native TRPC6 channels (Saleh *et al.* 2006). Figure 1*Aa* and *b* shows that 100 μM diC8-PIP₂ reduced the mean peak amplitude of whole-cell cation currents induced by 1 nM Ang II from -26 ± 8 pA (*n* = 7) to -2 ± 0.5 pA (*n* = 7, *P* < 0.01) at -50 mV. Figure 1*Ac* also shows that the Ang II-evoked mean current-voltage (*I*-*V*) relationship was inhibited by 100 μM diC8-PIP₂ at all membrane potentials between -100 mV and +90 mV.

To obtain precise information on the effect of PIP₂ on TRPC6 channel activity and to exclude the possibility of observing actions of PIP₂ on multiple cation conductances we studied the effect of PIP₂ on TRPC6 activity at the single channel level. Figure 1*Ba* shows that bath application of 1 nM Ang II evoked channel activity in a cell-attached patch, which had similar unitary properties to TRPC6 channels previously described (Saleh *et al.* 2006), and was maintained following excision of the patch into the inside-out configuration for up to 30 min. Figure 1*Ba* shows that bath application of 100 μM diC8-PIP₂ to the cytosolic surface of the inside-out patch reversibly inhibited the mean open probability (*NP*_o) of Ang II-evoked TRPC6 activity from 3.81 ± 0.84 to 0.43 ± 0.21 (*n* = 6, *P* < 0.05) at -50 mV. Previously it has been shown that DAG activates TRPC6 channels in mesenteric artery myocytes by a PKC-independent mechanism. Figure 1*Bb* shows that bath application of 10 μM 1-oleoyl-2-acetyl-*sn*-glycerol (OAG), a cell-permeable DAG analogue, to an inside-out patch also induced channel activity with TRPC6 properties (Saleh *et al.* 2006). Moreover, Fig. 1*Bb* shows that bath application of 100 μM diC8-PIP₂ reversibly inhibited mean *NP*_o of OAG-induced TRPC6 activity from 1.49 ± 0.03 to 0.09 ± 0.03 (*n* = 6, *P* < 0.01) at -50 mV. diC8-PIP₂ at 100 μM also inhibited OAG-evoked TRPC6 activity by $89 \pm 5\%$ (*n* = 3, data not shown) in the presence of the PKC inhibitor chelerythrine (3 μM) indicating that PIP₂-mediated inhibition of TRPC6 did not involve PKC (see Saleh *et al.* 2006). Figure 1*C* shows a dose-response

curve of diC8-PIP₂ on TRPC6 activity induced by 10 μM OAG in inside-out patches at -50 mV with 50% inhibition (IC_{50}) produced by 7.6 μM diC8-PIP₂.

Endogenous PIP₂ inhibits TRPC6 activity

The above data clearly demonstrate that diC8-PIP₂ has a profound inhibitory action on TRPC6 activity in mesenteric artery myocytes. Therefore we investigated the role of endogenous PIP₂ on TRPC6 activity by studying the effect of anti-PIP₂ antibodies on TRPC6 activity induced by Ang II and OAG in inside-out patches.

Figure 2Aa shows that bath application of anti-PIP₂ antibodies at 1:200 to an inside-out patch, in which TRPC6 channels were activated initially in the cell-attached mode by 1 nM Ang II (i.e. similar to Fig. 1B), increased TRPC6 activity (about 2-fold) but after 2 min there was a marked decrease in NP_o (Fig. 2Aa and B). Figure 2Ab illustrates that bath application of anti-PIP₂

antibodies at 1:200 dilution also increased OAG-evoked TRPC6 activity in inside-out patches by approximately 2-fold but unlike Ang II there was no subsequent inhibition of channel activity in the continued presence of the antibody (Fig. 2Ab and B).

Agents that deplete PIP₂ levels activate TRPC6 activity in un-stimulated mesenteric artery myocytes

These present results indicate that endogenous PIP₂ exerts a powerful inhibitory action of TRPC6 channels. In addition, previous work using an inhibitor of DAG lipase showed that these channels are also activated by constitutive DAG production (Saleh *et al.* 2006). We therefore investigated the effect of wortmannin and LY294002 on cell-attached patches from unstimulated myocytes since at high concentrations these agents are known to inhibit phosphoinositol (PI)-4-kinases leading to a reduction in the generation of PIP₂ and consequently a depletion of PIP₂ levels (Suh & Hille, 2005).




Figure 1. DiC8-PIP₂ inhibits Ang II-evoked whole-cell cation currents and single TRPC6 channel activity in rabbit mesenteric artery myocytes

Aa and b, inclusion of 100 μM diC8-PIP₂ in the patch pipette solution inhibits whole-cell cation currents evoked by 1 nM Ang II. Ac, mean $I-V$ relationship showing diC8-PIP₂ inhibits Ang II-evoked cation currents at all potentials tested ($n =$ at least 4 per point). Ba, 100 μM diC8-PIP₂ inhibited TRPC activity in an inside-out patch which was initially activated by 1 nM Ang II in a cell-attached patch (i/o, inside-out patch). Bb, 100 μM diC8-PIP₂ inhibited TRPC activity induced by 10 μM OAG in an inside-out patch. C, inhibitory action of diC8-PIP₂ on OAG-evoked TRPC6 activity in inside-out patches had an IC_{50} value of 7.6 μM ($n =$ at least 4 per point).

Figure 3*Aa* shows that bath application of 20 μM wortmannin evoked channel activity with a mean peak NP_o value of 1.79 ± 0.42 ($n = 14$) at -50 mV. Amplitude histograms of wortmannin-induced channel currents could be fitted by the sum of three Gaussian curves with three sub-conductance states of 15 pS, 28 pS and 44 pS which all had a reversal potential (E_r) of about 0 mV (Fig. 3*Ab* and *c*). These characteristics are identical to Ang II- and OAG-evoked TRPC6 channel currents (see Saleh *et al.* 2006). Figure 3*Ba*, *b* and *c* shows that bath application of 100 μM LY294002 evoked TRPC6 activity with a mean peak NP_o value of 0.37 ± 0.01 ($n = 60$) at -50 mV. TRPC6 activity induced by wortmannin and LY294002 was transient with no activity detected after 5 min in the presence of these agents. At concentrations that do not inhibit PI-4 kinases, wortmannin (50 nM) and LY294002 (10 μM) had no effect on TRPC6 activity.

Figure 3*Ca* shows that bath application of the poly-PIP₂ scavenger poly-L-lysine (50 $\mu\text{g ml}^{-1}$) to an inside-out patch activated channel activity with a mean peak NP_o value of 1.52 ± 0.34 ($n = 10$) at -50 mV. These poly-L-lysine-evoked channel currents had similar conductances/ E_r values to those induced by wortmannin and LY294002 (Fig. 3*Cb* and *c*).

Depletion of PIP₂ levels reduces Ang II- but increases OAG-evoked TRPC6 activity

These results show that endogenous and exogenous PIP₂ inhibits Ang II- and OAG-evoked TRPC6 activity and that PIP₂ also prevents channel opening induced by constitutive

generation of DAG. However, stimulation of AT₁ receptors by Ang II involves PLC-mediated PIP₂ hydrolysis leading to DAG production and TRPC6 activity (Saleh *et al.* 2006) whereas OAG acts on TRPC6 downstream of PIP₂ hydrolysis. Therefore we compared the effects of Ang II and OAG on TRPC6 activity in the presence of wortmannin in cell-attached patches.

Figure 4*Aa*, *b* and *C* shows that 15 min preincubation with 20 μM wortmannin reduced Ang II-evoked TRPC6 activity by about 75%. In contrast, Fig. 4*Ba*, *b* and *C* shows that 10 μM OAG produced little stimulation of TRPC6 activity in control cell-attached patches. However, following pre-treatment with 20 μM wortmannin for 15 min OAG-evoked TRPC6 activity was markedly increased by over 50-fold ($P < 0.001$).

Association of PIP₂ with TRPC6 proteins is reduced by pre-treatment with wortmannin and OAG

These data suggest that PIP₂ inhibits TRPC6 activity stimulated by Ang II and OAG but also that PIP₂ acts as a precursor for endogenous DAG production by Ang II. In contrast, sustained TRPC6 activation by OAG does not require PIP₂ as a DAG precursor. We investigated whether there is evidence for close association between PIP₂ and TRPC6 proteins using co-immunoprecipitation and Western blotting and whether association is altered in the presence of Ang II, OAG and wortmannin.

Figure 4*Da* illustrates that tissue lysates from mesenteric artery immunoprecipitated with anti-PIP₂ and then blotted with anti-TRPC6 antibodies displayed a band of




Figure 2. Anti-PIP₂ antibody potentiates TRPC6 channel activity

Aa, 1 : 200 dilution of anti-PIP₂ antibodies transiently increased TRPC6 activity in an inside-out patch which was induced by 1 nM Ang II in cell-attached mode. Subsequently the Ang II-evoked response was inhibited in the presence of the anti-PIP₂ antibody. *Ab*, 1 : 200 dilution of anti-PIP₂ antibodies produced a sustained increase of OAG-evoked TRPC6 activity in an inside-out patch. *B*, mean data of effect of anti-PIP₂ antibodies on Ang II- and OAG-induced TRPC6 activity ($n = 6$ for all conditions, ** $P < 0.01$).

~110 kDa which is the expected molecular weight for TRPC6 proteins. Complementary studies showed that association between TRPC6 and PIP₂ could be obtained using anti-TRPC6 for immunoprecipitation and anti-PIP₂ for blotting (data not shown). Figure 4*Da* also shows that incubation with 1 nM Ang II, 10 μM OAG and 20 μM wortmannin for 15 min before tissue lysis all reduced the association of TRPC6 proteins with PIP₂. Moreover, Fig. 4*Db* shows that wortmannin reduced, whereas Ang II and OAG had little effect, on total tissue PIP₂ levels detected with Western blotting. Figure 4*Da* and *c* illustrates that wortmannin, Ang II and OAG had no effect on expression levels of β-actin or TRPC6.

Discussion

The present data clearly show that in addition to acting as a precursor of DAG production endogenous PIP₂ also inhibits native TRPC6 channel activity in rabbit

mesenteric artery myocytes. Moreover, PIP₂ imposes a brake on TRPC6 activation by both Ang II and DAG. This latter result represents a novel physiological role for this important phospholipid in vascular smooth muscle excitation and may suggest that DAG activates TRPC6 channels by reducing this inhibitory action of PIP₂ on channel proteins.

DiC8-PIP₂ inhibited Ang II-evoked whole-cell currents and TRPC6 channel currents in inside-out patches. Moreover, DiC8-PIP₂ also inhibited OAG-induced TRPC6 activity in inside-out patches with an IC₅₀ of 7.6 μM, which is comparable to the affinity of PIP₂ binding to K⁺ channels (Zhang *et al.* 2003) and to fusion proteins composed of TRPC6 C-termini (Kwon *et al.* 2007). Anti-PIP₂ antibodies initially increased TRPC6 activity induced by Ang II, which is likely to be caused by removal of an inhibitory action of endogenous PIP₂. In the continued presence of anti-PIP₂ antibodies, Ang II-induced TRPC6 activity was greatly reduced, which presumably reflects a reduction in




Figure 3. Agents that deplete PIP₂ evoke TRPC6 channel activity

A and B, respectively, 20 μM wortmannin and 100 μM LY294002 transiently activated channel activity in cell-attached patches (a) which had similar amplitude histograms (b) and unitary conductances and E_r values (c). C, 50 μg ml⁻¹ poly-L-lysine activated channel currents in an inside-out patch (a) which had similar properties to those evoked by wortmannin and LY294002 (b and c).

PIP₂ required to produce endogenous PLC-mediated DAG production and TRPC6 opening. This is consistent with Ang II-evoked TRPC6 activity being almost completely blocked by U73122, a PLC inhibitor, indicating a pivotal role for PLC in the transduction pathway (Saleh *et al.* 2006). Predictably, anti-PIP₂ antibodies only potentiated OAG-evoked TRPC6 activity because OAG acts downstream of PLC-mediated PIP₂ hydrolysis. It has been shown that PIP₂ binds to expressed TRPC6 proteins in HEK293 cells (Tseng *et al.* 2004) and to fusion proteins composed of TRPC6 C-termini (Kwon *et al.* 2007), and the present co-immunoprecipitation work also shows that endogenous PIP₂ associates with native TRPC6 proteins in

unstimulated tissue lysates. The mechanism(s) by which anti-PIP₂ antibodies prevent the inhibitory action of PIP₂ is unknown. However, it is likely that recycling of PIP₂ means that this phospholipid can be free or bound to TRPC6. Therefore, a simple explanation is that the free PIP₂-anti-PIP₂ antibody complex binds less readily to TRPC6 and is less readily hydrolysed by PLC.

Further evidence for an inhibitory action of endogenous PIP₂ on TRPC6 activity is that wortmannin, LY294002 and poly-L-lysine, which all deplete PIP₂ levels, produced activation of TRPC6 channels in unstimulated myocytes. In mesenteric artery myocytes there is likely to be constitutive production of DAG because an inhibitor of




Figure 4. Wortmannin-induced depletion of PIP₂ decreases Ang II-evoked and increases OAG-induced TRPC6 activity in cell-attached patches

A shows that Ang II-evoked TRPC6 activity (a) is decreased after pre-treatment with wortmannin (b). B illustrates that OAG-induced TRPC6 activity (a) is markedly increased following pre-treatment with wortmannin (b). C, mean data showing effect of wortmannin on Ang II- and OAG-induced TRPC6 activity ($***P < 0.001$). Da shows a co-immunoprecipitation experiment in which mesenteric artery tissue lysates were immunoprecipitated (IP) with an anti-PIP₂ antibody and then Western blotted (WB) with an anti-TRPC6 antibody. In control conditions a band of ~110 kDa was observed which was absent after pre-treatment of the tissue with 20 μ M wortmannin, 1 nM Ang II and 10 μ M OAG for 15 min. Note that similar results were obtained with both anti-TRPC6 antibodies (Alomone and Santa Cruz). Db, immunoblot with anti-PIP₂ antibody showing reduction of total tissue PIP₂ levels after incubation with 20 μ M wortmannin for 15 min. Pre-treatment with 1 nM Ang II and 10 μ M OAG for 15 min had little effect on total tissue PIP₂ levels. Dc, immunoblot with anti-TRPC6 antibody showing pre-treatment of tissue with 20 μ M wortmannin, 1 nM Ang II and 10 μ M OAG and for 15 min had no effect on expression of TRPC6 protein. Note that all immunoprecipitation and immunoblot experiments were from at least $n = 3$.

DAG lipase, which is the main pathway for metabolism of DAG in smooth muscle (Severson & Hee-Cheong, 1989), activates TRPC6 channels (Saleh *et al.* 2006). It is likely that wortmannin and LY294002 evoked TRPC6 activity by inhibiting PI-4-kinases which produces a reduction in PIP₂ synthesis (Suh & Hille, 2005) due to recycling of this phospholipid. Thus leading to removal of its inhibitory action, and channel opening by constitutive generation of DAG. The transient nature of the excitatory effects of wortmannin and LY294002 is also consistent with the idea that opening of TRPC6 channels requires PIP₂-mediated DAG production. Therefore, depletion of PIP₂ will lead eventually to reduced DAG production and channel activation. Indeed, co-immunoprecipitation studies showed that wortmannin reduced both the association of PIP₂ with TRPC6 proteins and PIP₂ levels in unstimulated tissue lysates. Wortmannin also produced transient TRPC6 activity and greatly reduced noradrenaline-evoked TRPC6 currents in portal vein indicating that PIP₂ may have a similar action on TRPC6 in this preparation (Aromolaran *et al.* 2000; Inoue *et al.* 2001).

As expected, wortmannin reduced Ang II-evoked TRPC6 activity, which correlated with depletion of tissue PIP₂ levels and decreased association between PIP₂ and TRPC6 proteins demonstrated by Western blotting and co-immunoprecipitation. A notable result was that in the presence of wortmannin, OAG-evoked TRPC6 activity in cell-attached patches was enhanced by about 50-fold, whereas OAG-induced TRPC6 activity in inside-out patches was only increased 2-fold by an anti-PIP₂ antibody. These differences in potentiation of TRPC6 activity by depletion of PIP₂ probably relate to the relative activity of OAG alone in the different patch configurations with OAG evoking much larger activity in inside-out than in cell-attached patches (see Figs 1Ba and 4Ba). These differences in stimulation of TRPC6 by OAG may suggest that in inside-out patches OAG is more effective at removing the inhibitory action of PIP₂ and/or there are lower levels of PIP₂ in a patch following excision. However, what is conclusive from these data is that there are opposing inhibitory effects of PIP₂ and excitatory DAG actions on TRPC6 proteins.

An intriguing observation was that OAG reduced association between PIP₂ and TRPC6 proteins without reducing tissue PIP₂ levels. We are cautious in concluding any mechanism from these results although it is tempting to speculate that a simple model of mutual antagonism between DAG and PIP₂, perhaps at a similar binding site(s), may underlie the mechanism by which TRPC6 proteins are activated by DAG via a PKC-independent mechanism. However, it is also possible that DAG may partially activate TRPC6, which in turn allows entry of Ca²⁺ ions leading to subsequent activation of PLC and PIP₂ hydrolysis. Considerably more work including

identification of mutual DAG and PIP₂ binding sites within TRPC6 proteins and investigation of the role of PLC will be required to prove these possible activation mechanisms of TRPC6. Nevertheless, our data suggest that simultaneous depletion of PIP₂ and production of DAG is necessary for activation of native TRPC6 activation as previously proposed for TRP channels in *Drosophila* (Estacion *et al.* 2001; Hardie, 2003).

It is not clear why our data showing that PIP₂ has an inhibitory effect on native TRPC6 channels is opposite to the facilitatory response seen in expressed TRPC6 channels (Kwon *et al.* 2007; Lemonnier *et al.* 2008) although it should be noted that the work of Kwon *et al.* mainly focused on the role of PIP₃ in regulating TRPC6 activity. Lemonnier *et al.* (2008) may have observed potentiation of TRPC6 activity by PIP₂ through the phospholipid acting as a substrate for generation of DAG, particularly if there was an excess of expressed TRPC6 proteins compared with endogenous PIP₂ levels in HEK293 cells. However, our present results may explain why pre-treatment with low doses of carbachol, which are likely to lower PIP₂ levels, enhanced OAG-evoked TRPC activity expressed in HEK293 cells (Estacion *et al.* 2004).

In conclusion, endogenous PIP₂ imposes an inhibitory action on native TRPC6 channels in mesenteric artery myocytes. Removal of this tonic inhibitory effect leads to TRPC6 channel activation indicating that PIP₂ has a dual role of being a precursor for DAG production for channel activation and a direct inhibitor of the ion channel. This may represent an important novel physiological activation mechanism of native TRPC6 channels in vascular myocytes and may also be a common activation mechanism of other TRPC proteins stimulated by DAG via a PKC-independent pathway.

References

- Albert AP & Large WA (2003). Synergism between inositol phosphates and diacylglycerol on native TRPC6-like channels in rabbit portal vein myocytes. *J Physiol* **552**, 789–795.
- Albert AP & Large WA (2006). Signal transduction pathways and gating mechanisms of native TRP-like cation channels in vascular myocytes. *J Physiol* **570**, 45–51.
- Albert AP, Piper AS & Large WA (2005). Role of phospholipase D and diacylglycerol in activating constitutive TRPC-like cation channels in rabbit ear artery myocytes. *J Physiol* **566**, 769–780.
- Albert AP, Pucovsky V, Prestwich SA & Large WA (2006). TRPC3 properties of a native constitutively active Ca²⁺-permeable cation channel in rabbit ear artery myocytes. *J Physiol* **571**, 361–373.
- Albert AP, Saleh SN, Peppiatt-Wildman CM & Large WA (2007). Multiple activation mechanisms of store-operated TRPC channels in smooth muscle cells. *J Physiol* **583**, 25–36.

- Aromolaran AS, Albert AP & Large WA (2000). Evidence for myosin light chain kinase mediating noradrenaline-evoked cation current in rabbit portal vein myocytes. *J Physiol* **524**, 853–863.
- Asteggiano C, Berberian G & Beauge L (2001). Phosphatidylinositol-4,5-bisphosphate bound to bovine cardiac Na⁺/Ca²⁺ exchanger displays a MgATP regulation similar to that of the exchange fluxes. *Eur J Biochem* **268**, 437–442.
- Beauge L, Asteggiano C & Berberian G (2002). Regulation of phosphatidylinositol-4,5-bisphosphate bound to the bovine cardiac Na⁺/Ca²⁺ exchanger. *Ann N Y Acad Sci* **976**, 288–299.
- Beech DJ, Muraki K & Flemming R (2004). Non-selective cationic channels of smooth muscle and the mammalian homologues of *Drosophila* TRP. *J Physiol* **559**, 685–706.
- Bian J, Cui J & McDonaild TV (2001). HERG K⁺ channel activity is regulated by changes in phosphatidylinositol 4,5-bisphosphate. *Circ Res* **89**, 1168–1176.
- Estacion M, Li S, Sinkins WG, Gosling M, Bahra P, Poll C, Westwick J & Schilling WP (2004). Activation of human TRPC6 channels by receptor stimulation. *J Biol Chem* **279**, 22047–22056.
- Estacion M, Sinkins WG & Schilling WP (2001). Regulation of *Drosophila* transient receptor potential-like (TrpL) channels by phospholipase C-dependent mechanisms. *J Physiol* **530**, 1–19.
- Hardie RC (2003). Regulation of TRPC channels by lipid second messengers. *Ann Rev Physiol* **65**, 735–759.
- Helliwell RA & Large WA (1997). α 1-Adrenoceptor activation of a non-selective cation current in rabbit portal vein by 1,2-diacyl-sn-glycerol. *J Physiol* **499**, 417–428.
- Inoue R, Okada T, Onoue H, Hara Y, Shimizu S, Naitoh S, Ito Y & Mori Y (2001). The transient receptor potential protein homologue TRP6 is the essential component of vascular α 1-adrenoceptor-activated Ca²⁺-permeable cation channel. *Circ Res* **88**, 325–332.
- Kwon Y, Hofmann T & Montell C (2007). Integration of phosphoinositide- and calmodulin-mediated regulation of TRPC6. *Mol Cell* **25**, 491–503.
- Large WA (2002). Receptor-operated Ca²⁺-permeable non-selective cation channels in vascular smooth muscle: a physiologic perspective. *J Cardiovasc Electrophysiol* **13**, 493–501.
- Lemonnier L, Trebak M & Putney JW Jr (2008). Complex regulation of the TRPC3, 6 and 7 channel subfamily by diacylglycerol and phosphatidylinositol-4,5-bisphosphate. *Cell Calcium* **43**, 506–514.
- Liou HH, Zhou SS & HuAng CL (1999). Regulation of ROMK1 channel by protein kinase A via a phosphatidylinositol 4,5-bisphosphate-dependent mechanism. *Proc Natl Acad Sci U S A* **96**, 5820–5825.
- Liu X, Bandyopadhyay BC, Singh BB, Groscher K & Ambudkar IS (2005). Molecular analysis of a store-operated and 2-acetyl-sn-glycerol-sensitive non-selective cation channel. Heteromeric assembly of TRPC1-TRPC3. *J Biol Chem* **280**, 21600–21606.
- Ma HP, Saxena S & Warnock DG (2002). Anionic phospholipids regulate native and expressed epithelial sodium channel (ENaC). *J Biol Chem* **277**, 7641–7644.
- Otsuguro K, Tang J, Tang Y, Xiao R, Freichel M, Tsvilovskyy V, Ito S, Flockerzi V, Zhu M & Zholos AV (2008). Isoform-specific inhibition of TRPC4 channel by phosphatidylinositol 4,5-bisphosphate. *J Biol Chem* **283**, 10026–10036.
- Peppiatt-Wildman CM, Albert AP, Saleh SN & Large WA (2007). Endothelin-1 activates a Ca²⁺-permeable cation channel with TRPC3 and TRPC7 properties in rabbit coronary artery myocytes. *J Physiol* **580**, 755–764.
- Pian P, Bucchi A, Robinson RB & Siegelbaum SA (2006). Regulation of gating and rundown of HCN hyperpolarization-activated channels by exogenous and endogenous PIP₂. *J Gen Physiol* **128**, 593–604.
- Rohacs T (2007). Regulation of TRP channels by PIP₂. *Pflugers Arch* **453**, 753–762.
- Saleh SN, Albert AP, Peppiatt CM & Large WA (2006). Angiotensin II activates two cation conductances with distinct TRPC1 and TRPC6 channel properties in rabbit mesenteric artery myocytes. *J Physiol* **577**, 479–495.
- Severson DL & Hee-Cheong M (1989). Diacylglycerol metabolism in isolated aortic smooth muscle cells. *Am J Physiol Cell Physiol* **256**, C11–C17.
- Sours S, Du J, Chu S, Ding M, Zhou XJ & Ma R (2006). Expression of canonical transient receptor potential (TRPC) proteins in human glomerular mesangial cells. *Am J Physiol Renal Physiol* **290**, F1507–F1515.
- Suh B-S & Hille B (2005). Regulation of ion channels by phosphatidylinositol 4,5-bisphosphate. *Curr Opin Neurobiol* **15**, 370–378.
- Tseng PH, Lin HP, Hu H, Wang C, Zhu MX & Chen CS (2004). The canonical transient receptor potential 6 channel as a putative phosphatidylinositol 3,4,5-trisphosphate-sensitive calcium entry system. *Biochemistry* **43**, 11701–11708.
- Voets T & Nilius B (2007). Modulation of TRPs by PIPs. *J Physiol* **582**, 939–944.
- Xie LH, John SA, Ribalet B & Weiss JN (2008). Phosphatidylinositol-4,5-bisphosphate (PIP₂) regulation of strong inward rectifier Kir2.1 channels: multilevel positive cooperativity. *J Physiol* **586**, 1833–1848.
- Yue G, Malik B, Yue G & Eaton DC (2002). Phosphatidylinositol 4,5-bisphosphate (PIP₂) stimulates epithelial sodium channel activity in A6 cells. *J Biol Chem* **277**, 11965–11969.
- Zhang H, Cracium LC, Mirshahi T, Rohacs T, Lopes CM, Jin T & Logothetis DE (2003). PIP₂ activates KCNQ channels, and its hydrolysis underlies receptor-mediated inhibition of M currents. *Neuron* **27**, 963–975.