

Long-Term Interleukin-6 Levels and Subsequent Risk of Coronary Heart Disease: Two New Prospective Studies and a Systematic Review

John Danesh¹*, Stephen Kaptoge¹, Andrea G. Mann¹, Nadeem Sarwar¹, Angela Wood¹, Sara B. Angleman¹, Frances Wensley¹, Julian P. T. Higgins², Lucy Lennon³, Gudny Eiriksdottir^{4,5}, Ann Rumley⁶, Peter H. Whincup⁷, Gordon D. O. Lowe⁶, Vilmundur Gudnason^{4,5}

1 Department of Public Health and Primary Care, University of Cambridge, Cambridge, United Kingdom, **2** Medical Research Council Biostatistics Unit and Public Health Genetics Unit, Cambridge, United Kingdom, **3** Department of Primary Care and Population Sciences, Royal Free University College London Medical School, London, United Kingdom, **4** Icelandic Heart Association, Kopavogur, Iceland, **5** University of Iceland, Kopavogur, Iceland, **6** Division of Cardiovascular and Medical Sciences, University of Glasgow, Glasgow, Scotland, United Kingdom, **7** Division of Community Health Sciences, St George's, University of London, London, United Kingdom

Funding: This study was supported by programme and project grants from the British Heart Foundation (to JD, PHW, GDOL, and VG) and by the Raymond and Beverly Sackler Research Award in the Medical Sciences (to JD). SA is supported by a National Institutes of Health–Cambridge fellowship and her doctoral studies are cosupervised by T. Harris at the National Institute on Aging. Aspects of the study were supported by an unrestricted educational grant from GlaxoSmithKline (to JD). The funding bodies had no role in study design, data collection and analysis, decision to publish or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

Academic Editor: Colin Baigent, University of Oxford, United Kingdom

Citation: Danesh J, Kaptoge S, Mann AG, Sarwar N, Wood A, et al. (2008) Long-term interleukin-6 levels and subsequent risk of coronary heart disease: Two new prospective studies and a systematic review. *PLoS Med* 5(4): e78. doi:10.1371/journal.pmed.0050078

Received: September 6, 2007

Accepted: February 18, 2008

Published: April 8, 2008

Copyright: © 2008 Danesh et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abbreviations: BRHS, British Regional Heart Study; CHD, coronary heart disease; CI, confidence interval; ECG, electrocardiogram; IL-6, interleukin-6; MI, myocardial infarction; SD, standard deviation

* To whom correspondence should be addressed. E-mail: john.danesh@phpc.cam.ac.uk

These authors contributed equally to this work

ABSTRACT

Background

The relevance to coronary heart disease (CHD) of cytokines that govern inflammatory cascades, such as interleukin-6 (IL-6), may be underestimated because such mediators are short acting and prone to fluctuations. We evaluated associations of long-term circulating IL-6 levels with CHD risk (defined as nonfatal myocardial infarction [MI] or fatal CHD) in two population-based cohorts, involving serial measurements to enable correction for within-person variability. We updated a systematic review to put the new findings in context.

Methods and Findings

Measurements were made in samples obtained at baseline from 2,138 patients who had a first-ever nonfatal MI or died of CHD during follow-up, and from 4,267 controls in two cohorts comprising 24,230 participants. Correction for within-person variability was made using data from repeat measurements taken several years apart in several hundred participants. The year-to-year variability of IL-6 values within individuals was relatively high (regression dilution ratios of 0.41, 95% confidence interval [CI] 0.28–0.53, over 4 y, and 0.35, 95% CI 0.23–0.48, over 12 y). Ignoring this variability, we found an odds ratio for CHD, adjusted for several established risk factors, of 1.46 (95% CI 1.29–1.65) per 2 standard deviation (SD) increase of baseline IL-6 values, similar to that for baseline C-reactive protein. After correction for within-person variability, the odds ratio for CHD was 2.14 (95% CI 1.45–3.15) with long-term average (“usual”) IL-6, similar to those for some established risk factors. Increasing IL-6 levels were associated with progressively increasing CHD risk. An updated systematic review of electronic databases and other sources identified 15 relevant previous population-based prospective studies of IL-6 and clinical coronary outcomes (i.e., MI or coronary death). Including the two current studies, the 17 available prospective studies gave a combined odds ratio of 1.61 (95% CI 1.42–1.83) per 2 SD increase in baseline IL-6 (corresponding to an odds ratio of 3.34 [95% CI 2.45–4.56] per 2 SD increase in usual [long-term average] IL-6 levels).

Conclusions

Long-term IL-6 levels are associated with CHD risk about as strongly as are some major established risk factors, but causality remains uncertain. These findings highlight the potential relevance of IL-6–mediated pathways to CHD.

The Editors' Summary of this article follows the references.

Introduction

As atherosclerosis may be an inflammatory condition [1], there is interest in the relevance of various circulating inflammatory markers to cardiovascular diseases [2]. Inflammatory cascades are propagated by proximal mediators such as interleukin-6 (IL-6), a cytokine produced in various tissues. IL-6 exerts proinflammatory effects including stimulation of the liver to produce positive acute-phase proteins during tissue injury or infection [3–5]. Previous epidemiological studies of coronary heart disease (CHD) and inflammation have focused mainly on “downstream” acute phase reactants (e.g., fibrinogen [6] and C-reactive protein [7]) because they are comparatively stable within individuals over time. By contrast, investigation of IL-6 in CHD has been relatively limited because of its shorter half-life (<2 h) and greater within-person variability. Published prospective studies of IL-6 have yielded divergent odds ratios ranging from 1.0 to 3.0 [8–21]. However, as each report has typically comprised only a few hundred patients with CHD, the estimates involve wide, overlapping confidence intervals (CIs). A recent nonquantitative review [22] reported on published data from five prospective studies of IL-6 and CHD, but it comprised only about one-third of the currently published evidence, and it mixed results of studies involving different vascular outcomes and of different study designs. Even a more comprehensive and consistent review of published reports would not, however, have been able to assess appropriately IL-6–CHD associations, because no individual prospective study to our knowledge has yet made allowances for the cytokine’s within-person variability. Owing to fluctuations in IL-6 values over time, comparisons using only baseline values may yield biased estimates of the true association, which can be corrected, for the most part, by using data from paired measurements [23–27] (see Methods).

We report new data on IL-6 levels from two population-based prospective cohorts, the Reykjavik Study and the British Regional Heart Study (BRHS), which together comprise 24,230 predominantly middle-aged individuals with an average of almost 20 years of follow-up per participant. After exclusion of participants with any evidence of baseline coronary disease or stroke, 2,138 incident cases of CHD were available for the present analyses, more than four times as many CHD cases as in the largest previous study [13]. Paired measurements were made in random samples of participants in both studies in order to quantify (and correct for) long-term variation in IL-6 levels. We contextualised our new data by conducting an updated systematic review of all prospective studies of IL-6 and CHD based only in essentially general populations, comprising a further 3,592 CHD cases (defined as myocardial infarction [MI] or death attributed to CHD by World Health Organization [WHO] or similar criteria). Furthermore, we used new data from serial measurements made in our two prospective studies to correct for within-person variability in the studies included in the meta-analysis. The focus of the current report is on the magnitude of the association of IL-6 levels with CHD (rather than on the separate issue of risk prediction).

Methods

Study Population

The Reykjavik Study and the BRHS were initiated in 1967 and 1978, respectively, and have each been described in detail


previously [28,29]. In the Reykjavik Study, men and women born between 1907 and 1935, resident in Reykjavik, Iceland, and its adjacent communities on December 1, 1966, were identified in the national population register and were invited to participate in the study during five stages of recruitment between 1967 and 1991. A total of 8,888 male and 9,681 female participants aged 33–86 y without a history of MI were recruited (72% response rate). Nurses administered questionnaires, made physical measurements, recorded an electrocardiogram (ECG), performed spirometry, and collected fasting venous blood samples to prepare serum that was stored at -20°C for subsequent analysis. History of diabetes was self-reported. All participants have been monitored subsequently for cause-specific mortality (ascertained from central registers on the basis of a death certificate; ICD-9 codes 410–414 were used for CHD death) and for cardiovascular morbidity (diagnosis of MI was based on MONICA criteria), with a loss to follow-up of only about 0.6% to date. The BRHS enrolled 7,735 men aged 40–59 y (response rate 78%) randomly selected from general practice registers in each of 24 British towns during 1978–1980. Nurses administered questionnaires, made physical measurements, recorded an ECG, performed spirometry, and collected nonfasting venous blood samples from 5,661 men in 18 of the towns. Serum was obtained from the baseline blood samples and stored at -20°C for subsequent analysis. Participants were monitored for all-cause mortality (ascertained through the National Health Service central registers, using ICD-9 codes 410–414 for CHD) and for cardiovascular morbidity (using general practitioners’ reports and records, with diagnosis of MI based on WHO criteria), with a loss to follow-up of less than 1% to date. History of diabetes was, again, self-reported. Relevant institutional review boards approved each study, and participants provided informed consent.

Laboratory Measurements

IL-6 measurements were made in the Reykjavik Study and BRHS on available serum samples, using a sensitive enzyme-linked immunosorbent assay (ELISA; R & D Systems) in the Glasgow laboratory by the same technicians blinded to participants’ disease status. The intra-assay coefficient of variability was 7%, and the inter-assay coefficient of variability was 8%. Measurements were made in pairs of samples from a randomly selected subset of baseline participants in each study, comprising 258 participants in the BRHS (samples collected at a median interval of about 4 y apart) and 300 participants in the Reykjavik Study (samples collected at a median interval of about 12 y apart). Because the temperature of storage of the initial BRHS samples was -20°C , whereas it was -70°C at the resurvey, there was scope in this study for slight overestimation of within-person variability in IL-6 levels, encouraging comparison of its findings with those from the Reykjavik Study in which conditions at the two surveys were identical. Lipid and other biochemical measurements involved standard assays, as previously described [28–30].

Statistical Analyses

In both the Reykjavik Study and BRHS, two or three control participants were randomly selected and frequency matched to incident CHD cases in 5-y age bands (and, in the Reykjavik Study, on calendar year of recruitment and sex; and



* In aggregate, these three studies involved a total of approximately 450 CHD cases, or about 7% of the total included in the current meta-analysis

Figure 1. QUOROM Flow Diagram Summarising the Search Strategy for Meta-analysis of IL-6 and CHD Outcomes in Generally Healthy Populations

doi:10.1371/journal.pmed.0050078.g001

in the all-male BRHS by town of residence), thereby obtaining the same joint frequency distributions of these factors among controls and cases. To limit any biases due to preexisting disease, prespecified principal analyses were restricted to the 2,138 individuals with CHD and 4,267 control participants who did not have baseline evidence of CHD or stroke (i.e., without baseline ECG abnormalities or self-reported prevalent MI, stroke, or angina) in a nested case-control comparison. Subsidiary analyses excluded patients who had CHD outcomes in the first 5 y of follow-up. Odds ratios for CHD per 2 standard deviations (SDs) increase in the natural logarithm (\log_e) of IL-6 were calculated using unconditional logistic regression (Stata, version 9). The odds ratios for a 2 SD change is similar to a comparison of extreme thirds of IL-6 values in a population. Analyses to investigate the shape of the IL-6-CHD association were conducted for groups defined by fifths of baseline IL-6 values in controls, with corresponding 95% CIs estimated from floating variances that reflect the amount of information underlying each group (including the reference group) [31–33]. To avoid overadjustment, adjustment for other inflammatory markers (e.g., C-reactive protein) was reserved for subsidiary analyses.

Correction for Regression Dilution

Data from paired measurements in the Reykjavik Study, which coincided with approximately the midpoint of this study's follow-up duration, were used to quantify and to correct for within-person variability in IL-6 and other risk factors, which are likely to be measured with error owing to both within-person biological variability and laboratory variability. Adjusted regression dilution ratios were calculated by Rosner's regression method [26] to quantify the extent to which single measurements of the markers reflect their long term average (i.e., "usual") levels. Paired measure-

ments from the BRHS were used solely for comparison. Statistical significance of the difference in the strength of CHD association with IL-6 compared to other markers was determined from differences in the \log_e odds ratios per 2 SD increase in 2,000 bootstrap samples of the data [34]. Uncertainties in the regression dilution ratios calculated from the 300 participants with repeats in the Reykjavik Study were incorporated into the CIs for CHD associations by adding to the observed regression dilution ratios an error term sampled from the regression dilution ratio distribution on each bootstrap sample. This correction was applied to odds ratios both in the Reykjavik Study and in the BRHS. Correction for multivariate within-person variability in both IL-6 and error-prone confounders was achieved using a multivariable linear regression model [27] from which conditional expectations of usual levels of IL-6 and the error-prone confounders were predicted and then substituted in place of the baseline levels in the logistic model to obtain the corrected odds ratios. Odds ratios for \log_e C-reactive protein, \log_e erythrocyte sedimentation rate, von Willebrand factor, systolic blood pressure, and total cholesterol were calculated using methods described for \log_e IL-6.

Systematic Review

A systematic review of prospective, population-based studies of circulating IL-6 levels and CHD risk was conducted according to QUOROM (Quality of Reporting of Meta-analyses) guidelines (Texts S1 and S2). Studies published before May 2007 were sought using computer-based databases (MEDLINE, EMBASE, and Science Citation Index), and by scanning the reference lists of articles identified for all relevant studies and review articles (including meta-analyses). The computer-based searches combined free and MeSH search terms and combinations of key words related to IL-6 (e.g., "interleukin-6", "IL-6") and coronary disease (e.g., "coronary heart disease", "myocardial ischemia", "myocardial infarction", "CHD", "heart attack", "MI") without restricting language or publication date (flow chart in Figure 1). Data were extracted, where available, on study, geographic location, publication date, sample population, sampling methods (i.e., complete, random, etc.), sample source (plasma/serum), nature of sample (i.e., fresh or frozen, storage temperature), fasting status at time of blood sampling (duration of any fasting), blinding of lab worker to case versus control status, assay type, source and standard, analysis methods and units, case definition, any matching criteria, sample size (of total cohort and numbers with IL-6 measurements), numbers of cases (total, fatal CHD, nonfatal MI), numbers of controls, mean age, sex, time of baseline survey, duration of follow-up, mean or median IL-6 values, adjustment of confounders, and summary statistics. These details were extracted independently by two investigators using a prespecified data extraction form (Table S1). Discrepancies were resolved by discussion and by adjudication of a third reviewer.

Prospective (cohort or "nested" case-control) studies based in essentially general populations (i.e., where participants were not selected on the basis of preexisting disease) with samples taken at baseline (i.e., prior to the occurrence of events) were eligible for inclusion in the current review. Analyses were restricted to studies with at least 1 y of follow-up and with outcomes defined as MI or coronary death. Three studies of IL-6 (comprising approximately 450 CHD cases or

Table 1. Baseline Characteristics of Incident CHD Cases and Controls without Known Evidence of Coronary Disease at Baseline Nested within the Reykjavik Study and BRHS

Type of Marker	Variables	Reykjavik Study				p-Value	BRHS				
		Cases		Controls			Cases		Controls		p-Value
		n	Mean ± SD or n (%)	n	Mean ± SD or n (%)		n	Mean ± SD or n (%)	n	Mean ± SD or n (%)	
Questionnaire	Age (years)	1,768	53.9 ± 8.6	3,303	54.9 ± 8.9	Matched	370	52.1 ± 5.3	964	52.3 ± 5.4	Matched
	Male sex	1,768	1,259 (71%)	3,303	2,281 (69%)	Matched	370	370 (100%)	964	964 (100%)	All male
	Current cigarette/pipe/cigar smoker	1,768	1,068 (60%)	3,303	1,618 (49%)	<0.0001	369	197 (53%)	963	399 (41%)	<0.0001
	Current cigarette smoker	1,767	728 (41%)	3,299	1,055 (32%)	<0.0001	369	197 (53%)	963	399 (41%)	<0.0001
	Self-reported history of diabetes	1,768	42 (2%)	3,303	46 (1%)	0.011	370	9 (2%)	964	13 (1%)	0.164
Physical measurements	Height (m)	1,766	1.71 ± 0.09	3,289	1.72 ± 0.09	0.181	370	1.72 ± 0.06	964	1.73 ± 0.07	0.048
	Weight (kg)	1,762	76.5 ± 13.9	3,290	75 ± 13.5	<0.0001	370	75.9 ± 10.7	964	76.1 ± 11.3	0.765
	Body mass index (kg/m ²)	1,762	26 ± 3.9	3,282	25.4 ± 3.8	<0.0001	370	25.5 ± 3.1	964	25.4 ± 3.3	0.408
	Systolic blood pressure (mm Hg)	1,767	146 ± 24	3,290	142 ± 22	<0.0001	368	152 ± 21	964	146 ± 21	<0.0001
	Diastolic blood pressure (mm Hg)	1,765	91 ± 12	3,288	88 ± 12	<0.0001	368	85 ± 13	964	83 ± 13	0.001
Lipids and metabolic factors	Forced expiratory volume (l/s)	1,744	2.84 ± 0.87	3,226	2.87 ± 0.86	0.305	365	3.18 ± 0.7	950	3.28 ± 0.76	0.033
	Total cholesterol (mmol/l)	1,767	6.88 ± 1.19	3,297	6.40 ± 1.12	<0.0001	369	6.60 ± 1.02	961	6.16 ± 0.99	<0.0001
	High-density lipoprotein cholesterol (mmol/l) ^a	—	—	—	—	—	365	1.10 ± 0.26	945	1.15 ± 0.26	0.001
	Log _e triglycerides (log _e mmol/l)	1,674	0.155 ± 0.451	3,087	0.025 ± 0.439	<0.0001	365	0.645 ± 0.523	954	0.511 ± 0.532	<0.0001
	Log _e lipoprotein(a) (log _e mg/l)	1,764	-1.24 ± 1.61	3,302	-1.59 ± 1.73	<0.0001	368	1.09 ± 1.17	961	0.97 ± 1.06	0.080
Inflammatory markers	Fasting glucose (mmol/l)	1,756	4.6 ± 1.05	3,278	4.53 ± 0.76	0.004	369	5.77 ± 1.69	961	5.61 ± 1.29	0.073
	Log insulin (log _e mU/l) ^a	—	—	—	—	—	369	4.5 ± 0.8	962	4.4 ± 0.8	0.013
	Log _e IL-6 (log _e pg/ml)	1,768	0.78 ± 0.74	3,303	0.65 ± 0.76	<0.0001	370	0.97 ± 0.62	964	0.79 ± 0.63	<0.0001
	Log _e C-reactive protein (log _e mg/l)	1,751	0.49 ± 1.09	3,281	0.22 ± 1.11	<0.0001	290	0.8 ± 1.15	773	0.3 ± 1.2	<0.0001
	Log _e serum amyloid A protein (log _e mg/l) ^a	—	—	—	—	—	290	2.08 ± 0.73	773	1.92 ± 0.68	0.001
Other markers	Log _e erythrocyte sedimentation rate (log _e mm/h) ^b	1,692	1.98 ± 0.97	3,130	1.84 ± 0.96	<0.0001	—	—	—	—	—
	Leukocyte count (×10 ⁹ /l) ^a	—	—	—	—	—	357	7.7 ± 1.8	926	7.2 ± 1.8	<0.0001
	von Willebrand factor (IU/dl)	1,758	115 ± 46	3,283	113 ± 45	0.046	370	115 ± 42	964	111 ± 43	0.153
	Serum creatinine (μmol/l)	1,751	88.9 ± 23.5	3,268	87.8 ± 47.4	0.329	369	98.1 ± 13.4	961	99.1 ± 18.6	0.323
	Uric acid (μmol/l)	1,766	310 ± 71	3,297	299 ± 70	<0.0001	368	357 ± 66	960	350 ± 68	0.098
Other markers	Hemoglobin (mmol/l) ^b	1,752	149 ± 13	3,279	146 ± 13	<0.0001	—	—	—	—	—
	Hematocrit (%)	1,752	44.9 ± 3.5	3,280	44.3 ± 3.5	<0.0001	357	44.8 ± 3.1	926	44 ± 3.2	<0.0001

^aAvailable only in BRHS.^bAvailable only in Reykjavik Study.

doi:10.1371/journal.pmed.0050078.t001

7% of the total) [35–37] were excluded because they reported on composite cardiovascular outcomes rather than for MI or fatal CHD separately. To limit potential biases, analyses involved only within-study comparisons (i.e., cases and controls were directly compared only within each cohort). Results of the studies were combined using a random effects model. Associations of usual levels of IL-6 and CHD were estimated using correction factors derived from the Reykjavik Study and BRHS. Odds and hazard ratios were assumed to approximate the same measure of relative risk. Heterogeneity was quantified by the I^2 statistic [38,39], and subsidiary subgroup analyses were conducted using metaregression [40] to investigate several study-level characteristics potentially explaining heterogeneity, including population sampling

methods, duration of follow-up, source and type of IL-6 assay, blood storage temperature, geographical region, number of CHD cases, sex, degree of adjustment of other cardiovascular risk factors, and whether individuals with CHD at baseline were excluded from analysis. Test statistics from the metaregression based on the F -distribution were used to assess the statistical significance of these study-level characteristics in explaining heterogeneity.

Results

Baseline Correlates of IL-6 Levels

In the Reykjavik study, the median duration of follow-up of participants with CHD and control participants were 14 y and

Table 2. Relative Odds of CHD per 2 SD Increase in Baseline Log_e IL-6 Levels in Participants without Known Coronary Disease at Baseline in the Reykjavik Study and the BRHS


Study	Number of Cases (n)	Number of Controls (n)	Adjusted for Age, Sex, Period, Town, and Smoking Status		Adjusted for Age, Sex, Period, Town, Smoking Status, and Other Established CHD Risk Factors ^a		Adjusted for Age, Sex, Period, Town, Smoking Status, Other Established CHD Risk Factors, and CRP ^b	
			Odds Ratio (95% CI)	χ^2	Odds Ratio (95% CI)	χ^2	Odds Ratio (95% CI)	χ^2
Reykjavik Study	1,768	3,303	1.48 (1.31–1.68)	—	1.38 (1.22–1.57)	—	1.24 (1.07–1.42)	—
BRHS	370	964	2.12 (1.58–2.84)	—	1.94 (1.43–2.63)	—	1.51 (1.00–2.27)	—
Combined	2,138	4,267	1.57 (1.40–1.76)	59	1.45 (1.29–1.63)	40	1.26 (1.10–1.44)	11

Comparisons are reported per 2 SD increase. The SD for baseline log_e IL-6 values in the two studies combined was 0.74 log pg/ml, therefore 2 SD increase in log_e IL-6 corresponds to approximate quadrupling of IL-6 levels since $e^{2 \times 0.74} = 4.4$. Wald χ^2 statistics of whether IL-6 is significantly associated with CHD are presented for the combined analyses.

^aThe adjusted odds ratios for CHD in the combined analysis were 1.94 (1.60, 2.34) for the top fifth versus the bottom fifth, and 1.72 (1.46, 2.04) for the top quarter versus the bottom quarter of IL-6 levels.

^bBecause of missing values, the model with further adjustment for C-reactive protein (CRP) levels involved 1,935 CHD cases and 3,798 controls. Individuals with evidence of electrocardiogram abnormalities, previous MI, or history of angina at the baseline survey were excluded from these analyses. Period refers to 5-y calendar periods of recruitment; town refers to town of recruitment; established CHD risk factors included systolic blood pressure, total cholesterol, log_e triglycerides, body mass index, and diabetes history. Smoking status refers to current, former, and never smoking, further adjustment for smoking amount (i.e., cigarettes smoked per day) did not materially change the odds ratios.

doi:10.1371/journal.pmed.0050078.t002

**Figure 2.** Risk of CHD by Fifths of Baseline Circulating IL-6 Levels in a Pooled Analysis of Participants without Known CHD at Baseline in the Reykjavik Study and the BRHS

Fifths were calculated on the basis of the distribution of controls in the combined dataset. CIs were calculated using floating variances. Sizes of data markers are proportional to the inverse of the variance of the odds ratios. Adjusted for cohort, age, sex, recruitment period, and town of recruitment.

doi:10.1371/journal.pmed.0050078.g002

23 y, respectively; in the BRHS, the corresponding durations were 9 y and 17 y, respectively. IL-6 levels were significantly higher in those with CHD than in control individuals in both the Reykjavik Study and in the BRHS, as were levels of several established cardiovascular risk factors (Table 1). Mean IL-6 levels were slightly higher in the male-only BRHS compared with the mixed-sex Reykjavik study. As shown in Table S2, baseline IL-6 levels were positively and highly significantly ($p < 0.0001$) associated with age, smoking, systolic blood pressure, and concentrations of triglycerides, fasting glucose, and several inflammatory markers (including C-reactive protein and the erythrocyte sedimentation rate), and somewhat less strongly with body mass index ($p < 0.001$). IL-6 levels were inversely associated with total cholesterol concentrations ($p < 0.01$). With the exception of correlations of IL-6 with smoking and measured inflammatory markers (for which the Pearson coefficient generally ranged between 0.20 and 0.43), even the highly significant correlations of IL-6 with other characteristics tended to be relatively small in magnitude.

Baseline IL-6 Levels and Incident CHD

Table 2 provides odds ratios for CHD per 2 SD increase in baseline log_e IL-6 levels. The minimally adjusted odds ratios (95% CI) for CHD per 2 SD increase were 1.48 (1.31–1.68) in the Reykjavik Study, 2.12 (1.58–2.84) in the BRHS, and 1.57 (1.40–1.76) in the combined dataset. After further adjustment for smoking status and several other established cardiovascular risk factors, the odds ratio per 2 SD increase fell to 1.38 (1.21–1.57) in the Reykjavik Study, 2.08 (1.51–2.87) in the BRHS, and 1.46 (1.29–1.65) in the combined dataset. Odds ratios did not change materially after exclusion of CHD outcomes recorded during the first 5 y of follow-up (odds ratio [95% CI] per 2 SD increase: 1.41 [1.24–1.60]) or in analyses including individuals with prevalent CHD at baseline (1.55 [1.38–1.73]). CHD risk increased continuously with increasing fifths of circulating IL-6 concentrations (Figure 2). Odds ratios did not vary substantially by levels of several

Table 3. Regression Dilution Ratios (95% CI) for IL-6, Other Measured Inflammatory Markers, and Some Established Cardiovascular Risk Factors, Based on Paired Measurements in Up to 380 Participants in the Reykjavik Study Who Provided Samples Approximately 12 Years Apart

Markers and Risk Factors	Variables	Number of Participants Providing Paired Measurements	Regression Dilution Ratio ^a (95% CI)
Inflammatory markers	Log _e IL-6 (log _e pg/ml)	300	0.35 (0.23–0.48) ^b
	Log _e C-reactive protein (log _e mg/l)	370	0.54 (0.44–0.64)
	von Willebrand factor (IU/dl)	378	0.55 (0.45–0.65)
	Log _e erythrocyte sedimentation rate(log _e mm/h)	331	0.64 (0.51–0.76)
Established risk factors	Systolic blood pressure (mm Hg)	380	0.65 (0.54–0.76)
	Total cholesterol (mmol/l)	379	0.59 (0.51–0.67)
	Body mass index (kg/m ²)	378	0.90 (0.83–0.97)
	Smoking status (current versus never/former)	380	0.59 (0.49–0.68)

^aRegression dilution ratio calculated from paired measurements in the Reykjavik Study using Rosner's regression method, adjusted for baseline age, sex, smoking status, diabetes history, total cholesterol, log_e triglycerides, systolic blood pressure, and body mass index.

^b $p < 0.05$ in comparison of the regression dilution ratio for log_e IL-6 versus the regression dilution ratios for each of the other markers in the table.

doi:10.1371/journal.pmed.0050078.t003

established cardiovascular risk factors, except potentially by sex ($p = 0.03$) and cholesterol levels ($p = 0.02$), though the latter were only marginally significant findings in the context of multiple comparisons (Figure S1).

Correction of Odds Ratios for Within-Person Variability

The regression dilution ratio for log_e IL-6 was 0.41 (0.28–0.53) in 258 individuals who provided paired measurements on average 4 y apart in the BRHS, similar to the regression dilution ratio of 0.35 (0.23–0.48) in 300 individuals who provided paired measurements on average 12 y apart in the Reykjavik Study (mean log_e IL-6 levels at baseline and at resurvey were 0.76 [SD 0.58] and 0.98 [SD 0.64] pg/ml, respectively, in BRHS and 0.63 [SD 0.70] and 0.88 [SD 0.68] pg/ml, respectively, in the Reykjavik Study). Similar point

estimates for the regression dilution ratios of log_e IL-6 levels over 4- and 12-y intervals may reflect that much of the variability in this marker is related to shorter-term rather than longer-term factors, although CIs around these estimates were relatively wide. As there were no significant associations between within-person differences in log_e IL-6 levels and time intervals between measurements ($p = 0.53$), it is unlikely that increasing age of participants explains the generally higher mean log_e IL-6 levels at resurvey. Table 3 indicates that the degree of within-person variability for IL-6 levels over 12 y was generally greater than those for other measured inflammatory markers and established cardiovascular risk factors ($p < 0.05$ for each comparison). Figure 3 (with numeric details shown in Table S3) shows three odds




Figure 3. Odds Ratios for CHD per 2 SD Increase in Several Inflammatory Markers and Established Cardiovascular Risk Factors Shown with Progressively Increasing Degrees of Correction for Within-person Variability in a Pooled Analysis of Participants without Known Coronary Disease at Baseline in the Reykjavik Study and the BRHS

(A) Baseline exposure adjusted for baseline confounders.

(B) Usual exposure adjusted for baseline confounders.

(C) Usual exposure adjusted for usual confounders.

Sizes of data markers are proportional to the inverse of the variance of the odds ratios. Baseline (A) refers to measured values just at the initial examination. Usual (B and C) refers to an estimate of the long-term average values by employing data on repeat measurements from the Reykjavik Study (see Methods). Odds ratios are adjusted for cohort, age, sex, period and town of recruitment, smoking status, history of diabetes, total cholesterol, log_e triglycerides, systolic blood pressure, and body mass index.

doi:10.1371/journal.pmed.0050078.g003

ratios for \log_e IL-6 and several other risk markers: (i) assessment 1 lists the odds ratio per 2 SD increase in baseline levels of the relevant risk marker adjusted for baseline levels of established risk factors (i.e., uncorrected for within-person variability in either exposure or confounder levels) (Figure 3A); (ii) assessment 2 lists the odds ratio per 2 SD increase in usual levels of the relevant risk marker adjusted for baseline levels of established risk factors (i.e., corrected for within-person variability only in levels of the exposure) (Figure 3B); and (iii) assessment 3 lists the odds ratio per 2 SD increase in usual levels of the relevant risk marker adjusted for usual levels of established risk factors (i.e., corrected for within-person variability in levels of both exposure and confounders) (Figure 3C). In assessment 1, the odds ratio for \log_e IL-6 is similar to those for \log_e C-reactive protein and systolic blood pressure, but lower than that for total cholesterol. In assessment 3, the odds ratio for \log_e IL-6 is comparable to those for some established risk factors. Figure 3 shows that the impact on odds ratios of correction for within-person variability in confounder levels varied in magnitude among risk markers, reflecting the markers' differing strength of associations with levels of confounders (and the differing degrees of within-person variability among such confounders [Table S3]). Sensitivity analyses allowing for uncertainty in the regression dilution ratios for the exposure variables (but not in the usual confounder levels, Table S3 legend) gave slightly wider CIs for the corrected associations.

Updated Meta-analysis

We identified 17 relevant studies [8–21] (including the two current studies, ten newly identified studies [8–13,17,19–21], and five studies [14–16,18] included in the previous published review [22]; flow chart in Figure 1). These studies comprised a total of 5,730 patients with MI or CHD death and 19,038 control individuals (weighted mean age at entry, 62 y; weighted mean follow-up, 6 y; Table 4). All studies, with one exception [20], used enzyme-linked immunosorbent assays produced by the same manufacturer, and all but two studies [10,17] adjusted for age, sex, and established cardiovascular risk factors. Using only within-study comparisons, the combined odds ratio (95% CI) was 1.61 (1.42–1.83) per 2 SD increase of baseline IL-6 levels and 3.34 (2.45–4.56) per 2 SD increase of usual IL-6 levels (Figure 4). Alternatively expressed, the odds ratios per 1 SD increase in baseline and usual IL-6 levels were 1.26 (1.19–1.35) and 1.83 (1.56–2.14), respectively. There was evidence of moderate heterogeneity among the 17 separate findings ($I^2 = 58\%$, 95% CI 28%–75%), but, as displayed in Figure S2, little of it was explained by differences in sample size ($F_{1,15} = 0.91$, $p = 0.36$), geographical location ($F_{1,15} = 0.30$, $p = 0.59$), population source ($F_{2,14} = 0.76$, $p = 0.49$), whether individuals with prevalent CHD at baseline were excluded from subsequent analyses ($F_{1,15} = 2.08$, $p = 0.17$), sex ($F_{2,11} = 0.16$, $p = 0.85$), reported duration of follow-up ($F_{1,15} = 1.41$, $p = 0.25$), or temperature of blood storage ($F_{3,13} = 1.19$, $p = 0.35$).

Discussion

Previous reports on IL-6 levels and CHD have not been able to correct for within-person fluctuations in the levels of this short-acting cytokine, potentially yielding biased estimates. The present study, which made such correction on the

basis of paired measurements of IL-6, indicates that long-term average (“usual”) IL-6 levels are about as strongly associated with CHD risk as are some major established risk factors. Increasing IL-6 levels are associated with progressively increasing odds ratios for CHD (i.e., there are continuous, approximately log-linear relationships). There are moderate associations of IL-6 levels with some established risk factors (notably smoking, diabetes, and dyslipidemia) and with several downstream inflammatory markers, consistent with the key role of IL-6 in mediating inflammatory cascades [3–5,41–44]. The current findings do not, of course, establish causality, but they may have implications for understanding disease mechanisms and for further research strategies.

By showing that pathways mediated by long-term IL-6 levels are associated with CHD risk about as strongly as are some major established risk factors, the current data reinforce interest in the connection between inflammatory pathways and cardiovascular diseases. The data also underscore the need for investigations of proximal inflammatory mediators that can quantify and correct for within-person variability. Serious underestimation is likely without such correction because, as demonstrated in the current study, the variability in IL-6 levels is substantially higher than for downstream inflammatory markers and several established CHD risk factors. Hence, given the central role of IL-6 levels in inflammatory pathways and its continuous association with CHD risk, it warrants further investigation as a plausible potential therapeutic target. There are initial reports of reductions in circulating IL-6 concentrations in randomised trials of statins, although these agents have other effects, notably lowering low-density lipoprotein cholesterol [45–47]. Future investigations should involve complementary strategies to help judge causality, such as large-scale studies of specific genetic markers as proxies for circulating IL-6 levels [48] (although it is now uncertain whether the -174 G/C *IL-6* polymorphism is materially correlated with IL-6 levels [49]) and, possibly, use of selective IL-6 antagonists in early randomized trials [50], although the pleiotropic actions of IL-6 could complicate such an approach.

The strengths and potential limitations of the present report merit careful consideration. The current study reports new data from two population-based prospective cohorts comprising more than four times as many patients with first-ever CHD than in the previous largest report. Both the Reykjavik Study and BRHS identified participants in population registers, involved high response and follow-up rates, and involved robust ascertainment of incident MI and coronary death. Their broadly similar results support the generalisability of the current findings. Assay methods used in the current study were similar to those in earlier reports, yielding similar median IL-6 levels (Table 4). Although the current studies involved more prolonged blood storage than previous studies, they produced associations with CHD risk at least as strong as in earlier studies, arguing against underestimation due to sample degradation. Paired IL-6 (and other) measurements enabled approximate correction for within-person variability in a way that implies that long-term average IL-6 values are relevant to CHD risk, consistent with prevailing hypotheses of sustained low-grade inflammation, but it is not clear why mean \log_e IL-6 levels were generally higher at resurvey examinations.

In comparison with a previous nonquantitative review [22],

Table 4. Characteristics of 17 Prospective Studies of Circulating IL-6 Concentrations and CHD Risk in Essentially General Populations

Study	Location	Population/Sampling Method	Baseline Survey	Number of Cases of CHD	Number of Controls	Age Range (y)	Male Sex (%)	Mean Duration of Follow-up (y)	IL-6 Assay ^a		Blood Storage Temperature (°C)	Median (IQR) IL-6 in Controls (pg/ml)
									Source	Type		
Reykjavik ^b	Iceland	Population register/complete birth cohorts	1967–1991	1,768	3,303	33–86	69	20	R&D Systems	ELISA	–20	1.84 (1.22–2.87)
BRHS ^b	UK	General practitioners' list/random	1978–1980	370	964	40–59	100	16	R&D Systems	ELISA	–20	2.09 (1.48–3.30)
WOSCOPS [13]	Scotland	Coronary screening clinic/complete	1989–1991	503	916	45–64	100	5	R&D Systems	ELISA	–70	1.77 (1.20–2.76)
MONICA/KORA [11]	Germany	Population register/FU questionnaire	1984–2002	382	1,980	35–74	55	11	R&D Systems	ELISA	–80	2.1 (1.0)/1.9 (1.0)
WHIOS [16]	US	Trial screenings/complete	1994–1998	304	304	50–79	0	3 (median)	R&D Systems	ELISA	–70	1.47 (1.05–2.15)
ARIC [9]	US	Four U.S. communities/complete	1987–1989	304	365	45–74	~56	4.6	R&D Systems	ELISA	–70	1.63 (0.6)
HPFS [15]	US	Workforce/complete	1986	265	529	40–75	100	NS	R&D Systems	ELISA	–130	1.53 (0.98–2.88)
NHS [15]	US	Workforce/complete	1976	239	469	30–55	0	NS	R&D Systems	ELISA	–130	1.65 (1.15–2.65)
CHS ^c [10]	US	Medicare eligibility lists/random	1989–1990, 1992–1993	217	1,650	≥65	43	6.5	R&D Systems	ELISA	–70	1.90 ± 1.85 ^d
QCS [19]	Canada	Seven towns in Quebec	1985	210	1,772	35–64	100	13	R&D Systems	ELISA	–80	1.0 (2.0)
PHS [18]	US	Workforce/complete	1982–1984	202	202	40–84	100	14	R&D Systems	ELISA	–80	1.46 (range: 0.015–10.01)
EAS [21]	Scotland	General practitioners' list	1987	190	1,177	55–74	51	12	R&D Systems	ELISA	–50	2.01 (1.34–3.28)
Health ABC [8]	US	Medicare beneficiaries list Pittsburgh/Memphis	1997	188	1,950	70–79	45	3.6	R&D Systems	ELISA	–70	1.68 (1.16–2.54)
PRIME [14]	France, Northern Ireland	Workforce and community setting/complete	1991–1993	163	763	50–59	100	5	R&D Systems	ELISA	–80	1.25 (0.84–1.98)
RANCHO [12]	US	Residential community/complete	1972–1974	152	1,909	40–94	46	20	R&D Systems	ELISA	–70	2.12 (1.99–2.25)
FINRISK [20]	Finland	Population registers in four regions	1992–2001	151	541	25–46	64	9	Immulite chemiluminescent immunometric assay	Solid-phase	–20	M 1.31 (1.0–1.7); F 1.38 (0.9–2.0)
WHS [17]	US	Workforce/complete	1992–1995	122	244	>45	0	3	R&D Systems	ELISA	–70	1.30 (1.00–2.03)

^aThe BRHS, Reykjavik Study, and WOSCOPS used the R&D Systems standard for IL-6 (i.e., 50 pg recombinant human IL-6 in buffered protein base with preservatives, lyophilized). Such specific information on standards used was not available from the other studies.

^bCurrent studies.

^cControls in this study were randomly selected cohort members free of prevalent or incident cardiovascular disease and free of MRI-detectable infarcts, although some did not satisfy additional strict criteria of being free of subclinical cardiovascular disease established by detailed clinical investigations, including measurement of carotid intima-media thickness and assessment of abnormalities in wall motion on an echocardiogram.

^dGeometric mean of natural log transformed values, range.

ARIC, Artherosclerosis Risk in Communities; CHS, Cardiovascular Health Study; EAS, Edinburgh Artery Study; FINRISK, Finnish National Risk Factor Survey; Health ABC, Health, Aging, and Body Composition; HPFS, Health Professionals Follow-up Study; IQR, interquartile range; MONICA, MONITORing of trends and determinants in Cardiovascular disease; NHS, Nurses' Health Study; PHS, Physicians' Health Study; PRIME, Prospective Epidemiological Study of Myocardial Infarction; QCS, Quebec Cardiovascular Study; RANCHO, Rancho Bernardo Study; WHIOS, Women's Health Initiative Observational Study; WHS, Women's Health Study; WOSCOPS, West of Scotland Coronary Prevention Study.

doi:10.1371/journal.pmed.0050078.t004






Figure 4. Summary of 17 Prospective Studies Reporting Adjusted Associations of CHD Risk per 2 SD Increase in Baseline IL-6 Levels and Usual IL-6 Levels, with Both Adjusted Only for Baseline Levels of Several Established Risk Factors

(A) Baseline exposure adjusted for baseline confounders.

(B) Usual exposure adjusted for baseline confounders.

Baseline refers to measured values just at the initial examination. Usual refers to an estimate of the long-term average values. The correction for regression dilution (see Methods) in the two current studies (filled squares) was done using study-specific regression dilution ratios of 0.35 in Reykjavik study and 0.41 in the BRHS. A regression dilution ratio of 0.4 was assumed for other studies (empty squares) where such regression dilution ratio coefficients were unavailable. The odds ratio for a 1 SD increase in baseline and usual IL-6 levels are 1.26 (1.19–1.35) and 1.83 (1.56–2.14), respectively. Unshaded data markers were used in the second plot of this figure to indicate that correction for regression dilution involved use of coefficients from external studies, in contrast with the cohort-specific data on IL-6 variability available for the Reykjavik Study and BRHS. Sizes of data markers indicate the weight of each study in the analysis. Adjustment for within-person variability in levels of IL-6 and possible confounders requires access to individual participant data, currently available only in the Reykjavik Study and BRHS.

+, adjusted for age and sex; ++, adjusted for age, sex, and smoking status; +++, adjusted for age, sex, smoking status, and other established CHD risk factors; FINRISK, Finnish National Risk Factor Survey; HPFUS, Health Professionals' Follow-up Study; MONICA, MONITORing of trends and determinants in Cardiovascular disease; PRIME, Prospective Epidemiological Study of Myocardial Infarction; WHIOS, Women's Health Initiative Observational Study; WOSCOPS, West of Scotland Coronary Prevention Study.

doi:10.1371/journal.pmed.0050078.g004

the current meta-analysis involved almost five times as many incident CHD cases from population-based prospective studies (thereby enhancing statistical power), strictly defined CHD as MI or fatal CHD (thereby enhancing accuracy of disease classification), excluded (where possible) people with prevalent cardiovascular disease (thereby limiting potential biases), and explored in detail potential sources of heterogeneity (thereby probing possible causes of study diversity). This updated review suggests that the apparently divergent estimates reported in earlier studies were probably chiefly due to random error. Further studies are needed to test whether associations of IL-6 with CHD are importantly modified by lipid concentrations, as suggested by exploratory analyses in the current report. Studies are also needed with serial measurements in larger numbers of participants over different intervals in order to assess IL-6 variability in greater detail (e.g., assessment of any changes in mean levels and variability over time). This information will enable study-specific and time-dependent correction for regression dilution [23,51,52], whereas in the current meta-analysis the correction factor was derived from repeat measurements in only the Reykjavik Study. Adjustment for within-person variability in both exposure and confounder levels across all available studies will require access to individual participant

data, as demonstrated by such adjustment in the current analyses of the Reykjavik Study and BRHS.

Conclusion

Long-term IL-6 levels are associated with CHD risk about as strongly as are some major established risk factors, but causality remains uncertain. These findings highlight the potential relevance of IL-6-mediated pathways to CHD.

Supporting Information

Figure S1. Association of Baseline Circulating Log_e IL-6 Levels and CHD Risk in a Pooled Analysis of Participants without Known CHD at Baseline in The Reykjavik Study and the BRHS, Grouped by Several Participant Level Characteristics

Found at doi:10.1371/journal.pmed.0050078.sg001 (60 KB PPT).

Figure S2. Seventeen Prospective Studies of Baseline IL-6 levels and Risk of CHD, Grouped According to Several Study Characteristics

Found at doi:10.1371/journal.pmed.0050078.sg002 (8 KB PPT).

Table S1. Data Abstraction Form for Systematic Review of IL-6 and CHD

Found at doi:10.1371/journal.pmed.0050078.st001 (49 KB DOC).

Table S2. Comparison of Levels of Risk Factors and Other Characteristics by Thirds of Baseline IL-6 Levels among Non-CHD cases in a Pooled Analysis of the Reykjavik Study and the BRHS

Found at doi:10.1371/journal.pmed.0050078.st002 (103 KB DOC).

Table S3. Comparisons of Odds Ratios for CHD with Baseline and Usual (i.e., Corrected for Within-person Variability) Levels of Four Circulating Inflammatory Markers and Two Established Cardiovascular Risk Factors in a Pooled Analysis of Data for Participants without Known Coronary Disease at Baseline in the Reykjavik Study and the BRHS

Found at doi:10.1371/journal.pmed.0050078.st003 (41 KB DOC).

Text S1. Protocol for Systematic Review of Prospective Studies of IL-6 and Coronary Disease

Found at doi:10.1371/journal.pmed.0050078.sd001 (33 KB PDF).

Text S2. The QUOROM Statement Checklist

Found at doi:10.1371/journal.pmed.0050078.sd002 (48 KB DOC).

Acknowledgments

Mark B. Pepys, Kausik Ray, and Emanuele Di Angelantonio commented helpfully; J. Wheeler provided statistical support; E. Poorhang and P. Welsh conducted IL-6 assays.

Author contributions. JD, PHW, and VG conceived and designed the current work. SK and FW analysed the data. JD and SK drafted the paper. JD, SK, NS, AW, SBA, FW, JPTH, GE, AR, PHW, GDOL, and VG contributed to the writing of the paper. SK, AGM, NS, FW, and JPTH conducted the systematic review. LL collected the follow-up data in the BRHS. AR and GDOL supervised IL-6 assays.

References

- Hansson GK (2005) Inflammation, atherosclerosis, and coronary artery disease. *N Engl J Med* 352: 1685–1695.
- Pearson TA, Mensah GA, Alexander RW, Anderson JL, Cannon RO III, et al. (2003) Markers of inflammation and cardiovascular disease: application to clinical and public health practice: a statement for healthcare professionals from the Centers for Disease Control and Prevention and the American Heart Association. *Circulation* 107: 499–511.
- Kerr R, Stirling D, Ludlam CA (2001) Interleukin 6 and haemostasis. *Br J Haematol* 115: 3–12.
- Le JM, Vilcek J (1989) Interleukin 6: a multifunctional cytokine regulating immune reactions and the acute phase protein response. *Lab Invest* 61: 588–602.
- Woodward M, Rumley A, Tunstall-Pedoe H, Lowe GD (1999) Associations of blood rheology and interleukin-6 with cardiovascular risk factors and prevalent cardiovascular disease. *Br J Haematol* 104: 246–257.
- Meade TW, Ruddock V, Stirling Y, Chakrabarti R, Miller GJ (1993) Fibrinolytic activity, clotting factors, and long-term incidence of ischaemic heart disease in the Northwick Park Heart Study. *Lancet* 342: 1076–1079.
- Kuller LH, Tracy RP, Shaten J, Meilahn EN (1996) Relation of C-reactive protein and coronary heart disease in the MRFIT nested case-control study. *Multiple Risk Factor Intervention Trial*. *Am J Epidemiol* 144: 537–547.
- Cesari M, Penninx BW, Newman AB, Kritchevsky SB, Nicklas BJ, et al. (2003) Inflammatory markers and onset of cardiovascular events: results from the Health ABC study. *Circulation* 108: 2317–2322.
- Folsom AR, Chambless LE, Ballantyne CM, Coresh J, Heiss G, et al. (2006) An assessment of incremental coronary risk prediction using C-reactive protein and other novel risk markers: the atherosclerosis risk in communities study. *Arch Intern Med* 166: 1368–1373.
- Jenny NS, Tracy RP, Ogg MS, Luong LA, Kuller LH, et al. (2002) In the elderly, interleukin-6 plasma levels and the $-174G>C$ polymorphism are associated with the development of cardiovascular disease. *Arterioscler Thromb Vasc Biol* 22: 2066–2071.
- Koenig W, Khuseynova N, Baumert J, Thorand B, Loewel H, et al. (2006) Increased concentrations of C-reactive protein and IL-6 but not IL-18 are independently associated with incident coronary events in middle-aged men and women: results from the MONICA/KORA Augsburg case-cohort study, 1984–2002. *Arterioscler Thromb Vasc Biol* 26: 2745–2751.
- Langenberg C, Bergstrom J, Scheidt-Nave C, Pfeilschifter J, Barrett-Connor E (2006) Cardiovascular death and the metabolic syndrome: role of adiposity-signaling hormones and inflammatory markers. *Diabetes Care* 29: 1363–1369.
- Lowe GD, Rumley A, McMahon AD, Ford I, O'Reilly DS, et al. (2004) Interleukin-6, fibrin D-dimer, and coagulation factors VII and XIIa in prediction of coronary heart disease. *Arterioscler Thromb Vasc Biol* 24: 1529–1534.
- Luc G, Bard JM, Juhan-Vague I, Ferrieres J, Evans A, et al. (2003) C-reactive protein, interleukin-6, and fibrinogen as predictors of coronary heart disease: the PRIME Study. *Arterioscler Thromb Vasc Biol* 23: 1255–1261.
- Pai JK, Pischon T, Ma J, Manson JE, Hankinson SE, et al. (2004) Inflammatory markers and the risk of coronary heart disease in men and women. *N Engl J Med* 351: 2599–2610.
- Pradhan AD, Manson JE, Rossouw JE, Siscovick DS, Mouton CP, et al. (2002) Inflammatory biomarkers, hormone replacement therapy, and incident coronary heart disease: prospective analysis from the Women's Health Initiative observational study. *JAMA* 288: 980–987.
- Ridker PM, Hennekens CH, Buring JE, Rifai N (2000) C-reactive protein and other markers of inflammation in the prediction of cardiovascular disease in women. *N Engl J Med* 342: 836–843.
- Ridker PM, Rifai N, Stampfer MJ, Hennekens CH (2000) Plasma concentration of interleukin-6 and the risk of future myocardial infarction among apparently healthy men. *Circulation* 101: 1767–1772.
- St-Pierre AC, Cantin B, Bergeron J, Pirro M, Dagenais GR, et al. (2005) Inflammatory markers and long-term risk of ischemic heart disease in men: A 13-year follow-up of the Quebec Cardiovascular Study. *Atherosclerosis* 182: 315–321.
- Tuomisto K, Jousilahti P, Sundvall J, Pajunen P, Salomaa V (2006) C-reactive protein, interleukin-6 and tumor necrosis factor alpha as predictors of incident coronary and cardiovascular events and total mortality. A population-based, prospective study. *Thromb Haemost* 95: 511–518.
- Tzoulaki I, Murray GD, Lee AJ, Rumley A, Lowe GD, et al. (2007) Relative value of inflammatory, hemostatic, and rheological factors for incident myocardial infarction and stroke: the Edinburgh Artery Study. *Circulation* 115: 2119–2127.
- Lobbes MB, Lutgens E, Heeneman S, Cleutjens KB, Kooi ME, et al. (2006) Is there more than C-reactive protein and fibrinogen? The prognostic value of soluble CD40 ligand, interleukin-6 and oxidized low-density lipoprotein with respect to coronary and cerebral vascular disease. *Atherosclerosis* 187: 18–25.
- Clarke R, Shipley M, Lewington S, Youngman L, Collins R, et al. (1999) Underestimation of risk associations due to regression dilution in long-term follow-up of prospective studies. *Am J Epidemiol* 150: 341–353.
- Fibrinogen Studies Collaboration, Wood AM, White I, Thompson SG, Lewington S, et al. (2006) Regression dilution methods for meta-analysis: assessing long-term variability in plasma fibrinogen among 27,247 adults in 15 prospective studies. *Int J Epidemiol* 35: 1570–1578.
- Knuiman MW, Divitini ML, Buzas JS, Fitzgerald PE (1998) Adjustment for regression dilution in epidemiological regression analyses. *Ann Epidemiol* 8: 56–63.
- Rosner B, Willett WC, Spiegelman D (1989) Correction of logistic regression relative risk estimates and confidence intervals for systematic within-person measurement error. *Stat Med* 8: 1051–1069.
- Rosner B, Spiegelman D, Willett WC (1990) Correction of logistic regression relative risk estimates and confidence intervals for measurement error: the case of multiple covariates measured with error. *Am J Epidemiol* 132: 734–745.
- Jonsdottir LS, Sigfusson N, Gudnason V, Sigvaldason H, Thorgeirsson G (2002) Do lipids, blood pressure, diabetes, and smoking confer equal risk of myocardial infarction in women as in men? The Reykjavik Study. *J Cardiovasc Risk* 9: 67–76.
- Walker M, Whincup PH, Shaper AG (2004) The British Regional Heart Study 1975–2004. *Int J Epidemiol* 33: 1185–1192.
- Danesh J, Wheeler JG, Hirschfield GM, Eda S, Eiriksdottir G, et al. (2004) C-reactive protein and other circulating markers of inflammation in the prediction of coronary heart disease. *N Engl J Med* 350: 1387–1397.
- Easton DF, Peto J, Babiker AG (1991) Floating absolute risk: an alternative to relative risk in survival and case-control analysis avoiding an arbitrary reference group. *Stat Med* 10: 1025–1035.
- Greenland S, Michels KB, Robins JM, Poole C, Willett WC (1999) Presenting statistical uncertainty in trends and dose-response relations. *Am J Epidemiol* 149: 1077–1086.
- Plummer M (2004) Improved estimates of floating absolute risk. *Stat Med* 23: 93–104.
- Carpenter J, Bithell J (2000) Bootstrap confidence intervals: when, which, what? A practical guide for medical statisticians. *Stat Med* 19: 1141–1164.
- Harris TB, Ferrucci L, Tracy RP, Corti MC, Wacholder S, et al. (1999) Associations of elevated interleukin-6 and C-reactive protein levels with mortality in the elderly. *Am J Med* 106: 506–512.
- Kip KE, Marroquin OC, Shaw LJ, Arant CB, Wessel TR, et al. (2005) Global inflammation predicts cardiovascular risk in women: a report from the Women's Ischemia Syndrome Evaluation (WISE) study. *Am Heart J* 150: 900–906.
- Volpato S, Guralnik JM, Ferrucci L, Balfour J, Chaves P, et al. (2001) Cardiovascular disease, interleukin-6, and risk of mortality in older women: the women's health and aging study. *Circulation* 103: 947–953.
- Higgins JP, Thompson SG (2002) Quantifying heterogeneity in a meta-analysis. *Stat Med* 21: 1539–1558.
- Higgins JP, Thompson SG, Deeks JJ, Altman DG (2003) Measuring inconsistency in meta-analyses. *BMJ* 327: 557–560.
- Thompson SG, Sharp SJ (1999) Explaining heterogeneity in meta-analysis: a comparison of methods. *Stat Med* 18: 2693–2708.
- Pickup JC (2004) Inflammation and activated innate immunity in the pathogenesis of type 2 diabetes. *Diabetes Care* 27: 813–823.
- Ritchie SA, Connell JM (2007) The link between abdominal obesity, metabolic syndrome and cardiovascular disease. *Nutr Metab Cardiovasc Dis* 17: 319–326.
- Szekanecz Z, Kerekes G, Der H, Sandor Z, Szabo Z, et al. (2007) Accelerated atherosclerosis in rheumatoid arthritis. *Ann N Y Acad Sci* 1108: 349–358.

44. Yanbaeva DG, Dentener MA, Creutzberg EC, Wesseling G, Wouters EF (2007) Systemic effects of smoking. *Chest* 131: 1557–1566.
45. McCarey DW, McInnes IB, Madhok R, Hampson R, Scherbakov O, et al. (2004) Trial of Atorvastatin in Rheumatoid Arthritis (TARA): double-blind, randomised placebo-controlled trial. *Lancet* 19: 2015–2021.
46. Ray KK, Cannon CP (2005) The potential relevance of the multiple lipid-independent (pleiotropic) effects of statins in the management of acute coronary syndromes. *J Am Coll Cardiol* 46: 1425–1433.
47. Rezaie-Majd A, Maca T, Bucek RA, Valent P, Muller MR, et al. (2002) Simvastatin reduces expression of cytokines interleukin-6, interleukin-8, and monocyte chemoattractant protein-1 in circulating monocytes from hypercholesterolemic patients. *Arterioscler Thromb Vasc Biol* 22: 1194–1199.
48. Smith GD, Ebrahim S (2004) Mendelian randomization: prospects, potentials, and limitations. *Int J Epidemiol* 33: 30–42.
49. Sie MP, Sayed-Tabatabaei FA, Oei HH, Uitterlinden AG, Pols HA, et al. (2006) Interleukin 6 –174 g/c promoter polymorphism and risk of coronary heart disease: results from the rotterdam study and a meta-analysis. *Arterioscler Thromb Vasc Biol* 26: 212–217.
50. Maini RN, Taylor PC, Szechinski J, Pavelka K, Broll J, et al. (2006) Double-blind randomized controlled clinical trial of the interleukin-6 receptor antagonist, tocilizumab, in European patients with rheumatoid arthritis who had an incomplete response to methotrexate. *Arthritis Rheum* 54: 2817–2829.
51. Frost C, White IR (2005) The effect of measurement error in risk factors that change over time in cohort studies: do simple methods overcorrect for ‘regression dilution’? *Int J Epidemiol* 34: 1359–1368.
52. Boshuizen HC, Lanti M, Menotti A, Moschandreas J, Tolonen H, et al. (2007) Effects of past and recent blood pressure and cholesterol level on coronary heart disease and stroke mortality, accounting for measurement error. *Am J Epidemiol* 165: 398–409.

Editors’ Summary

Background. Coronary heart disease (CHD), the leading cause of death among adults in developed countries, kills one person in the US every minute. With age, “atherosclerotic plaques”—deposits of fats, calcium, and various cellular waste products—coat the walls of arteries, causing them to narrow and harden, interrupting blood flow through the body. When this occurs in the coronary arteries, which nourish the heart muscle, the end result is CHD. If a plaque breaks off the artery wall, it can get trapped in the arteries and completely stop the blood flow, causing death of the heart muscle. The technical term for this is “myocardial infarction” (MI), although it is more commonly known as a heart attack. Smoking, high blood pressure, high blood levels of cholesterol (a type of fat), being overweight, and being physically inactive all increase the risk of developing CHD, as do some inherited factors. Treatments for CHD include lifestyle changes (for example, losing weight and exercising regularly) and medications that lower blood pressure and blood cholesterol. In the worst cases, the narrowed artery can be widened using a device called a stent or surgically bypassed.

Why Was This Study Done? Atherosclerosis might, at least partly, be an inflammatory condition. Inflammation—an immune response to injury characterized by swelling and redness—involves the production of proteins called “cytokines,” which attract cells of the immune system to the site of injury. In atherosclerosis, damage to the artery walls seems to trigger inflammation, which helps the atherosclerotic plaques grow. Because of the potential involvement of inflammation in atherosclerosis, increased levels of circulating cytokines might be associated with an increased risk of CHD. If they are, cytokines might provide a new therapeutic target for the treatment of CHD. In this study, the researchers have asked whether prolonged moderate increases in the cytokine interleukin-6 (IL-6) in the bloodstream are associated with CHD risk. IL-6, which is produced very early in inflammation, survives only briefly in the human body and its levels fluctuate within individuals. Consequently, its relevance to CHD has been unclear in previous studies.

What Did the Researchers Do and Find? Between 1967 and 1991, nearly 25,000 healthy, mainly middle-aged people were enrolled into two studies—the Reykjavik Study and the British Regional Heart Study—and followed for about 20 years, during which time 2,138 people had a first-ever nonfatal heart attack or died of CHD. The researchers measured baseline IL-6 blood levels in these participants and in 4,267 similar

participants who had not had a CHD event. They also measured IL-6 levels in 558 healthy participants several years into the study to determine a “regression dilution ratio” for IL-6. This ratio gives an idea of the year-to-year consistency of IL-6 levels. When the researchers used this ratio to estimate the impact of prolonged increases in IL-6 levels on CHD, they found that increased long-term IL-6 levels more than doubled the risk for CHD in their study populations. The researchers then combined these new results with those of 15 previous relevant studies. This combined analysis indicated very similar findings to those in the new data.

What Do These Findings Mean? These findings indicate prolonged moderate increases in IL-6 levels are associated with risk of CHD as strongly as several major established risk factors, including blood pressure and blood cholesterol levels, but whether there is a cause-and-effect relationship remains unknown. More studies are needed to find out whether this result is generalisable to other populations, but the broad agreement between the Icelandic and British studies suggests that they should be. This study renews interest in IL-6-mediated inflammatory pathways and CHD.

Additional Information. Please access these Web sites via the online version of this summary at <http://dx.doi.org/10.1371/journal.pmed.0050078>.

- Read a related *PLoS Medicine* Perspective article
- The MedlinePlus encyclopedia has pages on coronary heart disease and atherosclerosis (in English and Spanish)
- Information is available from the US National Heart Lung and Blood Institute on coronary heart disease and atherosclerosis
- Information for patients and caregivers is provided by the American Heart Association on all aspects of heart disease, including inflammation and heart disease
- Information is available from the British Heart Foundation on heart disease and on keeping the heart healthy
- Further details are available about the Reykjavik Study and the British Regional Heart Study
- Wikipedia has pages on inflammation and on interleukin-6 (note that Wikipedia is a free online encyclopedia that anyone can edit; available in several languages)